



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

PAGE'S
INTRODUCTORY TEXT BOOK
OF
PHYSICAL GEOGRAPHY
2/.



TEXT-BOOKS

ON

GEOLOGY AND PHYSICAL GEOGRAPHY.

By DAVID PAGE, F.G.S.

Fifth Edition, price 1s. 9d.

INTRODUCTORY TEXT-BOOK OF GEOLOGY. With Illustrations and Glossarial Index.

"Of late it has not often been our good fortune to examine a text-book on science of which we could express an opinion so entirely favourable as we are enabled to do of Mr Page's little work."—*Athenæum*.

Third Edition, Revised and Enlarged, price 6s.

ADVANCED TEXT-BOOK OF GEOLOGY, Descriptive and Industrial. With numerous Illustrations and Glossary of Terms.

"An admirable book on Geology. It is from no invidious desire to underrate other works—it is the simple expression of justice—which causes us to assign to Mr Page's 'Advanced Text-Book' the very first place among geological works addressed to students, at least among those which have come before us. We have read every word of it with care and with delight, never hesitating as to its meaning, never detecting the omission of anything needful in a popular and succinct exposition of a rich and varied subject. The *ordonnance* of its material is clear, masterly, and philosophical. The exposition is often eloquent, without ever striving after rhetorical effect. The information is lucidly yet briefly given. . . . The three passages (quoted) sufficiently indicate the quality of the book; its merits as a text-book can only be estimated by the student himself. If it does not drive many a student, hammer in hand, into quarries and railway cuttings for immediate experience of geological phenomena, nothing will."—*Leader*.

"It is therefore with unfeigned pleasure that we record our appreciation of his 'Advanced Text-Book of Geology.' We have carefully read this truly satisfactory book, and do not hesitate to say that it is an excellent compendium of the great facts of Geology, and written in a truthful and philosophic spirit."—*Edinburgh Philosophical Journal*.

"We know of no introduction containing a larger amount of information in the same space, and which we could more cordially recommend to the geological student."—*Athenæum*.

Price Sixpence.

THE GEOLOGICAL EXAMINATOR. A Progressive Series of Questions adapted to the Introductory and Advanced Text-Books of Geology. Prepared to assist Teachers in framing their Examinations, and Students in testing their own Progress and Proficiency.

In Crown Octavo, price 6s.

HANDBOOK OF GEOLOGICAL TERMS AND GEOLOGY.

"Every science, like every manual art, must have its own technicalities and modes of expression: new objects require new names, and new facts new phrases to express their relations. There is no avoiding this necessity in any progressive branch of human knowledge; and the only thing that can be done to lessen the difficulty—next to the rigid exclusion of whatever seems superfluous—is to explain these terms in brief and simple language. This I have endeavoured to do, chiefly with a view to the requirements of the general reader, at the same time appending such details as might render the volume an acceptable Handbook of Reference to the student and professed Geologist. Thus the ordinary reader will generally find the information he requires in the first and second sentences of a definition; what follows is addressed more especially to the professional inquirer—to the student, miner, engineer, architect, agriculturist, and others, who may have occasion to deal with geological facts, and yet who might not be inclined to turn up half-a-dozen volumes, or go through a course of geological readings, for an explanation of the term in question."—*Author's Preface.*

In Crown Octavo, price 6s.

THE PAST AND PRESENT LIFE OF THE GLOBE.

Being a Sketch in Outline of the World's Life-System. With Fifty Illustrations, Drawn and Engraved expressly for the Work.

"Mr Page, whose admirable text-books of geology have already secured him a position of importance in the scientific world, will add considerably to his reputation by the present sketch, as he modestly terms it, of the Life-System, or gradual evolution of the vitality, of our globe. In no manual that we are aware of have the facts and phenomena of biology been presented in at once so systematic and succinct a form, the successive manifestations of life on the earth set forth in so clear an order, or traced so vividly from the earliest organisms deep-buried in its stratified crust, to the familiar forms that now adorn and people its surface. Without wearying the reader with an endless profusion of technical details and scientific nomenclature, such as might be proper to a dry handbook of paleontology, or diving at length into specific distinctions or physiological riddles, it has been the writer's aim to submit, in clear readable form, a trustworthy *résumé* of the science, brought down to the latest point of discovery, and treated from a higher and more comprehensive point of view than that of either mere geology or natural history taken by itself."—*Literary Gazette.*

"Mr Page is favourably known as the author of three or four useful elementary geological books, each of which we have noticed in due order of appearance. Together with the one now published, they form a serviceable series for primary instruction. The present volume, however, has somewhat higher pretensions than its predecessors, and is more readable continuously than any one of them. . . . On the Development Hypotheses Mr Page has condensed some sound and unanswerable argumentative information."—*Athenæum.*

"The object of this excellent and unpretending little work is clearly set forth in the title. It contains a pictorial and historical sketch of the life-scenes of our earth, from the earliest times until now, as inferred from the geological records. It is perspicuous in style, prettily illustrated, furnishing a very complete compendium of the subject. The spirit of the book is good. Mr Page is evidently a philosopher who thinks it no disgrace to be a Christian—a man of science who believes in an Author of science. He is no 'developmentist'; and his arguments concerning successions of life are singularly acute and conclusive. Some of his positions are strikingly original; as, for instance, his views on recurrent cycles of temperature. He holds creation to be still in progress, presiding over the constant introduction of new species. And as each advance in creation has been, on the whole, upward, so our author anticipates a still higher order of animated creatures on our earth, perhaps also a higher destiny for man. We strongly recommend this little book to our readers, and its calm and temperate spirit to many of our modern writers."—*British Quarterly Review.*

In the Press, by the same Author.

I.


INTRODUCTORY TEXT-BOOK OF PHYSICAL GEOGRAPHY. With Illustrations and Glossarial Index. Price 1s. 9d.

II.

ADVANCED TEXT-BOOK OF PHYSICAL GEOGRAPHY. With Illustrations and Glossary of Terms.

W. BLACKWOOD & SONS, EDINBURGH & LONDON.





INTRODUCTORY TEXT-BOOK
OF
PHYSICAL GEOGRAPHY

BY
DAVID PAGE, F.R.S.E. F.G.S.
AUTHOR OF "INTRODUCTORY AND ADVANCED TEXT-BOOKS OF GEOLOGY," "HANDBOOK
OF GEOLOGICAL TERMS AND GEOLOGY," "PAST AND PRESENT
LIFE OF THE GLOBE," ETC.

WILLIAM BLACKWOOD & SONS
EDINBURGH AND LONDON
MDCCCLXIII

~~180. c. 191.~~
201. f. 56.

The Right of Translation is reserved.

"If, preserving the etymological sense of the word *Geography*, we should, with many authors, undertake to limit this study to a simple description of the surface of the globe, and of the beings that are found there, we must at once renounce the idea of calling it by the name of *science*, in the exalted sense of this word. To describe without rising to the causes or descending to the consequences, is no more science than merely and simply to relate a fact of which one has been a witness. The geographer who thus understands his study, 'seems to make as little of geography as the chronicler of history.'"—Guvor's *Earth and Man*.



P R E F A C E.

THIS little treatise, like the Author's 'Introductory Text-Book of Geology,' is intended to convey in a simple but systematic manner the leading facts of the science to which it relates. The learner is not troubled with masses of details which it is impossible for him to remember, nor is he deterred from the study of a most interesting and important branch of knowledge by an array of astronomical, geological, and meteorological data, which, while they bear but distantly on his subject, are altogether unneeded at this stage of his progress. A great deal of valuable information may be acquired respecting the external conditions of our globe—its lands and waters, their extent and configuration, their climate and vegetable and animal productions—without burdening the mind with unimportant distinctions, on the one hand, or going, on the other, into the higher reasonings on which the phenomena of Physical Geography depend ; and it is the object of this initiatory volume to facilitate the acquisition of such knowledge, and prepare the learner, should he wish to proceed beyond this stage, for the study of the subject in its more exact and extended aspects. It should be remembered, however, that even in its simplest form the

science of Physical Geography is by no means a child's study, and that the comprehension of its truths will be greatly assisted by a previous acquaintance with the elements of General Geography. Under this conviction the Author has endeavoured to produce a *Text*- not a *Task*-book—a thing to be read and reasoned, and not a compilation of unconnected facts to be irksomely and uselessly committed to memory.

While the text has been prepared on a general plan, and capable of being studied in connection with any modern set of physical maps, the order has, in most instances, special reference to Keith Johnston's 'School Atlas of Physical Geography,' as yet by far the best and most available of similar publications.

GILMORE PLACE, EDINBURGH,
December 1862.

CONTENTS.

	PAGE
I. PHYSICAL GEOGRAPHY AS A SCIENCE.	
Its Aim and Objects—Theoretical and Practical Importance—Hints to the Learner,	9
II. THE EARTH—ITS GENERAL OR PLANETARY RELATIONS.	
Its Figure, Motions, and Dimensions—Density, Temperature, and Atmosphere—Technical Subdivisions, as Points, Circles, Zones, &c.,	14
III. THE EARTH—ITS INDIVIDUAL STRUCTURE AND COMPOSITION.	
The Rocky Crust—Its Constitution and Formation—Relative Age and Arrangement of Rock-Formations—Connection between Physical Geography and Geology, . . .	26
IV. DISTRIBUTION OF LAND AND WATER.	
Their Relative Positions and Areas—Their Subdivisions, Extent, and Nomenclature—Action and Reaction on each other,	35
V. THE LAND—ITS SUPERFICIAL CONFIGURATION.	
Contour or Horizontal Disposition—Vertical Relief or Elevation—Causes and Consequences of Configuration, . . .	44
VI. THE LAND—ITS HIGHLANDS.	
Mountains, their Characteristics and Origin—Mountain Chains and Systems—Table-Lands or Plateaux, . . .	52

	PAGE.
VII. THE LAND—ITS LOWLANDS.	
Plains and Deserts—Valleys and Minor Depressions,	70
VIII. THE WATER—ITS OCEANS AND SEAS.	
Their Areas and Configuration—Composition, Density, Depth, and Temperature—Waves, Tides, and Currents,	79
IX. THE WATER—ITS RIVERS AND LAKES.	
Springs and Streams—Rivers and River Systems—Lakes and Lacustrine Areas,	106
X. CLIMATOLOGY—THE ATMOSPHERE.	
Nature and Constitution of the Atmosphere—Its Heat and Moisture—Winds, constant, periodical and variable— Dew, Rain, Hail, Snow, &c.—Causes affecting Climate,	134
XI. PLANTS AND ANIMALS—THEIR DISTRIBUTION.	
Life as affected by External Conditions—Plant Distribu- tion—Animal Distribution—Interdependence of Plants and Animals,	162
XII. ETHNOLOGY—RACES AND VARIETIES OF MAN.	
Man as affected by External Conditions—Characteristics and Distribution of Races—Conditions of Civilisation and Progress,	176
XIII. GENERAL REVIEW AND DEDUCTIONS,	186
GLOSSARIAL INDEX,	190

PHYSICAL GEOGRAPHY.

I.

PHYSICAL GEOGRAPHY AS A SCIENCE.

Its Aim and Objects.

1. GEOGRAPHY (from two Greek words—*gê*, the earth, and *graphê*, a writing or description) embraces, in its widest sense, all that can be known of the superficial aspects of our globe—its lands and waters, their extent and configuration, their altitude and depth, the atmosphere that surrounds them, their varied conditions and climate, and, finally, the distribution of the plants and animals by which they are respectively peopled.

2. A science embracing so wide and varied a subject will readily present itself under several heads or departments; and thus we may have what is termed *Mathematical Geography*, which devotes itself to the size, form, motions, and general divisions of the earth as a planet forming part of the solar system; *Political Geography*, which relates to the arbitrary subdivision of the earth into empires, kingdoms, and states, with their populations, manners, religion, laws, industry, commerce, and other features distinctive of such subdivisions; *Descriptive or General Geography*, which restricts itself to a mere account of the external conditions of the lands and waters—their extent and configuration, their scenery, life, and other obvious features—without inquiring into the causes that produce these appearances; and, lastly, we may have *Physical Geography*, which, while it embraces all the natural conditions of the lands and waters depicted by descriptive geography, proceeds further to inquire into the causes that produce these results, and by which they are in the course of nature continually reproduced within certain limits of change and modification.

3. Physical Geography is thus the higher department, its aim being not only to describe the external aspects of the terraqueous (land and water) globe, but to inquire into those conditions of position, altitude, soil, heat, moisture, and the like, which govern the distribution of plants and animals on the land, and those conditions of depth, composition, and temperature which regulate, in a similar manner, the distribution of plants and animals in the ocean. The Land, its continents and islands, its mountains and valleys, its soil and climate, its scenery and life-arrangements—the Ocean, its seas and bays, its shoals and depths, its composition and temperature, its aspects and life-dispersion—are the special objects of Physical Geography, and its highest aim the discovery and expression of those laws which regulate the action and reaction of land, water, and atmosphere in the production of physical phenomena. Physical Geography thus rises above the mere description of external appearances, and seeks to explain the causes that produce them—arranging the whole into a system of world-machinery, whose modes of action can be understood, and whose results it is possible to determine.

4. Rising to this conception of his science, and viewing the beautiful and diversified field before him, the student of geography meets a problem in every phenomenon that presents itself, and finds a solution in every incident that occurs. Why, for instance, do two countries lying within the same parallels of latitude present such differences in climate? Why are mountain-heights perpetually enveloped in snow, while the lower ridges are clothed in verdure and blossom? Why does one region of a continent be arid and rainless, while another is deluged with periodical torrents? Why do the winds in certain latitudes blow steadily for weeks in one direction, while in another they are fitful and irregular? Why should one expanse of ocean be still and tideless, while another swells and falls with tides, and is traversed by currents? Why should the plants and animals that flourish in one region dwarf and die out if transferred to another that seems equally fertile in soil and genial in climate? Why should the men at the mountain-foot be tillers of fields and dressers of vineyards, while those a thousand feet higher are herdsmen and shepherds? Or why should one country be the scene of busy industry and successful commerce, of intellectual activity and mental culture, while another, as fair and even more fertile, remains the mere squatting-grounds of indolent, dependent, and semi-civilised hordes? These and a thousand similar questions press themselves upon the attention of every geographical observer; and while the facts may be detailed with clearness and accuracy by Descriptive Geography, to Physical Geography, in particular, we must turn for a rational solution of the phenomena presented.

5. In the prosecution of his subject, the student of Physical Geography appeals to Astronomy for what relates to the figure,

size, motions, and other primary conditions of our planet; to Geology for the structure and constitution of the rocky crust which forms, as it were, the groundwork of all geography; to Meteorology for much that belongs to climate and its allied phenomena; while from Chemistry and Physiology he derives important aid in dealing with the nature, growth, and dispersion of plants and animals. Though drawing in this manner from other sciences, it by no means follows that the student should be deeply read in Astronomy, Geology, or Meteorology. All that is necessary is, that he be able to perceive the connection and interbearings of these sciences, and be capable of appreciating the importance of their deductions as far as they relate to his own immediate study of Physical Geography.

Theoretical and Practical Importance.

6. The value of such a science must be obvious to the most casual observer. To determine the relative extent of the land and water that constitute this terraqueous surface—the varying altitudes of the one and the depths of the other, the climates of the one and the winds and currents that traverse the other, with the infinitely diversified mineral, vegetable, and animal productions of both—is not only a source of high intellectual enjoyment and culture, but a task of prime industrial necessity. Scattered over the earth's surface, separated by sea and mountain, enjoying different climates, and placed in proximity to different mineral, vegetable, and animal products, it is a natural necessity that different nations should trade and barter with each other. To ascertain the peculiarities of this varied surface, to learn the variety of its products—to know all, in fine, that relates to the home we tenant, and the comforts and necessities with which it is furnished, as well as the obstacles or facilities that lie in the way of obtaining them—is the sum and substance of Geography. The observation and reasoning required in geographical research, the amount of information obtained, and the curiosity gratified by faithful descriptions of distant and diverse regions, constitute on the one hand its theoretical value; acquaintance with their mineral, vegetable, and animal products, now so indispensable to civilised existence, the conditions under which these occur, and the capability of the latter for naturalisation in other countries, form, on the other hand, its economical or practical importance. To the navigator dependent on the winds and currents of the ocean; to the pioneer and settler in new lands; to the merchant-traveller in search of additional products; to the gardener and farmer who seek to naturalise the plants and animals of different regions; and even to the physiologist and psychologist who would study the influence of climate and other conditions on health as well as on mental peculiarity, Physical Geography becomes a

science of direct and important interest. Combining, therefore, its theoretical with its practical bearings, our science has paramount claims alike on the attention of the philosopher, the statesman, the sailor, the farmer, the merchant, and manufacturer.

Hints to the Learner.

7. To present an outline of this science intelligible to beginners, and preparatory to a more exact and extended course of study, is the object of the present treatise ; and the learner will best accomplish his purpose who step by step confirms the statements of the text by reference to the maps of his atlas. By this double process any little difficulty will be more readily mastered, while the facts will be fixed more clearly and permanently on the memory. Every principle thoroughly understood becomes a key to some other problem ; and hence the value of systematic treatment and the necessity of a clear comprehension of the successive stages by which the higher problems are attained. The study of General Geography—the positions of towns, courses of rivers, heights of mountains, and the like—is for the most part little better than a task of memory ; Physical Geography, on the other hand, requires reasoning at every step, and the student will find his reasonings greatly assisted, not only by the systematic use of his atlas, but by the habit of appealing to the phenomena presented by his own immediate district. There are few localities, however limited, that do not present their alternations of hill and dale, of lake and river, of warm winds and cold winds, of periods of drought and periods of rainfall, of plants that love the marsh and others that thrive only in the thirsty upland ; and by noting such distinctions, and the causes concerned in their production, the mind, by such training, will be better prepared for the comprehension of the phenomena of wider and more varied regions. As the botanist, zoologist, and geologist find the objects of their studies in every walk through the fields around them, so the student of Physical Geography will find the illustrations of his science in every locality he may visit. Every district has its own features of highland and lowland, its streams and lakes and rivers, its peculiar winds and rains and frosts, its special arrangements of plants and animals ; and he who understands best the governing causes of these local peculiarities will be best able to deal with the general problems of Physical Geography. There is nothing fortuitous in the economy of our planet : every breeze that blows, every cloud that sweeps across the firmament, and every shower that falls—fickle, uncertain, and local as these may appear—are as much the results of law and law-directed forces as the rising and falling of the tides, or the revolutions of the planets. Impressed with this conviction, and seeing how closely every incident in nature is connected with

another, the student will consider no occurrence as too trivial, and no fact as too insignificant, to deserve his attention.

RECAPITULATION.

In the preceding paragraphs we have endeavoured to show that the object of Physical Geography is to describe the external conditions of the globe, and discover the causes by which they are maintained or modified. It treats of the Earth's surface as composed of land and water, determines their extent and configuration, their altitude and depth, the climate and other conditions that influence the growth and distribution of the plants and animals by which they are respectively peopled, and, rising above these, endeavours to account for the mental and social peculiarities of different nations as seemingly dependent on external circumstances. Deriving from Astronomy what relates to the figure, dimensions, motions, and other primary features of our planet; from Geology the structure and composition of its rocky crust; from Meteorology the proximate causes of the diversity of climates; and from Chemistry and Physiology the more intimate nature of vegetable and animal life, Physical Geography proceeds to apply these adjuncts to its own proper field of inquiry, and from the whole deduces a rational and connected account of the cosmical phenomena by which we are surrounded. As a science of observation and deduction in connection with the external conditions of the beautiful planet we inhabit, it possesses high intellectual attractions; and as bearing on its mineral, vegetable, and animal products—their abundance, distribution, and capabilities of dispersion—it becomes to civilised nations a study of prime industrial importance.

II.

THE EARTH—ITS GENERAL OR PLANETARY RELATIONS.

Figure, Motions, Dimensions.

8. By the general relations of the Earth are meant those primary conditions of form, size, density, motion, and the like, which belong to it as a member of the planetary system. From these conditions arise all those multifarious actions and reactions that take place on its surface—the alternations of night and day, heat and cold, summer and winter, growth and decay, the winds and motions of the atmosphere, the tides and currents of the ocean, and in fact all that confers on it geographical diversity and change. Thus, on its axial rotation depends the recurrence of light and darkness ; from its revolution round the sun arises the succession of the seasons, with all their varied effects on vegetable and animal life ; from the unequal reception of the sun's heat by the terraqueous surface and atmosphere, result the various phenomena of what we call weather and climate ; while from the attraction of the sun and moon, and the earth's own proper motions, spring the flow and ebb of the tides, and the other great currents of the ocean. The consideration of these conditions belongs, no doubt, more especially to Astronomy and Physics, but as much may be here recapitulated as will enable the student to lay the foundation, as it were, of his own special science.

9. Astronomers have determined that the Earth we inhabit is one of a number of planets that revolve at different distances, and with different velocities, round the Sun as a common centre—constituting what is termed the *solar system* (Lat. *sol*, the sun). These bodies, some of which are nearer the sun than our earth, and others more remote, some vastly larger, and others smaller, are nearly all spherical in form, and move from west to east, in courses or *orbits* more or less circular. As at present known, this solar system consists of the great centre or *sun* ; nine large or primary *planets*—Vulcan, Mercury, Venus, Earth, Mars, Jupiter, Saturn, Uranus, and Neptune ; seventy-one *planetoids*, or small planet-like bodies, between the orbits of Mars and Jupiter, and occupying, as it were, the place of a large primary ; twenty secondaries or *satellites*, revolving round their primaries ; and an unknown


number of *comets*, which move round the sun in extremely elliptical orbits, and, consequently, become visible to us only at widely-distant periods. As the secondary planets revolve round their primaries, and these, again, round the sun, so the solar system itself may revolve round some vaster centre, and this order of things through systems and centres that baffle the grasp of our finite conceptions. The following table exhibits at a glance the names of the planets, their diameters, distances from the sun, times of revolution and rotation, and number of satellites :—

Names and order of planets.	Diameters in English miles.	Distance from Sun in miles.	Revolution in days.	Rotation in days and hours.	No. of satellites.
SUN, . . .	883,000	D. H. M. 25 0 0	...
Vulcan, . . .	785	18,500,000	19.70
Mercury, . . .	3,140	37,000,000	87.97	1 0 5	0
Venus, . . .	7,700	69,000,000	224.70	0 23 21	0
Earth, . . .	7,918	95,000,000	365.24	1 0 0	1
Mars, . . .	4,100	144,000,000	686.98	1 0 39	0
<i>Planetoids</i> ,	263,000,000	1,684.74	...	0
Jupiter, . . .	90,000	494,000,000	4,332.62	0 9 56	4
Saturn, . . .	76,068	906,000,000	10,759.30	0 10 29	8
Uranus, . . .	34,500	1,822,000,000	30,686.82	0 9 30	6
Neptune, . . .	42,000	2,869,000,000	60,624.63	...	1

10. As stated above, the distance of the Earth from its central orb is 95,173,000 miles—or, in round numbers, ninety-five millions of miles. Its time of revolution round the sun is about $365\frac{1}{4}$ days, or more precisely, 365 days, 5 hours, 48 minutes, and 50 seconds; and this period of revolution we designate *a year*. Besides this annual revolution round the sun, the earth rotates or turns on its own axis in 24 hours—or, more exactly, in 23 hours, 56 minutes, and 4 seconds; and this period of rotation constitutes *a day*. In other words, the earth turns on its own axis three hundred and sixty-five times during the course of a single revolution round the sun; and thus the more frequent and obvious motion of daily rotation has been taken as the unit of measurement for the longer and less apparent motion of annual revolution. In these movements of annual revolution and daily rotation, the earth is attended by a minor or secondary body, which revolves round her as she round the great central luminary of the system. This secondary planet or *satellite* (Lat. *satelles*, an attendant) is the Moon, which is 2160 miles in diameter, and which completes her revolution round her parent orb (at a distance of 240,000 miles) in 27 days, 8 hours—or, in round numbers, in 28 days, or one lunar *month*. These motions, and times of motion, are determined and influenced by the attraction and gravitation

of the sun and other planets ; and it is by the same great forces that all the heavenly bodies that lie beyond our system are held in harmonious order, and in all likelihood in similar systems or fraternities.

11. The sun being the great centre of light, heat, and other ethereal influences, it necessarily follows that, during the earth's rotation on her axis, only one half of her surface can be exposed at a time to these influences ; hence the alternations of *day* and *night*, and all the phenomena that accompany these alternations. But day and night are of unequal and varying lengths at certain localities, according to the *seasons*, and these seasonal successions are caused by the facts, *first*, that the orbit or path of the earth's revolution round the sun is not a perfect circle, but an ellipse or oval ; and, *second*, that in performing this revolution, her axis is not perpendicular, but inclined or slanting at an angle of $66\frac{1}{2}$ degrees to the plane of her orbit. The accompanying diagram will assist in explaining the consequences of this elliptical orbit, and obliquity of axis. Thus, *s* is the sun, with the earth represented at four different points in her annual orbit. At *A* and *B* the light and heat of the sun strike at the equator or middle line, and consequently day and night are of equal duration. At any intermediate position, day and



The Seasons.

night are respectively lengthened and shortened : when at *c* the north pole is in darkness ; and when at *D* the south pole is in the same state. When the point presented to the sun is at *e* (which is on the 22d December), it is midsummer to all the southern parts of the earth, and winter to all the north ; but as the exposed part advances towards the point *f*, the northern regions gradually receive more and more heat, till on the 21st of June it becomes their midsummer. Such are the main systematical motions of the earth that regulate our days, years, months, and

seasons; let us now advert to her own proper dimensions, as determined by the astronomer and mathematician.

12. The body which thus rotates on its own axis, while it revolves round the sun, and is in turn revolved around by the moon, is not, strictly speaking, a sphere or globe. A diameter taken along its axis, or the ideal line round which it rotates, is only 7899.170 miles, while one taken in the opposite direction is 7925.648 miles. The one diameter thus exceeds the other by about $26\frac{1}{2}$ miles, thereby causing a deviation from the true globular form, and producing what is termed an *oblate spheroid* (Gr. *sphaira*, a sphere; *eidos*, like); that is, a figure flattened in the direction of its axis, and bulging out all around somewhat in the shape of an orange. Such a figure can be produced by rapidly spinning a ball of yielding material (like soft clay or putty) round its own axis, when the tendency which all rotating bodies have to fly off from the centre (centrifugal force) causes the mass to bulge out at the circumference, and to flatten in proportion at the ends or *poles* of the axis. To this centrifugal force arising from rotation, and to some original yielding condition of the earth's mass, is its oblate or spheroidal form usually ascribed. The earth's mass, as is well known, is kept together by the force of *gravitation*; and had it remained at rest, its form would have been perfectly spherical; but the moment it began to turn on its own axis, the particles of its mass began to obey another law—viz., that of *centrifugal force*, which exerts itself at right angles to the axis of rotation, and in proportion to the distance from that axis. Hence the greater bulging-out of the earth's mass at the equator, where the distance from the axis is greatest; and hence also the gradual declension of centrifugal force as we proceed towards the poles.

13. It has been proved, we have said, by the astronomer and mathematician, that the true figure of the earth is that of an *oblate spheroid*, or ellipsoid of revolution; but of its *general spherical or globular form*, the ordinary observer has many evidences, among which may be mentioned the following:—1. As a vessel sails away from the land we first lose sight of her hull, next of her lower or mainsails, and lastly of her topsails and pennants, thus clearly showing that she is passing over a spherical surface. 2. The reverse of this also holds true; for the mariner as he approaches the land first sees the mountain-tops; and on gradually nearing it, the lower grounds, stage by stage, make their appearance. 3. In travelling any considerable distance, north or south, new stars gradually come into view in the direction to which the traveller is advancing, while others disappear in the direction from which he is receding. 4. Many navigators, by constantly sailing in one direction, or nearly so, whether due east or due west, have returned to the port from which they set out, thus making what is termed the *circumnavigation* of the globe. 5. The shadow which the earth casts on the moon during an eclipse is always circular. And, lastly, the earth belonging to

a system or brotherhood, the other members of which are globular, the fair presumption is that she also is of the same form.

14. Calculating from the dimensions given in the preceding paragraphs, the *mean diameter* of the earth will be 7912.402 miles, and its *mean circumference* or girth, 24,858 miles. It is usual, however, to speak, in round numbers, of the diameter being 8000, the radius or distance from the surface to the centre as 4000, and the circumference as 25,000 miles. The superficial area of a globe of these dimensions amounts to 197 millions of square miles; and of these about 51 millions consist of land, and 146 millions are occupied by water. The proportion of land to that of water may, therefore, be said to be, in round numbers, as *one to three*; or, in other words, while one-fourth of the earth's surface consists of dry land, the other three-fourths are covered by the waters of the ocean. The solid contents of the mass have in like manner been computed to exceed 260 thousand millions of cubic miles—an amount which, though expressible in figures, is altogether beyond the grasp of human conception. It is enough, then, for the learner to be convinced of the magnitude of these amounts, without troubling himself with the remembrance of their numerical expressions.

Density, Temperature, Atmosphere.

15. The density of the globe, as compared with the materials at its surface, has been determined with considerable precision. The average or mean density of the most prevalent rocks is about $2\frac{1}{2}$ times that of water; while the density of the whole mass, as determined by astronomical experiment, is $5\frac{1}{2}$ times that of water. It is evident from this that the interior of the earth cannot be composed of the same materials that constitute its outer portion; for these, under the law of gravitation, would be so compressed at the depth of a few miles as to give a greater mean density to the whole mass than its astronomical relations will allow. Thus, it has been calculated that water, at the depth of 362 miles, would be as dense as quicksilver; and that the density of marble, at the centre of the earth, would be 119 times greater than what it is at the surface. Either, then, the interior of the earth is composed of materials differing altogether in nature from those known at its surface, or the compression of gravitation must be counteracted by some highly expansible force, such as heat, so as to maintain the mean density which astronomy and physics have determined. It is thus that geologists, and geographers, adopting the language of geologists, speak of the "crust of the globe," and of the "interior of the globe"—meaning by the former that outer portion composed of rocks and rock-masses which can be observed and examined, and by the latter that inner portion which is placed beyond such investigation, and respecting which we can offer little more than conjecture.

16. Closely connected with the density of the globe is its temperature, or the amount of heat that pervades it. As one of the orbs of the solar system, the earth has a variable superficial or atmospheric temperature; it has also a temperature peculiar to the rocky crust; and, judging from volcanic action, there is also a high and more remarkable interior or central temperature. Respecting the *superficial temperature*, which constitutes the great theme of Climatology, to be noticed hereafter, it may be stated, in the mean time, that it is influenced from day to day, and from season to season, by the heat of the sun; that it varies according to the latitude, being greatest at the equator and gradually decreasing towards the poles; that it is greatly modified by the extent and distribution of sea and land—the sea and sea-coasts being more equable than inland continents, which experience extremes of heat in summer and extremes of cold during winter; that it is also modified by the absorbent or radiating nature of the soil, according as this may be dark or light coloured, dry or moist, porous or compact; and, lastly, that it is notably affected by elevation above the mean level of the sea, the higher being the colder regions. The *temperature of the accessible crust*, on the other hand, is affected either by the direct heat of the sun, by heat generated chemically among its own materials, or by heat derived by conduction from the interior. During summer, for instance, the earth is warmed to a certain depth by the heat of the sun; during winter this heat is given off to the surrounding atmosphere; and though the heat of one summer and the cold of one winter may differ from the heat and cold of others, still, on an average of seasons, the results are pretty equable. It may, therefore, be stated in general terms, that in summer the crust of the earth at small depths is cooler than at the surface; that during winter the crust at these depths is warmer than at the surface: but that, at the depth of 80 or 90 feet, the variations of summer and winter become wholly insensible. Respecting the *heat of the interior*, we see it abundantly manifested in hot springs, volcanoes, and the like; and have, by direct experiment, been enabled to arrive at some important facts relative to its descending rate of increase. Thus, it has been found by experiments in coal-pits, artesian wells, and metalliferous veins, that after passing the depth at which the surface heat becomes inappreciable, the temperature begins to rise, and this gradually and steadily for every fathom of subsequent descent. On an average this rise takes place at the rate of 1° of Fahrenheit for every 55 or 60 feet of descent; and, calculating at this increase, a temperature (2400° Fahr.) would be reached at a depth of 25 miles or thereby, sufficient to keep in fusion such rocks as basalt, greenstone, and porphyry. We know little, however, of the deportment of heat under such an enormous pressure as must exist at these depths, and can only indicate the line of reasoning that leads to the general belief that the solid or rocky crust forms but a comparatively thin film, and that the great interior mass exists in a

state of high incandescence or molten fluidity. Intense as this interior heat may be, and active as it undoubtedly is in the production of volcanic phenomena, the surface temperature of the globe is scarcely if at all affected by it (according to Fourier only $\frac{1}{17}$ of a degree), owing to the weak conducting properties of the rocky crust.


17. Besides these original conditions of form, size, motions, density, and temperature, there is also that of its *atmosphere*, or aerial envelope which surrounds it on every side, and becomes an integral and indispensable portion of its constitution. This atmosphere, as is well known, is mainly composed of two gases—nitrogen and oxygen—79 parts of the former to 21 of the latter, with a small percentage of carbonic acid and other extraneous impurities. As at present constituted, the air is indispensable to vegetable and animal life; both alike breathe it, and by this breathing the functions of vitality in both are alike sustained. Being an elastic or compressible medium, the air nearest the sea-level is denser than that at considerable elevations; and by calculating the rate at which this rarity takes place, it has been estimated that at the height of 45 miles above the sea the atmosphere becomes so rare or light as to be inappreciable. We have thus surrounding the earth, and partaking in all its movements, a gaseous envelope or atmosphere, whose pressure at the sea-level is estimated at $14\frac{1}{2}$ lb. avoirdupois on every square inch of surface, but which gradually becomes lighter and lighter or more attenuated as we ascend, till at the height of 45 or 50 miles its presence is inappreciable. Through this envelope the heat and light of the sun are equally diffused and modified; it is also the recipient and diffuser of all watery vapours arising from the earth; and from local alterations in its density or expansibility, caused by heat and the like, arise all aerial currents or winds—whether steady and regular, or fitful and irregular. The atmosphere is thus the great medium through which light, heat, moisture, and other vital influences are communicated to the plants and animals that adorn the earth's surface; it is also the great laboratory in which all meteorological and electrical phenomena are elaborated; and hence all the varied aspects and results of winds, clouds, rains, snow, hail, thunderstorms, and the like, that constitute the essentials of climatic diversity.

Technical subdivisions—Points, Circles, Zones.

18. In treating of the earth thus constituted and surrounded, geographers make use of certain terms and technicalities expressive of distance, position, and the like, on its surface, and with these it will be necessary for the learner at this stage to render himself familiar. Thus, the direction from which the earth moves in its daily rotation is called the *West*; that towards which it

moves the *East*; the point to which a spectator looks, with the east on his right hand and the west on his left, is called the *North*; and that behind him the *South*. In this rotation from west to east, any point on the earth's surface turns towards the sun in the morning, and away from him after mid-day, and thus the terms *Sunrise* and *Sunset* (the sun being a stationary centre) express not real but apparent phenomena. The imaginary line on which the earth rotates is termed its *axis*, and the terminations of this axis its *poles*—that towards the north being known as the *North Pole*, and that towards the south as the *South Pole*. From an idea of the earlier geographers, who were inhabitants of the northern hemisphere, that the north pole was uppermost (though in reality there can be neither upper nor under in a globe freely rotating in space), it is customary in our maps and charts to place the Northern part at the top and the Southern at the bottom, with the east on the right hand and the west on the left. A line distant alike from either pole, and dividing the earth into two equal portions (Lat. *æquus*, equal), is called the *Equator*; the northern being known as the *Northern hemisphere*, and the southern as the *Southern hemisphere*. In the same way we may speak of the *Eastern hemisphere* and *Western hemisphere*, by supposing the globe divided by a line passing through the poles, and thus necessarily cutting the equator at right angles. Reckoning London as a fixed point on such a line, Europe, Asia, and Africa will lie in the eastern hemisphere, and North and South America in the western. (See Map, page 37.)

19. Like other circles, the circumference of the earth is divisible into 360 equal parts or *degrees*, and in this case each degree will equal about $69\frac{1}{2}$ British statute miles. At the distance of $23\frac{1}{2}$ degrees north and south of the equator (as shown in the accompanying diagram), are two parallel lines called the *Tropics* (Gr. *trōpē*, a turning), from their marking the turning-points of the sun in the ecliptic; that on the north being known as the *Tropic of Cancer*, and that on the south the *Tropic of Capricorn*, because these constellations occupy a corresponding point in the heavens. At the same distance ($23\frac{1}{2}$ degrees) from either pole is a parallel line—that on the north being called the *Arctic Circle* (from the northern




constellation, *Arctos*, the Bear), and that on the south the *Ant-arctic Circle*, because opposite (*anti*, opposite) to that of the north. The space or belt between the Tropics is called the *Torrid Zone*, because the sun, being always vertical in some part of that space, produces a greater degree of heat than in regions where his rays strike more obliquely. The spaces between the tropics and the arctic and antarctic circles, on either side, are termed the *Temperate Zones* (north and south); and those between these circles and their respective poles, the *Frigid Zones*. It is customary for the botanist and zoologist to subdivide these zones more minutely into equatorial, tropical, subtropical, warm-temperate, cold-temperate, subarctic, arctic, and polar; but of these subdivisions and their characters when we come to treat of the distribution of vegetable and animal life.

20. The equator has been described as a circle exactly between the poles, and cutting the globe into two equal portions: a line which cuts the equator obliquely (at an angle of $23\frac{1}{2}^{\circ}$), and touching upon opposite sides of the tropics, is called the *Ecliptic* (Gr. *ekleipsis*, an eclipse). The ecliptic is the orbit described apparently by the sun round the earth, but in reality by the earth round the sun; and is named from the circumstance that all eclipses can happen only when the moon is in the same plane or very near it. The points where the ecliptic cuts the equator are called the *equinoctial points* or *nodes*, because when the sun is in these parts of his course, the day and night are of equal duration. These equinoxes take place twice a-year—namely, on the 21st March and 21st September. The term *Greater Circles* is sometimes applied to the equator and ecliptic, because they encircle the earth at its thickest part—that is, cut it at planes passing through the centre; and that of *Lesser Circles* to the tropics, arctic, and other circles above described. Besides these circles, another set of lines, drawn from pole to pole over the earth's surface, and cutting the equator at right angles, are termed *Meridians* (Lat. *meridies*, mid-day), because, when any of these lines is opposite the sun, it is mid-day or 12 o'clock with all the places situated along that meridian on the same side of the globe; while on the other side it is midnight. When it is noon, for instance, in any particular part of Britain, it is midnight at an opposite and corresponding part near New Zealand; and so on for the intermediate hours at directly opposite parts along the same meridian. Thus, diametrically opposite parts on the globe are said to be the *Antipodes* of each other (Gr. *anti*, opposite, and *pous*, *podos*, foot), because the feet of their inhabitants are placed, as it were, in opposition to each other.

21. The position of a place on the earth's surface is determined by what is termed its *latitude* and *longitude*,—terms which arose from a notion of the ancients that the earth was longer from east to west than from north to south—or, in other words, had length (*longitudo*) and breadth (*latitudo*). The latitude of a place is, therefore, its distance north or south from the equator; hence, we

speak of north latitude (N. Lat.) in the northern hemisphere, and of south latitude (S. Lat.) in the southern hemisphere. Longitude, on the other hand, is measured east and west of any fixed meridian—different countries adopting different meridians (usually that of their capitals), and Britain that of the Royal Observatory at Greenwich; hence English geographers speak of east longitude (E. Long.), and west longitude (W. Long.), according as a place may be situated eastward or westward of the meridian of Greenwich. The circles on the earth being divided into 360 degrees, and each degree, again, into 60 minutes, each minute into 60 seconds, and so on, the position of a place can be indicated with great precision; and for the sake of brevity certain signs are employed, as $4^{\circ} 6' 12''$, meaning thereby 4 degrees, 6 minutes, and 12 seconds. The distance between the equator and either pole being only the fourth part of the earth's circumference, the latitude of a place north or south of the equator can never exceed the fourth part of 360, or 90 degrees. Longitude, on the other hand, being measured east and west of a fixed meridian, embraces a whole hemisphere, or 180 degrees; and as the meridians all converge towards the poles, the degrees of longitude become less and less as we approach these extremities. The lines of latitude being necessarily parallel to each other, geographers speak of *parallels of latitude*—designating those that lie near to the equator as *low* latitudes, those that approach the poles as *high* latitudes, and those that intervene as *middle* latitudes.



Parallels and Meridians.

RECAPITULATION.

In the foregoing chapter we have noticed those primary conditions of form, motion, size, density, temperature, and the like, which belong to the earth as a member of the solar system. Those conditions lie, as it were, at the foundation of all change and diversity on the earth's surface—day and night, heat and cold, summer and winter, meteorological fluctuation and climate, oceanic tides and currents, geological waste and reconstruction, and, in fine, all that confers diversity on land and water, and consequently on vegetable and animal life, being directly or

remotely dependent on the original constitution of the globe, and its connection with the solar system. It is necessary, then, that the student should render himself familiar with the planetary relations of the earth—its spheroidal form, its daily and annual revolutions, its density, temperature, magnitude, and admeasurements, as well as with those terms and technicalities by which its several portions and positions are known and described. As an oblate spheroid, then, the earth has an equatorial diameter of 7925.648, and a polar diameter of 7899.170 miles; rotates on its own axis in 23 hours, 56 minutes, and 4 seconds, or in *one day*; revolves round the sun in an elliptical path or orbit in 365 days, 5 hours, 48 minutes, and 49 seconds, or *one year*; and is in turn attended by the moon, which performs her revolution in 27 days 8 hours, or *one lunar month*. As the mean circumference of the earth is 24,858 miles, and as she rotates on her axis once in 24 hours, it is clear that any spot on the equator must be borne round at the rate of more than 1000 miles an hour. With every successive removal from the equator, however, the circles of latitude become smaller and smaller, and as they are all carried round within the same period of 24 hours, their rate of motion must be proportionally less. At the 60th parallel, for example, this motion is only 500 miles an hour, and at the absolute poles it ceases to exist. The mean density of the rocks known at the surface of the earth is about $2\frac{1}{2}$ times that of water, while that of the entire mass is $5\frac{1}{4}$, thereby showing that the interior consists of different material, or of materials in a different condition from those in its rocky crust. The earth's internal temperature, as distinct from the climatic temperature received from the sun, increases according to the depth; and this increasing temperature, though little affecting the surface, seems sufficient to keep the interior in a state of fusion, and to be the cause of volcanoes, hot springs, and other thermal phenomena. The earth, besides her own proper mass, is also surrounded by an aerial envelope or atmosphere, consisting in the main of 79 parts nitrogen and 21 oxygen, with a fractional percentage of carbonic acid; and this gaseous envelope, being an elastic medium, is denser at the earth's surface (where it presses with a weight of $14\frac{1}{2}$ lb. on the square inch), and becomes rarer and rarer as we ascend in space, till at the height of 45 or 50 miles its presence becomes inappreciable. This atmosphere is the medium through which the sun's light and heat are equally diffused; the laboratory in which all meteorological phenomena—winds, rains, clouds, storms, &c.—are elaborated; and, in fine, the source whose varied

mutations are the proximate causes of all climatic diversity. In treating of the earth, various terms and technicalities are necessarily employed by geographers, and with these the learner should early render himself familiar. Thus, the imaginary line on which the earth turns in her daily rotation is termed the *axis*; the extremities of this axis, north and south, the *poles*; the circle between the poles, which cuts the globe into two equal portions, the *equator*; these two equal portions are termed *hemispheres* or *half-spheres*; and the varying belts of surface temperature as we proceed northward and southward from the equator, where the sun's heat is greatest, are respectively known as the *torrid*, *temperate*, and *frigid zones*. The *ecliptic* is the apparent path of the sun round the earth; *meridians* are those lines drawn from pole to pole over the earth's surface, and at right angles to the equator, along which the sun is vertical at his highest daily ascension, or *noon*; *latitude*, the distance of any place, measured in degrees, south or north of the equator; and *longitude*, the distance of the same place east or west from any arbitrary or convenient meridian,—that in Britain being the meridian of Greenwich—that in France, Paris—and that in Germany, the Faroe Islands.

III.

THE EARTH—ITS INDIVIDUAL STRUCTURE AND COMPOSITION.

The Rocky Crust—Its Constitution and Formation.

22. HAVING considered, in the preceding chapter, some of the more obvious relations of the globe as a member of the Solar System, and the terms usually employed to express their connection, we now proceed to describe the leading features of its own individual structure. As the knowledge of its general relations was derived chiefly from Astronomy, so a knowledge of its individual structure is mainly obtained from the teachings of Geology. As the one set of facts could be taught without going deeply into the problems of the astronomer, so the other may be understood without entering largely into the reasonings of the geologist. What, for instance, is the nature of the earth's rocky crust? How are its rocks arranged, and how does this arrangement affect its superficial character? What has stretched out the level plain and upheaved the rugged mountain? What renders one soil obnoxious and sterile, and another genial and fertile? And as the earth's crust is continually undergoing modification under the operation of external and internal forces, what the effects of such modifications on the general geography of the globe? These and similar problems Geology endeavours to solve, and, with a little explanation, these solutions may be rendered intelligible to the student of Physical Geography.

23. Geologists speak of the "crust of the globe" just as the housewife talks of the "crust of her loaf." The crust of the loaf is one thing, the inside of it another. The crust or exterior portion of the earth, composed of rocks and rock-materials that can be seen and handled, is one thing—the interior, of which we can know nothing by direct observation, is a very different thing. By observation and comparison we can determine a great many truths respecting the structure and composition of the one; respecting the other, we can offer at best little more than conjecture. This outer portion, or rocky crust, is the great theatre of all geographical phenomena—the foundation of the land and

waters—the arena of climatic influences—the field of vital development—and, as such, it behoves the geographer to know something of its history and constitution.

24. On the most cursory inspection of quarries, railway-cuttings, sea-cliffs, and ravines, the observer will find a great portion of the rocks arranged in layers, or lying one above another in beds or strata (Lat. *stratum*, spread out). These are said to be *stratified*, and generally consist of sandstones, clays, shales or consolidated muds, limestones, and other similar rocks. He will also find another set not spread out in layers, but rising hard and massive in no determinate arrangement. These are termed the *unstratified*, and consist of such rocks as the granites, greenstones, basalts, and lavas. How, he will naturally inquire, have two sets



A A A, Stratified; B B, Unstratified Rocks.

of rocks, so dissimilar in character and arrangement, been produced? and as we can only interpret nature's productions by a knowledge of nature's operations, we must seek for the answer to this question in what is now taking place around us. And, first, if we turn to any lake, estuary, or bay of the ocean, we will find that the mud, sand, and gravel carried down by the rivers, or washed from the cliffs by the waves and tides, are deposited and arranged along the bottom of these receptacles in layers or strata more or less horizontal, and in course of ages, one above another, precisely like the shales and sandstones of the quarries and sea-cliffs. Here, then, we are entitled to infer that rocks arranged in layers or strata have been formed through and by the agency of water (Lat. *aqua*)—that is, have been deposited as sediment (Lat. *sedere*, to settle down) in water, or brought together and assorted by the action of moving water; and hence they are termed *aqueous*, *sedimentary*, or *stratified*. If we turn, in the next place, to the volcano or burning mountain, we find that lava and other molten rock-matters are discharged from its crater, and these, when cooled and consolidated, age after age, form mountain-masses, and fill up chasms and rents, precisely as the granites and greenstones and basalts do among the stratified rocks with which they are associated. Here, again, we are entitled to infer that greenstones, basalts, and other similar rocks, are the products of fire (Lat. *ignis*, fire); and hence they are designated *igneous*, *eruptive*, or *unstratified*.

25. In the crust of the earth, then, we have two main sets of rocks, the stratified and the unstratified—the one formed through and by the agency of water, the other through and by the agency of fire. The former are chiefly the products of aqueous and

atmospheric waste, the latter the products of igneous reconstruction; and between these two forces, the aqueous and igneous, the crust of the earth has ever been held in habitable equilibrium. Were the aqueous and atmospheric forces to operate uncontrolled, all the higher portions of the dry land would in the course of ages be worn down, and the whole reduced to a dreary uniformity of level; but to prevent such a contingency, the volcanic forces are perpetually exerting themselves from below, and once more upheaving the crust into dry land and diversity of surface. The great design of creation seems clearly to be diversity in time as well as diversity in space. Whether under the open air or under the waters of the ocean, hill and dale, level plain and rugged mountain, are ever attended by diversity of soil, diversity of winds and currents, heat and cold, and all the other climatic influences on which diversity of plant-life and diversity of animal-life are so intimately dependent. To the primary geological oscillations of the earth's crust—its waste and reconstruction—its submergences and upheavals—are we, therefore, indebted for all that confers on its surface geographical variety and diversity.

Relative Age and Arrangement of Rock-Formations.


26. If, then, the crust of the globe be in a state of oscillation between the aqueous and atmospheric forces that waste and wear from without, and the igneous forces that reconstruct and upheave from within—if cliffs and hills are worn down, and lakes and estuaries filled up and converted into alluvial plains—if plains are thrown up into mountains and the sea-bed into dry land—if large tracts of the earth are gradually raised higher and higher above the ocean while other regions are gradually submerged—it is clear that different portions of the rocky crust must be of different ages and composed of different materials. The present distribution of sea and land has undergone many noted modifications, even within the historic era; and if such changes have been accomplished within a period so brief as a few thousand years, what may we believe to have taken place during the thousands of ages that preceded human history? So numerous and repeated have been these changes, as clearly shown by geology, that every portion of the existing land has been at one time or another beneath the waters; and that which now constitutes the bed of the ocean has, in like manner, been the dry land of former epochs. The record of these changes lies in the stratified rocks, each period producing its own sedimentary deposits, and these deposits constituting what geologists term a *formation*, as having been formed at a certain period, and under the peculiar conditions of that period. Further, as the sediment of existing lakes

and estuaries and seas imbed the remains of plants and animals that have lived in these areas, or have been drifted from the land—leaves, branches, trunks, bones, teeth, shells, and crusts ; so the sediments of former ages contain the remains of plants and animals that then existed, these remains being petrified or converted into stone. These petrifications, or *fossils* as they are termed, differ widely in many instances from the plants and animals now peopling the globe ; and this difference may be said to increase with the depth and age of the strata that contain them. As a general rule, the older strata will be the deepest seated—the older, harder and more crystalline than the recent and superficial ; and the older the strata the greater will be the difference between their fossil plants and animals and the plants and animals now existing. Founding on principles such as these, geologists have arranged the stratified rocks into formations and systems according to their relative antiquities—each division representing (as nearly as can be determined) a different arrangement of sea and land, and a different aspect of vegetable and animal life during the period of its deposition. Thus, beginning at the recent and superficial, which contain few extinct forms of life, and descending through systems whose fossils belong chiefly to extinct races, we have :—

<i>Systems.</i>	<i>Periods.</i>
1. POST-TERTIARY, or RECENT ACCUMULATIONS, 2. TERTIARY SYSTEM,	} CAINOZOIC (Recent Life).
3. CRETACEOUS, or CHALK SYSTEM, 4. OOLITIC, or JURASSIC SYSTEM, 5. TRIASSIC, or UPPER NEW RED SANDSTONE,	
6. PERMIAN, or LOWER NEW RED SANDSTONE, 7. CARBONIFEROUS SYSTEM, 8. DEVONIAN, or OLD RED SANDSTONE, 9. SILURIAN and CAMBRIAN,	} MESOZOIC (Middle Life).
10. METAMORPHIC, or CRYSTALLINE STRATA,	} PALÆOZOIC (Ancient Life).
	} HYPOZOIC (Beneath Life).

Or arranging them in diagrammatic form, one above another, as they occur in the British Islands, we have the intelligible sequence represented on the following page.

27. As the stratified rocks have been arranged in chronological sequence, so also the unstratified or igneous admit of a similar though less precise arrangement. As at present, so during every former epoch of the earth's history, the volcanic forces have been in a state of activity—breaking up the stratified crust, throwing out lava and other products in the formation of hills, and filling up earthquake fissures with molten rock-matter from below. In course of the earth's mutations, these



CHRONOLOGICAL SUCCESSION OF BRITISH STRATIFIED SYSTEMS.


earlier products become submerged beneath the waters of the ocean and overlaid by newer sedimentary deposits, again to be broken through by another set of igneous operations, and this consecutively and continuously from the most distant ages up to the present day. As a general rule, the earliest igneous rocks will be associated with the earliest stratified deposits, and the older and deeper-seated will be more uniform and crystalline in texture than the newer and more superficial. Founding on principles such as these, geologists usually arrange the igneous or unstratified rocks in the following chronological order:—

1. Volcanic, associated chiefly with tertiary and upper secondary strata.
2. Trappean, associated chiefly with secondary and upper primary strata.
3. Granitic, associated chiefly with primary and metamorphic strata.

Or arranging them diagrammatically as they occur among the stratified systems, we have something like the connection represented on page 31.

28. The consideration of these systems and subdivisions forms the special subject of Geology; but as each formation occupies a distinct portion of the earth's crust, consists of different kinds of rocks, and presents different soils and surfaces, the following summary may be of use to the student of geography. The *Post-Tertiary* consists for the most part of sands, gravels,

peat-mosses, alluvial silt, and other superficial accumulations that have been recently formed or are still in the course of formation. From the nature of their formation, they occupy the flats or lower portions of plains, river-valleys, and deltas, and as a consequence present little variety of surface or scenery. The *Tertiary System* consists, in most areas, of clays, sands, gravels, marls, and occasional limestones; are in general superficial and spread over plains and lowlands, and are rarely elevated into hills of any magnitude. Subjected to the long-continued influences of elevation and meteoric waste, the system presents greater variety of surface than the post-tertiary; and the districts round London, or the "London Basin," as it is termed, the "Hampshire Basin," and "Paris Basin," may be taken as average types of its scenery. The *Cretaceous System* is composed of sands and sandstones, clays, shales, and, notably in many areas, of thick beds of chalk or earthy limestone. Subjected to atmospheric and other waste, this group is often worn into rounded hills and valleys, as is typically illustrated by the "downs" and "coombs" of the south-eastern counties of England. The *Oolitic System*, as developed in Europe, consists of clays, shales, sandstones, and limestones; the calcareous members in general predominating. From the harder nature of the calcareous strata, they have resisted the effects of external waste better than the softer clays and shales; and hence the undulating and pleasing character of much of the oolitic landscape. The *New Red Sandstone*, which comprises the *Triassic* and *Permian Groups*, consists, as its name imports, chiefly of sandstones, sandy marls, and limestones. The sandstones and marls being of a soft and disintegrating nature, have largely yielded to ex-



ternal influences ; and hence much of that red tenacious clay and moist pasture-land which characterises so large a portion of new red sandstone districts. The *Carboniferous System*, so called from its yielding in Britain our main supply of coal, is composed of sandstones, shales, fire-clays, ironstones, limestones, and coals ; and, unless where cut through by hills of greenstone and basalt, is generally overlaid by thick tenacious clays, which confer on its landscape a cold, moorland, and uninviting appearance. The *Old Red Sandstone* consists mainly of reddish-coloured sandstones and conglomerates, with subordinate beds of limestone ; and, being frequently upheaved into hills, and cut through by ravines and valleys, presents greater diversity of surface and more amenity of landscape. The *Silurian System*, consisting of flaggy slates, schists, and limestones, and often upborne at high angles by the older granitic hills, presents a bolder landscape than the old red, but less abrupt and wild than the *metamorphic* and *crystalline schists*, whose splintery peaks and crags constitute so prominent a feature in all highland and mountainous scenery.

29. Such are the leading features of the stratified systems, and by such, with a little practice, they may be readily detected in the various areas they occupy. As with the stratified so with the unstratified ; each great group—Volcanic, Trappean, and Granitic—has its own peculiar and distinctive features. The *Volcanic*, as already stated, are generally associated with the more recent formations, and consist of trachytes or greystones, various lavas, scorïæ, and similar products. They rise up in dry rocky hills more or less conical and crateriform ; and these are perhaps more frequently grouped round some common centre than arranged in linear directions. The *Trappean*, so termed from the terraced aspect of many of the hills they compose, consist of greenstones, basalts, felstones, porphyries, trap-tuffs, and other kindred rocks, and are generally associated with the secondary and upper primary strata. They are usually arranged in hill-ranges more or less persistent, and, from their higher antiquity and longer subjection to wasting influences, are more worn into rounded heights, exposed crags, slopes, and terraces, which confer on the landscape a beauty and diversity peculiarly their own. The *Granitic*, or oldest series of igneous rocks, consists of granites, syenites, porphyries, serpentines, and the like, which, from their more ancient and deep-seated relations, are generally hard and crystalline in texture and massive in structure. They constitute the nucleus or backbone, as it were, of all the higher and older mountain-chains, elevating the metamorphic schists into splintery peaks and abrupt ridges, or presenting of themselves broad massive shoulders of cold, sterile moorland and unprofitable heath. Vesuvius, Etna, and the hills of Auvergne may be taken as typical examples of the Volcanic ; the Ochils and Sidlaws of the Trappean ; and the Grampians, the Wicklow, and Cornish mountains of the Granitic.

- Connection between Geology and Physical Geography.

30. Understanding the nature of the preceding subdivisions, the student will be able to attach some intelligible idea to such phrases as "the river cutting its way through secondary strata;" "lying in an irregular tertiary basin;" "crossing a plateau of triassic sandstones;" "a cold, retentive soil, derived from the subjacent carboniferous rocks;" "intersected by a low range of trap-hills, whose grassy slopes and terraces;" "bounded by sterile granitic ridges, whose snow-clad summits;" and hundreds of others that are of continual occurrence in geographical descriptions. He will still more fully perceive the intimate connection of Geology with his own immediate science, when he reflects that many of the soils which give character and colouring to vegetable growth are derived directly from the disintegration of the subjacent rocks, and are further affected in their fertility by the porous or retentive nature of the beds on which they rest. The soil derived from the Chalk, for instance, is light and absorbent; hence the short sweet herbage of the "downs" of the south of England; that derived from the marls of the New Red Sandstone is stiff, retentive, and less fitted for tillage than for pasture; the disintegration of trap-rocks, rich in soda, potash, and lime, yields a fertile, genial soil, which their structural fissures ever keep dry and pulverulent; while the scanty disintegration of an imperious granite is poor, cold, and barren. We have seen, moreover, that every geological formation is less or more characterised by its own special features. The soft rounded outline of a chalk range is altogether distinct from the slopes, terraces, and conical heights of secondary trap-hills; while these, on the other hand, are widely different, both in outline and in vegetable covering, from broad-shouldered mountains of granite, or the splintery peaks of metamorphic strata. In fine, these formations and rock-groups constitute the framework on which the geographical features are moulded; and their inherent characters, as subject to external and internal change, have produced, and are ever producing, new variety of superficial aspect. Geology and Geography are inseparably connected; and thus it is that some acquaintance with the nature and sequence of the rock-formations that constitute the solid crust, and with the causes that produce them, become so indispensable to the student of Physical Geography.

RECAPITULATION.

In the preceding chapter we have directed attention to the rocky structure of the globe as that which gives colour and character to all external phenomena, and is, therefore, of per-

manent importance to the student of Physical Geography. We have spoken of the "crust," composed of materials that can be seen and investigated, as distinct from the "interior," of which nothing can be known by direct observation. This crust, composed of stratified and unstratified, or of water-formed and fire-formed rocks, is held in habitable equilibrium between the disintegrating forces of water from without, and the reconstructing forces of fire from within. To these two opposing powers are chiefly owing the continuous geological modifications of the earth's crust—each modification representing a period during which certain rocks were formed, and the remains of plants and animals that then lived entombed in them in greater or less perfection. These formations being arranged by the geologist in chronological sequence, and each varying in mineral character, and consequently conferring on the landscape different aspects, it is of importance in Physical Geography to know the order of this arrangement, and the nature of these distinctions. These *formations*, we stated, had been arranged into certain *systems*, and these systems again grouped into certain *periods*, according to the leading features of their fossils, thus:—

1. Post-Tertiary.	}	CAINOZOIC.
2. Tertiary.		
3. Cretaceous.	}	MESOZOIC.
4. Oolitic.		
5. Triassic.		
6. Permian.	}	PALÆOZOIC.
7. Carboniferous.		
8. Devonian.		
9. Silurian.		
10. Metamorphic.	.	HYPOZOIC.

As with the stratified so with the unstratified—their arrangement into VOLCANIC, TRAPPEAN, and GRANITIC expressing the relative antiquities and nature of the great igneous groups that give character and individuality to the hills and mountain-ranges of the globe. So far, then, as diversity of surface, fertility of soil, character of landscape, and similar peculiarities, are concerned, the connection between Physical Geography and Geology is intimate and inseparable; and hence the necessity that the student of the one should be less or more acquainted with the leading facts of the other.

IV.

DISTRIBUTION OF LAND AND WATER.

Their Relative Positions and Areas.

31. HAVING noticed the general conditions of the Earth as a member of the solar system, and having also adverted to the structure of its own rocky crust as the groundwork of all geographical diversity, we now proceed to consider the more intimate relations of its surface as composed of Land and Water. When we speak of the earth's surface as composed of land and water, we mean that all the more elevated portions of the crust stand out as dry land, while the lower and more depressed are covered by the waters of the ocean. Geologically speaking, the relations of land and water are continually, though slowly, undergoing change and modification; geographically speaking, and within the limits of such mutation, what are the respective areas of land and water, their positions on the earth's surface, their actions and reactions on each other in consequence of their different extents, positions, capacities for heat, &c. ? and what the influence of these reactions on currents, winds, rains, and all those kindred phenomena that constitute climatic diversity ? The present distribution of land and water tends to the production of certain physical and vital results, and the explication of these results forms the sum and substance of our science ; but no portion of the present distribution could be altered either in its position, extent, or configuration, without being accompanied by a modification of these results, so intimately is the one element bound up with the other in the production of a definite and harmonious whole.

32. As formerly stated, the superficial area of the globe has been computed at 197,000,000 square miles. Of this amount about 51,500,000 consists of dry land, and about 145,500,000 of water. The proportion of land to that of water is, therefore, nearly as *one to three*; or, expressing it fractionally, while *one-fourth* of the earth's surface is composed of dry land, the other *three-fourths* are covered by the waters of the ocean. Considered in hemispheres, northern and southern, the proportions of land and water are nearly as follows :—


Northern Hemisphere,	Land,	38,000,000 square miles.
Do.	do.	Water, 60,500,000 "
Southern Hemisphere,	Land,	13,500,000 "
Do.	do.	Water, 85,000,000 "

The boundaries of this terraqueous arrangement (Lat. *terra*, land ; *agua*, water) are extremely irregular. Here the land spreads out in broad unbroken masses, there the waters stretch away in vast continuous expanses ; here the land rises up in mere specks and fragments, there the water inserts itself among them in areas equally limited and irregular. Here the land juts boldly out into the ocean, there the ocean runs sharply into the land ; here the land trends uninterruptedly for leagues, there the ocean embays itself with gentle curves into the bosom of the land. Although this distribution seems to obey no regular order, and has been continually changing throughout all geological time, yet we may rest assured—and not the less assured because we are unable to discover it—that there is a law governing the insensibly varying relations of sea and land, and defining their respective amounts. The boundary between the land and water is known as the *shore-line* or *coast-line*—the former having more especial reference to the margin washed by the waves, the latter to the terrestrial verge that opposes the ocean. The space alternately covered and laid bare by the tides is termed the *beach* or *strand*, and the fringe of land bordering on, and more or less influenced by, its proximity to the sea, is spoken of as the *sea-board* of a country or continent. This sea-belt is generally marked by the peculiarity of its plants and animals, as well as by the character of its inhabitants—these differing in many points from those of the interior or inland districts. In like manner, on the configuration of the shore-line depends much that relates to the climate as well as to the industry and commerce of a country. A regular and unbroken coast is generally an exposed and ungenial one, and one unfavourable to maritime enterprise ; while one indented with gulfs and creeks and bays has more of genial diversity, and is better fitted for the purposes of navigation and commerce.

33. A glance at the Map of the World will show that the land-masses are situated chiefly in the northern hemisphere—there being about three times as much land in the northern as there is in the southern hemisphere. And if we look at the distribution of this land as regards the zones of temperature, we will find that the greatest proportion is situated in the North Temperate Zone, or in that which contains all the better portions of Europe, Asia, and North America ; while only comparatively narrow or isolated tracts are cut directly by the equator or central Torrid line. Another glance at the map will show that, while the land lies in broad and unbroken masses towards the north, it stretches towards the south in curious gradually-tapering projections. Again, if the globe be divided into two hemispheres, as shown in the ac-

THE WORLD.

Western Hemisphere.



Eastern Hemisphere.



companying illustration—the one having London for its pole, and the other having that of Antipodes Island, near New Zealand—the former will be found to contain nearly all the land-masses, and may therefore be called the *Continental* or *Land Hemisphere*, while



Land and Water Hemispheres.

the other contains the greater proportion of the water-expanes, and may for that reason be termed the *Oceanic* or *Water Hemisphere*. These and other noticeable facts connected with the arrangement of sea and land have been adverted to by geographers; and though it must be confessed that many of their comparisons are more curious than suggestive, there can be no doubt that the present distribution is the result of some great cosmical law—a law concerning the nature of which neither Geology nor Geography can give any certain indication. This much, however, Geology has shown, that the present arrangement is not the arrangement that obtained in former ages, and that continents existed where seas now roll, and seas extended where continents are now established. We can readily account for the minor modifications of coast-lines—why some portions, composed of hard enduring rocks, should stand out in bold projections, while others, consisting of soft and wasting material, should be worn into bays and irregular recesses; but of the greater forces that slowly upheave some regions, and as gradually submerge others, we are altogether ignorant. But while we fail to account for the present distribution of sea and land, and while we find that it is continually undergoing modifications under the operation of geological forces, we know that these changes are so silent and gradual as scarcely for ages to interfere with our ideas of geographical permanence and stability.

Their Subdivisions, Extent, and Nomenclature.

34. Admitting this condition of the terraqueous surface, it will be seen, on further reference to the map, that the Land is broken up into two main masses,—that of the Eastern Hemisphere—embracing Europe, Asia, and Africa, and described, from its being the only portion known to the ancients, as the *Old World*,—and that of the Western Hemisphere, including North and South America, and known (from its comparatively recent discovery by Columbus in 1492) as the *New World*. Properly speaking, there are only two great continuous land-masses or *continents* (Lat. *con* and *tenens*, holding together)—that of the Old World and that of the New; but geographers usually speak of six continents or principal sections—viz., those of *Europe, Asia, Africa, North America, South America, and Australia*. Australia and its contiguous islands, together with those of the Southern Ocean, are frequently grouped as the sixth great division, under the title of *Oceania*; but of this and other minor distinctions when we come to treat of the continents in detail. The areas of these respective continents or *quarters*, as they are sometimes termed (though the “four quarters” of our forefathers embraced only Europe, Asia, Africa, and America) have been computed, with their respective islands, as follows:—

Old World, or Eastern Continent,.....	81,250,000 square miles.	
Europe,.....	3,725,000	”
Asia,.....	16,165,000	”
Africa,.....	11,360,000	”
New World, or Western Continent,.....	14,900,000	”
North America,.....	8,080,000	”
South America,.....	6,820,000	”
Oceania, or Maritime World,.....	4,200,000	”

The islands included in the above estimate are sometimes solitary and independent, as Iceland, Spitzbergen, Nova Zembla, Madagascar, New Zealand, &c.; are occasionally curiously connected with the extremity of some peninsula or continent, as Terra del Fuego, Sicily, Ceylon, Van Dieman's Land, &c.; but, on the whole, are most frequently found in clusters or *archipelagos* (a term originally applied to the isles of Greece), as the West Indian Islands, the Isles of Greece, the East India Islands, the Japan Isles, and many others that must at once present themselves on the most cursory inspection of the Map of the World.

35. In describing the features and peculiarities of the land, geographers make use of the following designations, which, though familiar in everyday language, may here, for the sake of method, be briefly recapitulated. Thus, a *continent*, as already indicated, is any extensive region uninterrupted by seas; an *island*, any

smaller portion surrounded by water ; a *peninsula* (Lat., almost an island), a portion nearly surrounded by water ; an *isthmus*, the narrow neck that connects two adjacent masses of land, or a peninsula, with the mainland ; and a *cape*, *promontory*, *head-land*, or *ness*, a point of land jutting out into the water. Besides these terms, which refer to the contour or disposition of the land as surrounded by water, there are others which relate to surface configuration or vertical relief. Thus, extensive flats are known as *plains*, *steppes*, *prairies*, *pampas*, and the like ; smaller flats, as *valleys*, *dales*, *straths*, *carsees*, &c. ; lands elevated more or less abruptly above the general surface are spoken of as rising into *hills*, or, still higher, into *mountains* ; hills and mountains may stand less or more apart from each other and be *isolated*—may occur in *groups*, as if connected with a common centre—or they may stretch away in determinate courses known as *chains* and *ranges* ; while level elevated tracts are spoken of as *table-lands* and *plateaux*.

36. Though encircling the globe on every side, and in all its parts most intimately connected with one great ocean, the Water is still divisible into certain areas that are more or less defined by the intervention of the land. Thus, on the west of the Old World, and between it and the New World, extends one main division known as the *Atlantic* (so called by the ancients from its washing the western base of Mount Atlas in Africa) ; on the west of the New World, and between it and the Old, expands another natural division, termed the *Pacific* (from its comparative freedom from storms) ; while between Australia and its contiguous islands on the east, and Africa on the west, lies the *Indian Ocean*. It is usual, however, to speak of *five* great oceans—viz., the *Atlantic*, *Pacific*, *Indian*, *Arctic*, and *Antarctic*—the two latter being respectively within the arctic and antarctic circles. In geographical descriptions it is also useful to employ the terms, "North and South Atlantic," and "North and South Pacific," and to speak of the expanse that stretches away in unbroken vastness between the Indian Ocean and the south pole as the "Great Southern Ocean." The areas of these respective expanses are usually stated as follows :—

Pacific Ocean,	50,000,000 square miles.
Atlantic Ocean,	25,000,000 "
Indian Ocean,	17,000,000 "
Arctic Ocean,	? "
Antarctic Ocean,	? "

Being ignorant of the amount of land that lies within the arctic and antarctic circles, and also of the exact extent of many islands both in the northern and southern oceans, the above must be held as approximations merely ; and further, as amounts that do not embrace the areas of *inland seas*—such as the German, Baltic, Mediterranean, Red, Black, Caspian, &c.—which occupy, as will

hereafter be seen, considerable and important spaces on the surface of the globe.

37. In treating of the waters of the globe, though there is, strictly speaking, only one great ocean, the term *ocean* is applied to the large uninterrupted expanses above defined ; smaller areas are known as *seas* ; gradual bendings of the water into the land as *bays* ; deeper indentations as *gulfs* ; minor and sudden indentations as *creeks*, *inlets*, *arms*, &c. ; the narrow belts or openings connecting two adjacent seas as *straits* or *channels* ; and where the sea stretches inland to receive the waters of a river, such an expanse is known as a *firth* or *estuary*. Besides these, there are other local and restricted terms—as the *fjords* or rocky inlets of Norway—the *lochs* or lake-like sea-arms of Scotland—the *lagoons* or shallow intercepted sheets that occur along the shores of the Adriatic and other seas ; but these will be best described under the respective areas to which they belong. Again, referring to the depth of the ocean, whose bed seems to be as irregular and varied as the surface of the dry land, geographers speak of *deeps* and *pits* ; of *shoals* and *banks* ; of *sounds* that may be readily reached by the sounding-line ; of *roads* and *roadsteads* for anchorage ; of *harbours* or landlocked inlets for shelter ; and similar terms, whose meanings are so obvious as to require no special definition.

38. Besides the great oceans and inland seas above alluded to, and which all consist of salt water, there are other considerable volumes that belong more especially to the land, and which consist mainly of fresh water—comparatively few being brackish or saline. These are the *springs* issuing from the earth's crust, and more or less impregnated with the mineral substances through which they have percolated ; the *streams* or runnels of water formed by the union of several springs ; the *rivers* formed by the union of streams, and often traversing whole continents with gradually-increasing volumes ; and the *lakes*, which occupy depressions in the land, and most frequently lie along the courses of rivers, though occasionally occurring isolated, and apparently fed by springs, or, if receiving the waters of a river, have no river of discharge, but give off their surplus water by evaporation. The amount of surface occupied by these inland waters it is impossible to estimate with anything like accuracy, but the main volumes, as will be seen when we come to treat of the "River-systems," are the North American lakes, the lakes of northern Europe and central Asia, and the greater rivers—as the St Lawrence, Mackenzie, and Mississippi in North America—the Amazon, Orinocco, and La Plata in South America—the Nile and Niger in Africa—the Ganges, Indus, Yang-tse-kiang, Lena, Yenesei, and Obi in Asia—and the Volga and Danube in Europe.

Action and Reaction of Land and Water.

39. Such is the relative distribution, and such the general features of the land and water that form the surface of the terraqueous globe. Whatever may have been their distribution in former ages as revealed by Geology, one thing is obvious, that in the present era they are, within certain limits of change, mutually adapted and harmoniously adjusted. The land absorbs and radiates the sun's heat more readily than the water, and thus, while parched and thirsty in summer, it is refreshed and vivified by the moisture evaporated from the more extensive ocean; while in winter its tendency to grow colder is modified by the heat given off by the ocean, whose slower radiation renders it, as it were, a great storehouse of latent heat for the exigencies of the land. Besides this interchange of heat and moisture, there is also the interchange of aerial currents and winds caused by the unequal capacity of the two great surfaces for heat, and thus, as will be more fully explained hereafter, the whole machinery of climate—hot winds, cold winds, vapours, rains, and the like—is set agoing by the primary differences existing between land and water. As at present distributed, a certain amount of moisture is evaporated from the ocean and carried to the land, a certain amount of heat is interchanged, certain winds blow at certain seasons, certain tides rise and fall, and certain currents flow in performance of certain functions; but all this would be changed by the slightest alteration either in the relative areas, in the relative configurations, or in the relative positions of land and water. In their present areas, configurations, and dispositions, the two elements are harmoniously co-adapted for the production of certain results, geological, climatological, and vital; and the student must readily perceive how the force of tides and waves and currents would be altered, the climate of regions changed, and their plants and animals affected by the slightest disturbance in the existing terraqueous arrangements. The terraqueous arrangements of time past, with all their accompanying phenomena, belong to Geology; the arrangements of the present constitute the sum and substance of Physical Geography.

RECAPITULATION.

The facts detailed in the preceding chapter are so simple and obvious, that they require little recapitulation. The surface of the globe, or that which comes in contact with the atmosphere, is partly occupied by land and partly by water—the former forming more than a fourth, and the latter about three-fourths of the entire area. On a cursory inspection of the map of the

world, the land will be seen to resolve itself into two main masses—the Eastern hemisphere, or Old World, and the Western hemisphere, or New World; and these by geographers are usually further subdivided into the so-called quarters or continents of Europe, Asia, and Africa in the former, North and South America in the latter, together with a south or insular division, embracing Australia, &c., under the name of Oceania. Of these continents the greater portion lies in the northern hemisphere, the broader masses spreading out towards the north, and gradually narrowing towards the south, in the cape-like projections of South America, Africa, Hindoostan, and the Malayan peninsula. The ocean, though encircling the globe on every side, and stretching from pole to pole, is also arranged in areas more or less defined by the intervention of the land-masses. The main divisions are the Atlantic, the Pacific, Indian, Arctic, and Antarctic Oceans, which for convenience of description are frequently subdivided into North and South Atlantic, North and South Pacific, Northern, Southern, and Polar Seas. The technicalities employed in treating of the relations of land and water—shore-line, coast-line, island, peninsula, isthmus, cape, bay, gulf, strait, channel, and the like—are so obvious, and of such frequent use in everyday language, as scarcely to require any special explanation. The physical results arising from the present distribution of sea and land are manifested chiefly in tides, currents, unequal reception and radiation of the sun's heat, and as a consequence winds, vapours, rains, and all the other phenomena that give rise to climatic diversity. What may be the cause of the present arrangement of sea and land—why the land should lie chiefly in the northern hemisphere, and largely within the temperate zone, while it disappears in a succession of narrow cape-like lobes towards the south—neither Geology nor Geography can determine. One thing, however, is certain, that it is part of a great geological sequence of continuous oscillations between sea and land, each change being attended by its own physical and vital phenomena; and that, as in the past history of the globe so in the present, no alteration in this terraqueous arrangement can take place without a corresponding alteration in all the essentials of Physical Geography.

V.

THE LAND—ITS SUPERFICIAL CONFIGURATION.

40. HAVING glanced at the distribution of land and water, and their general relations to each other as composing the terraqueous surface of the globe, we may now proceed to consider their special features,—the mountains, table-lands, plains, and valleys of the one—the composition, depth, temperature, tides, and currents of the other. And, first, in the present chapter, of the Land, whose features are more within our reach, and have been longer and more minutely the subjects of geographical investigation. How, for instance, is it disposed on the surface of the globe; and what the outline of its form as regards the ocean by which it is on every side surrounded? This, in geographical language, is its *contour*, or line of horizontal configuration. How, again, does it rise above the water; and what the surface inequalities presented to the atmosphere by which it is enveloped? This is its *vertical relief*—contour and relief embracing all that relates to the superficial configuration of the Land.

Contour or Horizontal Disposition.

41. Though Geology and Geography are alike unable to account for the present disposition of sea and land, there are still some features in the contours of the respective continents that demand a passing notice. Not that we can account for these features, nor that they are deserving the attention some geographers have bestowed upon them, but simply because they are arrangements productive of obvious results, and as such cannot be overlooked in our review of physical causation. Thus, though the greater bulk of the land lies in the northern hemisphere, the greatest extension of the Old World continent is from east to west, while that of the New World is from north to south. In this way the Old World, lying largely along the same zones, presents a greater uniformity of external conditions; while the New World, crossing the zones—frigid, temperate, and torrid—is subject to a greater diversity of

temperature, and, as a consequence, to all the conditions that arise from this diversity. 2. Both continents attain their greatest dimensions from east to west along the same parallel of latitude—namely, that of 50° north—a disposition that places much of North America, Europe, and Asia within the temperate zone, while only the narrower portions of South America, Africa, and the East India Islands lie directly under the burning heat of the equator. 3. Both continents, as formerly noticed, spread widely towards the north, where they nearly approach each other, and terminate broadly along the same parallel (72° N.), or nearly so; while towards the south they gradually grow narrower, and terminate in far-separated promontories. 4. In like manner, the direction of all the principal spurs and peninsulas in both continents (California, Florida, Greenland, Scandinavia, Spain, Greece, Hindoostan, Malay, &c.) is towards the south—a feature by far too general not to suggest to the inquiring mind a sameness in the producing cause. 5. These southward-trending spurs, moreover, are in most instances curiously accompanied by an outlying island or islands—as South America by Terra del Fuego and the Falkland Islands, Africa by Madagascar, Hindoostan by Ceylon, and Australia by Tasmania. 6. The general disposition of the continents and larger islands is in the direction of their principal mountain-axes; so that, given the direction of the mountain-chains, we know the longitudinal disposition of a continent; or, given the direction of a continent, we can foretell the strike of its mountains. 7. The almost complete separation of South America from North America, of Africa from the Old World, and the complete severance of Australia from Asia, as well as the curious resemblance that obtains between North America with the West India Islands on its south-east, Europe with the Grecian islands on the south-east, and Asia with the Indian Archipelago on the same quarter, are curious coincidences that have long been noticed by geographers. 8. The general tendency of islands to arrange themselves in clusters or archipelagos is a fact also suggestive of a common geological cause and origin. And, lastly, the numerous indentations of the sea that confer irregularity and extent of coast-line on the land in the northern hemisphere is a feature that strongly contrasts with the uniform and unbroken coasts of South America, Africa, and Australia in the southern hemisphere. Indeed, this last relation—viz., that of a broken and deeply-indented coast-line, furnished with peninsulas, gulfs, inland seas, and harbours—is one of the most important in Physical Geography, as on it depend greater diversity of climate and productions, and all those facilities for navigation and commerce which confer on nations their wealth, power, and independence. Europe and North America stand pre-eminent for their extent of coast-line, the former having one mile of coast for every 170 square miles of surface, and the latter one mile of coast for every 260 square miles; a proportion more than double, and

in some instances even four times, that of the other continents; hence one great physical reason for the maritime enterprise, commerce, industry, and civilisation of their inhabitants.

42. Besides the preceding analogies, there are several others that have been noticed by geographers; and though such comparisons are often more fanciful than real, they are all less or more suggestive, and may occasionally lead to satisfactory theory. But whatever may have been the causes that produced the present arrangement of the land-masses, no change could take place in their relative situations without being attended by corresponding changes in the nature of their climate and in the character of their vegetable and animal inhabitants. Had they lain chiefly within the tropics, been situated partly in the northern and partly in the southern hemisphere, with a broad belt of tropical ocean between, been arranged either longitudinally or latitudinally in parallel zones, or been broken up into smaller masses by the more frequent intervention of the ocean, a totally different set of climatic agents would have prevailed, and been consequently attended by a totally different distribution of plants and animals. As it is, and in the words of an eloquent authority, "each of these terrestrial masses, considered as a whole, as an individual, has a particular disposition of its parts, and the forms which belong to it—a situation relatively to the rays of the sun, and with respect to the seas or the neighbouring masses, which is not found identically repeated in any other. All these various causes excite and combine, in a manner infinitely varied, the play of the forces inherent in the matter which composes them, and secure to each of them a climate, a vegetation, and animal life; in a word, an assemblage of physical characters and functions which are peculiar to it, and which really give it something of individuality." It is this *individuality* of the land-masses, and the study of its character, which constitutes, in fact, the very purport and essence of Physical Geography.

Vertical Relief or Elevation.

43. Next to the *contour* or horizontal configuration of the land, the most important feature is its *vertical relief* or elevation above the level of the ocean. As the position of an island a few degrees north or south, its disposition along the parallels of latitude or along the meridians, or its configuration as broken up by seas or lying in one unbroken mass, must each of them influence the nature of its climate, so its vertical relief is productive of similar results—every rise of 300 or 350 feet being sufficient to alter the temperature by one degree of Fahrenheit. It is therefore possible to determine with satisfaction the cause of the climate of a particular spot.

existing position and contour of the continents, we know, at least, that the main instruments of their surface diversity into mountain and valley, table-land and plain, are the vulcanic forces acting from within. These, in obedience to some great law, have been continuously and gradually shifting place from the earliest traces of geologic history; at one time, and extensively, along the old granitic ranges of Europe, Asia, and Africa—at another, and with equal intensity, along the secondary hills of the same continents—and now, with apparently unabated vigour, along the main ridges of America, the islands that skirt the eastern shores of Asia, and the groups that stud the bosom of the Pacific. These successive shiftings and upheavals form, in fact, part and parcel of the great machinery of the globe—a machinery by which the sea-bed is upheaved into new land, and old surfaces submerged beneath the ocean—and this continuously, successively, and, in all likelihood, in an order which geology shall one day or other be enabled to determine. Besides the more obvious operations of the earthquake and volcano, there is also that slow but gradual elevation and depression of vast regions (*e.g.*, the uprise of northern Scandinavia, the depression of South Greenland, the uprise of the Arctic islands, &c.) which, in course of time, must effect most important changes; and to these combined forces, assisted by the wasting and wearing of air and water, are to be mainly ascribed the principal features in the vertical relief or superficial diversity of the Land.

44. Among these features may be noticed—*first*, that in all the continents there is a gradual rise from the sea-shore towards certain points or ridges in the interior—the spaces occupied by a few depressed areas, as the Caspian, &c., being too limited to affect the results of the general configuration. These ridges or points of maximum swell form the grand *watersheds* of their respective continents, and a glance at the courses of the principal rivers will at once direct the eye to the line of these mountainous elevations. 2. In all the continents this line of greatest elevation does not occupy the centre, but is placed more towards one side than another; hence arises two slopes of unequal length—the long side forming the *slope*, and the shorter the *counter-slope* of the continent. In the Old World the long slopes are turned towards the north, and the short towards the south; while in the New World the gentle slope is towards the east, and the short and rapid one towards the west. 3. Though this difference of arrangement in the slope and counter-slope takes place in the two worlds, there is still in both a general flattening of the land towards the north, and a general rise and irregularity of its surface towards the south—a disposition that tends to counteract the effects of the northern cold, and to temper the burning heats of the tropical regions. “If this order were reversed,” says M. Guyot, “and the elevations of the lands went on increasing to-

wards the north, the most civilised half of the globe, at the present day, would be a frozen and uninhabitable desert." 4. That the grand elevations or mountain-ranges lie in the direction of the greatest length of a continent ; in fact, it is the great lines or *axes* of upheaval that confer on the continents their main relief or configuration. Thus, in the New World, whose greatest length is from north to south, the ranges of the Andes and Rocky Mountains, the Cordilleras of Brazil, and the Alleghanies of the United States, stretch all more or less regularly in the same direction. In the Old World, on the other hand, whose greatest length is from east to west, the ranges of the Pyrenees, Alps, Carpathians, Caucasus, Himalayas, and Altai, lie chiefly in the same direction, throwing the great rivers to the north and south of these watersheds. The backbone, if we may so speak, of the New World, is the great mountain-chain that extends uninterruptedly from Patagonia on the south to Russian America on the north ; while the backbone or framework of the Old World is similarly composed of a series of ranges that extend with but little interruption from the Pyrenees on the west to Kamtschatka on the east. 5. As with the continents so with the principal islands and peninsulas, they are all less or more continuously traversed by mountain or hill ranges in the direction of their greatest length—as witness Scandinavia, Italy, Kamtschatka, Malaya, the islands of the Indian Archipelago, Madagascar, California, and Greenland.


45. Reserving the arrangement of the world's highlands into mountain-chains and systems for subsequent consideration, it may be noticed, in the mean time, that the loftiest ridges in the vertical relief of the land are those of the Himalaya in Central Asia, whose highest peak is, in round numbers, 29,000 feet ; next, those of the Andes, in South America, 24,000 feet ; the Cordilleras of Mexico, in North America, 17,740 feet ; the Alps, in Europe, 15,740 feet ; the Abyssinian Mountains, in Africa, 14,720 feet ; and the highest ascertained points in Malaya, Australia, and Polynesia respectively, 15,000 feet, 7,000 feet, and 16,000 feet. These altitudes, however, convey no idea of the general relief of the several continents, whose superficial aspects depend more on the table-lands and plains, the hill-ranges and valleys, than on the extreme heights of their principal mountains. Indeed, so small a proportion do these extreme ridges bear to the great mass of the continents, that it has been calculated by Humboldt that the Alps, spread equally over the surface of Europe, would raise the general level not more than 21 feet ; while the vast mountain-chains of Asia, treated in a similar manner, would not add more than 150 feet to the general elevation of that continent. Blending all their heights and hollows into a general average, the *mean elevations* of the respective continents above the sea-level have been calculated as follows :—

Europe, 1342 feet ; Asia, 2264 feet ; Africa, 1800 feet ; North America, 1496 feet ; and South America, 2302 feet ; or, taking the whole—continents and islands, mountains and plains—the mean elevation of all the land has been estimated at somewhat less than 1000 feet. As the surface of the dry land presents its plains and mountains, and the sea-bed its shoals and depths, so it is usually imagined that the depressions of the one sink in proportion to the elevations of the other. For this belief, however, there is no foundation in fact. On the contrary, judging from the greater extent of the ocean, the soundings we already possess, and the small mean elevation of the dry land, it seems more reasonable to conclude with Sir John Herschel, that “the mean height of the surface of the dry land most probably does not exceed one-fifteenth of the mean depth of the ocean.” At all events, in the absence of reliable soundings for extreme depths, and with so much of the ocean-bed unexplored and unknown, it is better to avoid such comparisons, and to deal with the vertical relief of the land on its own merits, and as a thing in a great measure already determined.

RECAPITULATION.

It will be seen, from the facts stated in the preceding chapter, that the three great elements in the configuration of the land are relative position, contour, and vertical relief. By *relative position* is meant the place an island or continent occupies on the surface of the globe ; and just as this position is frigid, temperate, or tropical—as it lies in broad parallelism along the same zone, or crosses meridianally two or more zones—so will its climate be genial or ungenial, uniform or diversified. The causes that have determined the position of the existing continents lie far as yet beyond the indications of science, though we may rest assured that in the gradual shiftings of volcanic energy from centre to centre, and in the alternate uprise and submergence of large tracts of the earth's crust, we behold the results of a definite and continuously operating law. By *contour* is meant the outline or figure which the land receives by being surrounded by the water ; and just as this outline is simple and uniform, or irregular and indented by seas, bays, and gulfs, so will its external conditions be varied, and its coasts better fitted for the purposes of navigation and commerce. This contour depends mainly, of course, upon the original inequalities of the crust, but partly also upon the waste and degradation which all coast-lines suffer from the ceaseless action of the ocean. Combined, as in the case of Europe,

with favourable relative position, it exercises most important influences on climate, productions, industry, enterprise, and civilisation. By *vertical relief* is meant the elevation of the land above the level of the ocean; and just as this is regular or irregular, low or lofty, so will the whole character of a country be determined. A few hundred feet more or less of elevation is sufficient to change the whole physical aspects of a country—converting arable fields and vineyards into pasture-lands, pasture-lands into pine forests, and pine forests into regions of everlasting snow and glacier. Relative position, contour, and relief, being alike influential in modifying the external conditions of a region, the accuracy of their determination is of prime importance in all geographical descriptions. Latitude and longitude determine the position; a sufficient number of similar observations fix the contour; and trigonometrical survey or barometrical observation (mercurial or aneroid) can define the relief or outline of elevation. Position and contour are readily shown on maps by actual outline; relief can be indicated by a scale of shading, or, more accurately, by a system of contouring, as shown on Plate I. (Chartography) of the 'Physical Atlas.' Thus, taking the shore-line, which is all necessarily on the same level, as the first contour, we may have similar lines taken at every 50 feet, or every 100



Contour and Profile.

feet, of ascent, and these (when the observations are sufficiently numerous) will exhibit with great accuracy the risings and fallings of the surface configuration. Such a series of lines gives, as it were, the moulding or model of the surface, and a section of

this model in any direction will show the *profile* for that direction. Suppose an island, for example, with its contours taken at every 100 feet of ascent to be represented by fig. 1, then fig. 2 will show the same in profile or absolute elevation in the line of section A B. The steeper the ascent, the closer, of course, will be the lines of contour; and the gentler the slope, the more widely apart will these lines be—a greater extent of surface having to be travelled over. It is by such sections that the profiles of the different continents are usually shown, as will be more fully illustrated when we come to treat of the Mountains and Table-lands in the succeeding chapter.

VI.

THE LAND—ITS HIGHLANDS.

Mountains, their Characteristics and Origin.

46. HAVING directed attention to the general configuration of the land as dependent on its contour and relief, we now proceed to notice the special features of that relief as exhibited in its mountains and mountain-chains—its table-lands and plateaux. And, first, of the mountains, which have long attracted the attention of geographers as one of the most obvious causes of climatic, vital, and political diversity. Defining a mountain as “any portion of the earth’s crust rising considerably above the surrounding surface,” it must be remembered that mountain-ranges are not mere *ridges* rising abruptly on one side and descending as abruptly on the other, but are in reality elevated tracts, often of great breadth as well as length, and consisting of rounded heights, lofty peaks, and boldly-escarped plateaux, with intervening valleys. As, *geographically* speaking, a mountain-range is not a single ridge of elevation, but a brotherhood of many elevations; so, *geologically* speaking, it is not a simple upheaval, the result of one paroxysmal outburst, but the work of innumerable volcanoes and earthquakes operating through many ages. A mountain-chain ten or twenty thousand feet in height and hundreds of miles in length, is thus a thing of slow and gradual growth—upheaval after upheaval, and eruption after eruption contributing to augment the mass of final accumulation. And as the volcanic forces necessarily upheave and disrupt the stratified systems through which they pass, bearing them up on the mountain-flanks, or imbedding them amongst their ejectments, so we can judge of the relative age of mountain-ranges from their associated strata, and say whether they were formed during primary, secondary, or tertiary eras. It is in this manner that Geology enables Geography to account for the external characters of mountains, and to explain why some should be massive and rounded in outline, others serrated, with splintery peaks and pinnacles—some conical and dome-shaped, and others again terraced by crags, or sloping away in long gentle declivities.

47. Understanding the general character and formation of mountainous elevations, we may explain that the term *mountain* is usually applied to heights of more than 2000 feet—all beneath that height being regarded as *hills*, and those of still minor elevation as *hillocks*. A *mountain-chain* or *mountain-range* is a series of elevations having their bases in contact, and their axes or lines of elevation continuous over a considerable extent of country, as the Grampians, Urals, or Andes ; a *group* consists of several ranges more or less connected ; and a *system*, of several groups that evidently belong to the same set of geological operations. Mountain-summits are distinguished by such terms as *cones*, when gradually tapering to a point, as in volcanoes ; *domes*, when more massive and rounded ; *peaks*, when abrupt and insulated ; and by the French word *aiguilles*, or *needles*, when still more pointed, splintery, and detached. Their outlines or profiles are spoken of as *undulating*, *serrated*, *rugged*, and the like, as the case may be ; and their sides are said to consist of *slopes*, *terraces*, *escarpments*, and *precipices*, or of a combination of these—such features depending partly upon their geological structure, and partly on the amount of waste and degradation to which they have been subjected. In general, mountain-ridges have a long, gentle declivity on one side, and a short and abrupt one on the other ; and where this occurs, the longer declivity is spoken of as the *slope*, and the shorter one as the *counter-slope*. This slope and counter-slope is analogous to what in Britain is known as “crag and tail”—most of our hills presenting a bold precipitous front to the west or north-west, and a long slope or tail to the east or south-east—a configuration which has obviously been produced by denuding currents, as the land gradually uprose from the waters of the pleistocene ocean. The depressions and narrow valleys which occasionally intersect mountain-chains are known as *defiles* and *passes* ; and these, offering, as they often do, the only means of transit across mountain-barriers, have ever been objects of political and commercial importance. Some of the most celebrated deeds in history are associated with these passes, and modern engineering is still in search of their easy gradients and facilities.

48. With these explanations, the principal MOUNTAIN SYSTEMS, whose direction and extent will be most conveniently seen by referring to the Atlas, may be here enumerated, the learner remembering that there is much in this arrangement that is arbitrary, or, at all events, not established by actual observation. Thus, in EUROPE, geographers usually distinguish the *British System*, comprising the Grampian, Cheviot, Cumbrian, Cambrian, and Hibernian ranges ; the *Iberian* or Spanish, embracing the Pyrenees, the Cantabrian Mountains, the Sierra Morena, Sierra Nevada, and other ranges of the Spanish peninsula ; the *Alpine*, including the Alps proper, Apennines, Carpathians, Balkan and Hellenic ranges ; the *Scandinavian*, or mountain-chain of Norway and

Sweden ; the *Sarmatian*, or central high ground of Russia ; the *Uralian*, and *Caucasian*. In ASIA we have the *Western System*, comprising the Taurus, Anti-Taurus, Lebanon, and Elburz ranges ; the *South-eastern System*, embracing the Hindoo Koosh, the Himalaya, and the mountains of Birman, Siam, and Cochin-China ; the *Eastern System*, comprising the Kuen-lun, the Peling, Yun-ling, Kihan-shan, In-shan, and other Chinese ranges ; and the *North-eastern System*, extending from the Bolor Tagh in the centre of the continent to Behring Straits, and comprehending the Thian-Shan, the Altai, Yablonoi, and Stanovoi ranges. In AFRICA, the *Atlas System* ; the *Guinea System*, embracing the Kong and Cameroon mountains ; and, so far as is known, the *Eastern System*, comprehending such as the Quotlambi, Lupata, and other contiguous ranges. In NORTH AMERICA, the *Pacific System*, better known as the Rocky Mountains, in the west ; and the *Atlantic or Appalachian System* of the Alleghany Mountains on the east. In SOUTH AMERICA there is the great system of the *Andes* on the west ; the *Brazilian System* on the east ; and the system of *Parimè*, between the rivers Amazon and Orinocco on the north. Of AUSTRALIA and its mountain-chains we know too little as yet to enable us to arrange them into groups or systems, though one main ridge, extending along the eastern coast from Torres Straits on the north to the extreme point of Van Diemen's Land on the south, would seem to indicate a sameness and continuity of geological upheaval. Besides the above, other minor systems have been named, but enough is here indicated to lead the learner to the appreciation of the fact, that the principal mountain-chains may be arranged in groups or systems that have arisen apparently from the same set of geological causes acting more or less continuously along the line or direction of their elevation.

49. As mountain systems exercise very decided influences on the natural history of the globe, and as they generally appear in ranges, consisting either of one central chain with branches or spurs running off at right angles, or of several chains running less or more parallel to each other, various theories have been advanced to account for their upheaval, their parallelism, and their geographical connections. Thus, as their central masses generally consist of igneous rocks (see Atlas, Geological Map of Europe), which have been protruded from below, and as this protruding force must have acted along the line of least resistance in the crust, the question arises, What is the determining cause of these directions ? According to Elie de Beaumont, a French geologist, every system of mountains occupies a portion of a great circle of the globe—a cleft or fissure being more easily made in that direction than in any other ; and he endeavours to show that ranges of the same age are parallel to one another, even when in opposite hemispheres. Professor Hopkins of Cambridge, treating the subject from a purely mathematical point of view,

has also shown that, when the upheaving force acts on a single point, the lines of upheaval must radiate from that point; hence lofty central mountains with diverging spurs. He further shows that, when the expansive force acts uniformly over a wide area, the lines of greatest tension or upheaval must be in the direction either of the length or of the breadth of that area, and that if the crust yields in more places than one, the fissures would necessarily be parallel. Of course, this uniformity of system has been considerably obscured, if not modified, by subsequent geological changes; and it must also be remembered that, whatever may give the initial direction to a mountain-range, its subsequent growth is ever more a matter of *accumulation* than of *upheaval*.

Mountain-Chains and Systems.

50. Accepting such generalisations as initiatory steps towards the explanation of one of the most important problems connected with the history of our planet, we may now briefly advert to the character of the *mountain-chains or ranges* of which the preceding systems are respectively composed. Under the *British System* are embraced the northern or Ross-shire range, the Grampians, the Cheviots, the Cumbrian or Cumberland mountains, the Cambrian or Welsh, the Devonian, and the Hibernian or Irish—all having less or more a south-west and north-east strike, and all belonging to the earlier or primary geological periods. Their geological structure—huge granitic bosses flanked and capped by the crystalline schists—confers on them considerable boldness and diversity of scenery, though their minor elevation prevents that massive grandeur and ruggedness so often displayed by loftier ranges. None of them rise to the height of perpetual congelation, but in an insular and northern situation like Britain, their cold, heath-clad, and inhospitable summits exercise a decided influence alike on the scenery, climate, and productions of the country.—The *Iberian or Hesperian System* embraces the Pyrenees, Cantabrian Mountains, mountains of Toledo, Sierra Nevada, Sierra Morena, and other associated elevations that give character to the rocky table-land and peninsula of Spain. Less ancient than the British system, but loftier in their altitude, and more extensive in their ranges, they exercise a still more decided influence on the external conditions of their region. Ranging chiefly in an east and west direction, and upheaving rocks of primary and secondary formation, rising in many parts above the line of perpetual snow (which ascends from 8000 feet in the Pyrenees to 11,000 feet in the S. Nevada), and being intersected by numerous deep defiles and narrow valleys, they create great diversity of scenery, climate, and production, and are of themselves the frequent storehouses of the minerals and metals.

51. Under the *Alpine System* are usually comprehended the

whole of those extensive and lofty mountains which form Switzerland as a centre, ramify in ranges more or less persistent, and confer on southern Europe its distinctive and peculiar character. These ranges have many minor and local subdivisions, but for our present purpose it will be enough to arrange them into Eastern and Western Alps proper; the Gallo-Francia mountains, including the Jura, Vosges, &c.; the Apennines, traversing the entire peninsula of Italy; the Slavo-Hellenic, embracing the Herzegovina, Balkan, and Pindus chains, which lie between the Adriatic and the plain of the Danube; and the Hercynio-Carpathian range, which rises in rugged lines between the valley of the Danube and the great central plain of Europe. The component members of this grand Alpine system are of various geological ages—ranging from the granites and crystalline schists of the Western Alps, through the secondary limestones and altered shales of the Jura, to the tertiary beds of the sub-Apennines and the recent lavas and scoræ of Vesuvius and Etna. Connected with the older range of the Pyrenees on the west, and with the still active cones of Vesuvius, Etna, and the Lipari Islands on the east, the Alpine system may be said to have been on the increase from the earliest geologic ages to the present moment—gaining accession after accession, and this even since the tertiary period, to much of the Auvergne, Apennine, and Hellenic ranges, and to the Alps proper, an additional altitude of not less, perhaps, than 4000 or 5000 feet. Being of different geological structures and altitudes, the different members of the system present great diversity of character and aspect. Rising in many places above the snow-line, which sinks from 9000 feet in the Alps to 6000 feet in the Carpathians, they are rugged with peak and precipice, glacier and narrow gorge, as in the Alps—swelling and sloping in outline, as the Apennines—crateriform and terraced, as the hills of Auvergne—rocky and precipitous, as in the Pindus and Balkan ranges—or rich in mineral treasures, as the mountains of Transylvania.

52. The *Scandinavian System*, as the name implies, embraces the whole of the mountainous highland of Norway and Sweden, and extends, in a north-eastern direction, from the Naze to the North Cape—a distance of more than a thousand miles. It consists rather of a series of plateaux than of a continuous ridge; these mountain-elevations narrowing from a breadth of 200 miles in the south to 60 or 40 in the north, and being distinguished as the Hardaangar or Langefeld in the south, the Dovrefeld in the middle, and the Kiølen mountains in the north. Intersected by numerous ravines and gorges, the range presents a steep face and rugged coast-line of fiords and precipices to the North Sea on the west, and a terraciform slope to the shores of the Baltic on the east. A great portion of the range rises above the limit of perpetual snow (which ascends from 2400 feet at the North Cape to 5000 feet in the Langefeld), but beneath that limit there is

much of crag and forest. The whole consists of granitic, crystalline, and silurian rocks, rich in minerals and metallic ores.—The *Sarmatian System* (from the ancient Sarmatia or Poland) is meant by geographers to embrace that swell of country which stretches diagonally through Russia from the plain of Poland to the Uralian Mountains, and which forms the long watershed of Northern Europe—turning the waters of the Vistula, Duna, Dwina, and Petchora to the Baltic and White Seas, and those of the Volga, Don, and Dneiper to the Caspian and Black Seas. There are no true mountains throughout its extent—the Valdai Hills attaining an elevation of only 1100 feet; but it forms a great watershed, and as such exercises considerable influence on the Physical Geography of Europe.—The *Uralian System* embraces the well-known range of the Ural Mountains, which form the natural boundary between Europe and Asia, and the watershed between the extensive basins of the Volga and Obi. Departing from the usual strike of the Old World mountains, the Urals run in a true meridional direction for more than 1600 miles, and consist of round-backed plateau-shaped masses of very moderate elevation, generally not exceeding 2000 feet, and culminating in a few points only at upwards of 5000 feet. The range consists of an axis of igneous rocks flanked by crystalline strata and the older palæozoic formations, and is rich in the precious minerals and metals.—The *Caucasian System*, which forms another part of the boundary between Europe and Asia, extends in one immense chain of 750 miles from the Black Sea to the Caspian, and separates the basins of the Kuban and Terek on the north, from those of the Kur and Rion on the south. The range is a bold and lofty one, full of glens and mountain-fastnesses, and rises in many places above the snow-line, which there attains the altitude of 11,000 feet. The culminating point is Mount Elburz, near the centre of the chain; its height being 18,493 feet, or 2749 feet higher than Mont Blanc.

53. Turning next to Asia, which is the great headquarters of mountains and mountain-chains in the Old World, we may begin with the *Western System*, under which is comprised those numerous ranges that lie between the Levant and Black Sea on the west, and the Indus on the east, and give feature and character to the highlands of Asia Minor, Persia, and Afghanistan. Among these may be noticed the Taurus and Anti-Taurus, whose hilly and irregular ranges encircle the table-land of Turkey in Asia, and whose culminating point is Mount Argish, attaining an elevation of not less than 13,000 feet. The Lebanon range, proceeding transversely from the Taurus southward along the Syrian coast to Mount Hermon, 10,000 feet high, and thence through Palestine into the peninsula of Sinai, where Horeb, 8593, and Sinai, 7497 feet, may be said to terminate the chain. The mountains of Armenia, intimately associated with the Taurus and Anti-Taurus, and but prolongations of the same ranges, trend-

ing eastward into the Elburz, and southward into the Zagros and hills of Kurdistan—their culminating points being the snowy Ararat in Armenia, 17,210 feet, and the Sheikhiya, 10,120 feet, in Kurdistan. The Elburz range, a further prolongation of the Taurus and Armenian mountains, skirting the southern shores of the Caspian, where it attains, in the slumbering volcano of Demavend, an altitude of 14,675 feet, and thence stretches eastward into the Hindoo Koosh and mountains of Central Asia. The Hindoo Koosh or Hindu Koh, separating Afghanistan and the Punjab from Independent Tartary, and forming the watershed between the Amoo and the Indus. Its maximum elevation is about 20,000 feet, and, trending eastward in broad massive ridges into the Kuen-lun and Himalaya Mountains, it may be regarded as the commencement of the great central system of the Asiatic continent.

54. Closely connected with, and following the Western, is the great *South-Eastern System* of Asia, which comprises not only the highest mountains in that continent, but the highest known elevation on the surface of the globe. Pre-eminent in the system stands the chain of the Himalaya ("abode of snow"), stretching in a somewhat south-easterly direction between the basin of the Ganges and the upper basin of the Brahmapootra, forming the northern boundary of India, and constituting the southern buttress of the great central table-land. The range extends about 1500 miles in length, varies from 100 to 300 miles in breadth, and rises in many points (upwards of forty, it is said) to an altitude of 23,000 feet; the three highest peaks being Everest, Gahurishanka or Deodunga, 29,002 feet—Kinchinjunga, 28,156 feet—and Dwhalagiri, 26,826 feet. The snow-line rises, according to position in the range, from 13,000 to 16,000 feet, and a large portion being thus perpetually covered with ice and snow, the Himalayas present every possible feature of mountain grandeur—peak and precipice, gorge and glacier, rugged ravine and headlong waterfall. Abruptly separated from the Himalaya by the valley of the Brahmapootra, but still holding more or less in the same axial direction, are the mountains of Assam; and beyond these lie the well-defined ranges of Burmah, Siam, and Cochin-China—all trending in a southerly direction, separated by intervening river-valleys, and giving contour and character to the Cambodian and Malayan peninsulas. Little is known of these peninsular mountain-ranges, or of their elevations; but approaching the equator, we know that to great elevations they are covered with impenetrable forest-growth. Outlying the system, but still connected with the same area, are the mountains which give conformation to the triangular-shaped peninsula of Hindoostan. These are the Vindhya chain, which form, as it were, the northern barrier of the Deccan—the Western Ghauts, that support it on the west—and the Eastern Ghauts, that guard it on the east—the two latter chains converging into

the loftier heights of the Nilgherri Hills on its southern border. The plateaux of the Deccan rise, stage by stage, from 1500 feet in the north to 4000 or 5000 feet in the south ; the Western Ghauts from 3000 to 5700 feet ; the Eastern from 3000 to 4000 feet ; and the Nilgherries in many points from 6000 to 7000 feet—attaining their culmination, 8760 feet, in the peak of Dodabetta.

55. Starting from the same central knot of the Bolor, and extending eastward to the Pacific, we have next what is termed the *Eastern System of Asia*, a series of vast and partially known ranges, associated with high desert table-lands in Thibet and upper Tartary, and alternating with alluvial river-plains in the eastern districts of China. The system commences with the chain of the Kuen-lun, rising between the valleys of the upper Indus and Brahmapootra on the south, and those of the Amoor and Yarkhand on the north, and extending in an easterly direction for nearly 1200 miles, at an altitude of from 15,000 to 18,000 feet. Lying, as it does, in the same line with the Elburz on the west, and prolonged into the Pe-ling on the east, the whole looks like one great range, and is, no doubt, geologically dependent on the same axial elevation. Holding eastward, as we have said, into the main ridge of the Pe-ling Mountains, that rise between the basins of the Yang-tse-kiang and Hoang-Ho, the system at the same time diverges, fan-like, into the Yun-ling chain, between Thibet and China ; the Nan-ling between the basins of the Yang-tse-kiang and Canton River ; and the conjoint chains of the Kihan-Shan, In-shan, and Khingan, that trend in a north-easterly course, and form the southern wall of the great Mongolian Desert.

56. Commencing in like manner with the central knot of the Bolor, and stretching away in broad, more or less parallel, lines to Kamtchatka and Behring Strait, occurs the *Northern System of Asia*, flanked by the arid deserts of Tartary and Mongolia on the south, and descending by gradual stages to the great plain of Siberia in the north. The system embraces the Thian-shan or Celestial Mountains, ranging in an easterly direction for nearly 1400 miles, ascending in greater portion above the line of perpetual snow, and exhibiting throughout a number of volcanic cones, some of which, like Pe-shan and Ho-tscheou, are still active at elevations of more than 10,000 feet. In north-easterly association we have next the Altai Mountains, comprising several parallel ridges, the Daurian Mountains, and the Yablonoi Mountains—all rising between Siberia and Mongolia, and separating the basin of the Amoor from those of the Yenesei and Lena. These closely associated ranges are said to have an average elevation of from 5000 to 8000 feet ; embrace several active and many extinct volcanoes, rise high in numerous points above the snow-line, and culminate occasionally in peaks of 10,000 and 11,000 feet high. Still northward and eastward, but ramifying more irregularly, occur the Aldan, Stannovoi, and other chains that

terminate in the volcanic system of Kamtschatka. These ranges are for the most part covered with snow (the snow-line sinking to 4500 and 4000 feet in Eastern Siberia), and contain numerous volcanic cones, which, like Schiwelutsch, 10,548 feet, and Klient-schewsk, 16,131 feet, are still in active eruption.

57. Of the African continent, to which we next turn, our knowledge is yet too limited to enable us to do more than merely advert to some of the more prominent mountain regions as likely belonging to separate and independent systems. In the extreme north we have the *Atlas System* rising between the Mediterranean seaboard and the Sahara, and extending from Tripoli on the east to the Atlantic on the west. Geologically, it is evidently connected with the systems of Southern Europe, and consists of three or four parallel ranges, which ascend, stage by stage, from the basin of the Mediterranean, and increase in height from east to west—being about 2000 feet in Tripoli, 4500 in Tunisia, 7700 in Algeria; while in Morocco, Mount Miltzin (*Atlas*) ascends to 11,400 feet, and Jebel Tedla to 13,000 feet, or above the line of perpetual congelation. Several secondary spurs proceed from the main ranges—one northward, and terminating in Cape Spartel, and several others southward into the desert plateaux of the Sahara.—Next in importance is the *Abyssinian System*, connected with and forming the lofty table-land (*Amhara*) of Abyssinia and Upper Ethiopia. This plateau, which is 8000 feet above the sea-level, is supported and traversed by several clustering ranges of great elevation, and in many points above the line of perpetual snow, which is there about 14,000 feet. The two most persistent chains, under the names of the Samen and Taranta, strike in a northerly direction between the upper forks of the Nile and the Red Sea, and, skirting the shores of the latter, are prolonged into the lower hills of Egypt, which, at the Gulf of Suez, connect themselves with Sinai and the mountains of Syria. The culminating points in the Samen or upper range are Ras Detschen, 15,986 feet; Buahat, 15,000; Abba Jarret, 14,707; and Shubattai and Beyeda, each 12,000 feet. In the Taranta or lower range the heights descend from 9000 to 6000 and 5000 feet towards the Red Sea and the plain of Egypt.—In Western Africa the mountains are by no means well known; but under the *Guinea System* are usually embraced the Kong and Cameroon ranges—the former rising between the Gulf of Guinea and the Niger, and generally averaging from 1200 to 3000 feet (*Soracte*, 1278, and Mount Rennel, 3200 feet), and the latter stretching eastward and unknown into the centre of the continent, and rising in many points to elevations of 4000, 6000, 9000, and even 13,000 feet.—In Southern Africa the surface exhibits a series of sandstone plateaux, resting on and intersected by granitic rocks, rather than a series of well-defined mountain-chains—these flats (*karoos*) rising step by step from south to north at elevations of 2000, 4000, and 6000 feet above the sea-level. The steps of this ascent consist

of rocky walls and flat-topped mountains, intersected by narrow gorges (*kloofs*) which form the only means of passage from terrace to terrace, and these mountains (*bergs*) may, for the sake of reference, be denominated the *Cape System*. The system consists, 1st, of the Zwellendam range, about 20 miles inland, stretching for nearly 200 miles in length, and attaining in Table Mountain an elevation of 3816 feet; 2d, the Zwarte or Black range, about 30 miles further northward, and separated from the Zwellendams by the Kannaland Karoo; and, 3d, the northern chain, consisting of the Roggeveld, Nieuwveld, Winter Mountains, Sneeuwveld, Witte-bergen, and other contiguous ranges, separated from the Zwarte range by the Great Karoo, and attaining heights of 4000, 6000, and even 10,000, as in the Compass Berg in the Sneeuwveld or snowy range.—From this point begins what may be termed the *Eastern System*, consisting of the Drakenberg or Quotlamba Mountains (10,000 feet), the Lupata Mountains (8000 or 10,000 feet), and other ranges, which hold northward in parallel lines and increasing altitudes towards the equator, where several of the higher peaks (Kenia and Kilmandjaro, 17,000 and 20,000 feet) are said to be covered with perpetual snow.

58. Of Australia and its mountain-chains, it has been already stated (par. 48) that we know too little to enable us to arrange them into groups and systems, though one main ridge, extending along the eastern coast from Torres Strait on the north to the extreme point of Tasmania in the south, would seem to indicate a sameness and continuity of geological upheaval. This chain consists chiefly of granites and porphyries, intersected and overlaid in many places by recent volcanic products; is extremely rugged and inaccessible on the coast side, but slopes gradually towards the interior; and seldom exceeds 5000 feet in elevation.—Mount Kosciusko, 6500 feet, and Sea-View, 6000 feet, in Australia—and Mount Humboldt, 5502 feet, and Benlomond, 5010 feet, in Tasmania—being the highest known points of elevation. Of the mountains that occur in the islands of the Pacific and the Indian Archipelago it may be remarked that they are chiefly active volcanoes, and though often occurring in obvious linear connections, and of great altitudes (6000, 10,000, and 14,000 feet), they are merely to be regarded as chains in embryo, and the rudiments of systems yet to be elaborated.

59. Having reviewed the mountain systems of the Old World, we now turn to those of the New, where the arrangements are altogether on a simpler and more uniform plan. In North America we have first the *Eastern or Atlantic System*, so called from its general proximity and parallelism to the Atlantic seaboard. It separates the waters that flow eastward into the Atlantic from those that flow westward into the basins of the Mississippi and St Lawrence; and though trending in one continuous direction from the Gulf States to the St Lawrence, may be said to consist of two main divisions—the Alleghanies or Appalachians on

the south, and the Green and White Mountains on the north, separated from each other by the narrow cross-valley of the Hudson. In length the system is nearly 2000 miles, has an average breadth of 100 or 130 miles; and though its mean altitude is only about 2500 feet, it yet ascends, in Mount Washington in New Hampshire, to 6652 feet—in Black Mount, between Tennessee and Carolina, to 6420 feet—and in Mount Katahdin in Maine, to 5360 feet. In the northern section of the system the ridges of the Notre Dame, the Green, White, and Adirondack Mountains are more or less irregular and interrupted; but in the southern, the Alleghanies consist of several closely parallel chains of great continuity, though frequently cut across by ravines and river-courses. Geologically, the system consists chiefly of the older or palæozoic rocks, flanked on both sides by an extensive development of carboniferous strata; and its eastern slopes comprise some of the finest and most diversified country in the American Union.

60. Interiorly, and beyond these eastern ranges, the country is one immense plain till we come to the *Western* or *Pacific System*, which, under the familiar name of the Rocky Mountains, stretches in unbroken ranges from the Isthmus of Panama to the shores of the Arctic Ocean. This extensive system, which, after all, is but the northern prolongation of the great backbone of the New World, consists of two main ridges—the Pacific or Oceanic, skirting the western seaboard from Cape Lucas in California to Cape Elizabeth in Russian America, and the Rocky Mountains proper, extending in double, and sometimes in treble, chains from Panama to the Arctic shores. The former of these ranges forms the watershed between the Pacific on the west and the Colorado, Columbia, and Colville on the east. Though continuous as one great range, it consists of several members, such as the Sierra de S. Lucia and Sierra Nevada in California, whose highest points are Mount St John, 8000 feet, and Mount Tsashti, 14,000 feet; the Cascade range in Oregon territory, culminating in Mounts Hood and Jefferson, 15,000 feet, and Mount St Helens, 15,750 feet; and the Sea Alps in the north, having their highest points in Mount Fairweather, 14,783 feet, and Mount St Elias, 14,970 feet. Geologically, the range is of comparatively recent origin, contains many extinct and dormant volcanoes, and evidently connects itself with the still active series of the Aliaski peninsula and the Aleutian Islands. The latter or Rocky Mountain range forms, on the other hand, the long watershed between the tertiary valleys of the Colville, Frazer, Columbia, and Colorado on the west, and the great Lake region and the plains of the Mississippi on the east. It consists, in like manner, of several members, which, though obeying the same axial direction, yet separate and converge so as to constitute a series of plateaux of varying magnitude and elevation. At the southern extremity of the range we have, first, the volcanic chain of Guatamala, rising in the craters of Atitlan and Agua to 12,500 and 15,000 feet; second, the cluster-

ing or transverse mountains of Anahuac or Southern Mexico, ascending in Orizaba and Popocatepetl to 17,347 and 17,720 feet; third, the Cordillera of Cohahuela and Potosi, and the Sierra Verde and Madre, which support the lofty table-land of New Mexico, and ascend in Pike's Peak and Long's Peak to 10,000 and 12,000 feet; fourth, the Wind River Mountains, between Nebraska and Oregon, attaining in Fremont's Peak an elevation of 13,568 feet; and, lastly, the northern and parallel ranges of the Rocky Mountains proper, which rise in Mount Hooker to 15,700 feet, and in Mount Brown to 15,970 feet. Geologically, the whole of this vast range is of ancient formation, with the exception of the volcanic mountains of Guatamala and Mexico, and these are evidently portions of the still active development of Central America and the West India Islands.

61. In South America the pre-eminent system is that of the *Andes*, which extends along the Pacific seaboard from Terra del Fuego on the south to the Isthmus of Panama on the north, there connecting itself with the onward prolongation of the Rocky Mountains. As a mountain-range, the Andes forms one of the most definite and persistent on the globe—skirting in unbroken ridges the entire Pacific shore for nearly 4500 miles in length, and varying in breadth from 40 to 350 miles. In some places the range consists of a single ridge—in other places, of two or more ridges supporting lofty but narrow plateaux; and in general it presents a steep slope towards the Pacific, from which it is distant from 20 to 80 miles; while towards the east it descends by gradual stages into the broad plains of the Orinocco, Amazon, and La Plata. According to Humboldt, the mean elevation of the Andes is 11,830 feet, and the extent of surface covered by their bases not less than 531,000 geographical square miles. Geologically, the system is composed of granites, greenstones, and porphyries, flanked by metamorphic schists and palaeozoic strata, but exhibits throughout a greater number of active craters than any other mountain-chain, and, as a consequence, is largely crowned and covered, especially on its western slopes, by vast accumulations of lava, scorise, and other volcanic products. Though presenting one continuous axis, the range consists of several members, known by the countries in which they occur—as the Patagonian, the Chilian, the Bolivian, the Peruvian, and the Columbian Andes. The Patagonian section consists of a single range of moderate elevation, but ascending in several points (Mount Darwin, Mount Stokes, the volcanoes of Yanteles and Minchinmadiva) to 6400 and 8000 feet. As the snow-line descends in Southern Patagonia to 3000 feet, much of the range is perpetually frozen, and glaciers, unknown in other parts of the Andes, make their appearance in the higher glens and gorges. The Chilian Andes extend, in like manner, in one immense ridge; and though their mean elevation is inferior to those of Bolivia, they yet contain the giant Aconcagua—the culminating cone of the system, and the highest known point in the New

World continent. The snow-line in this portion of the system rises from 8000 to 10,000 feet; and high above it in perpetual winter rise the lofty peaks of Aconcagua, 23,910 feet; Tupungata, 15,000 feet; and the volcanoes of Chilian and Villarica, 16,000 feet. Next in northward order occur the Andes of Bolivia, rising in two parallel ranges—the Cordillera of the Coast and the Cordillera Real—and supporting between them the table-land of Desaguadero, 13,000 feet above the sea, 500 miles in length, from 30 to 40 miles wide, and enclosing Lake Titacaca at an elevation of 12,846 feet. In this, the central portion of the system, the snow-line ascends from 15,000 to 18,000 feet; and high above it, in the western range, rise Sahama, 22,950 feet—Chiquibamba, 21,000—and the volcanoes of Gualatieri and Arequipa, 20,000 and 22,000 feet; and in the eastern range, Sorate 21,286, Illimani 21,140, and Cochabamba 17,000 feet. Northwards from the Bolivian plateau, the Andes open up in three parallel ridges, known as the eastern, central, and western Cordilleras of Peru—the western being the highest, and separated from the Pacific by a sandy desert 120 miles in breadth. In these Cordilleras the highest points are Sasaguanca, in Lima, 17,904 feet, Vilcanota 17,525, and the Knot of Pasco 11,800 feet. Leaving the Cordilleras of Peru, we next meet, in northward order, the Colombian Andes, or the Andes of Quito—a lofty volcanic portion of the system, which rises in double or treble ridges, one main portion extending in the direction of Panama, and another bending north-eastward to the Caribbean Sea. In this region, much of which lies directly under the equator, the snow-line rises to 15,000 or 16,000 feet; and hence all the higher peaks and volcanoes—Chimborazo 21,424, Cotopaxi 18,875, Antisana 19,132, Coyambe 19,535, and Tolima 18,120 feet—are covered with perpetual snow. Such are the various portions that constitute the giant system of the Andes—a system which, whether in its extent and linear continuity, its boldness and altitude, its high inhabited table-lands, its mineral riches, or its physical influences, is, even more than the Himalayas, the most remarkable on the globe.

62. The system next in importance in South America is that of *Brazil*, occupying the eastern portion of the continent, extending in several parallel ranges from the plains of the Plata on the south to those of the Amazon on the north, and spreading inland for nearly 1800 miles in a broad plateau, whose mean elevation is about 3200 feet. These ranges or ridges of table-land are separated from each other by the affluents of the Amazon and the St Francisco on the one hand, and by those of the Paraguay and Parana on the other, and succeed each other, ridge and plain, with wonderful continuity. Proceeding from the Atlantic westward, we have first the Sierra Espinhaço, whose culminating heights are Itarube 5960, Piedade 5830, and Itacolumi 5750 feet, and the Sierra do Mar or Sea-range, which

attains in Morro dos Candos an elevation of 4476 feet; second, the Sierra Tabatinga, forking northward into the S. Irmaos and S. Mangabeiros; third, the Cordillera Grande, whose chief heights are from 6000 to 7000 feet; and lastly, the Sierra de los Vertientes and other inferior ridges, that gradually descend into the great central plain of the continent. Geologically, the Brazilian system is eminently primary, consisting of granitic protrusions and crystalline schists rich in the precious minerals and metals; abounding, from the character of its rocks, in picturesque beauty; and from its tropical situation and minor elevation, clothed to the summits with an exuberant and varied vegetation. The last and only other mountain system of South America is that of *Parimè*, which occupies the oval tract of country lying between the Amazon and Orinocco, and forms the high ground from which descend many of the minor affluents of these gigantic rivers. This plateau, whose mean elevation is from 1600 to 2000 feet, is traversed in an east and west direction by several closely-set ridges (Sierra Acarai, *Parimè*, Pacaraima, Imataca, &c.) which, though of no great general elevation, yet ascend in Duida to 7149 feet, in Poraima to 7450, and in Maravaca to 10,500 feet. Like the mountains of Brazil, the system of *Parimè* consists of granitic bosses, crested and flanked by crystalline schists, and has not inaptly been described as a primary island rising from the vast tertiary and recent expanses of the Orinocco and Amazon. Lying almost directly under the equator, the higher sierras are clothed with impenetrable forest-growth; while the lower grounds, according to the seasons, are alternately arid wastes, or covered with a carpeting of the most luxuriant grasses.

63. Such are the principal mountains of the world as arranged by geographers into groups and systems. The arrangement may not in every case be a natural one—that is, the mountains composing some so-called “system” may not strictly belong to the same set of geological causations—but the arrangement, such as it is, greatly facilitates reference, and aids our comprehension of the effects produced by any mountain-group on the climate and vital economy of the region in which it is situated. The arrangement has also its topographical advantages, for little can be done in the way of correct description till the objects to be described have been arranged and classified according to some principle of similarity or sameness of origin. But whatever may be the ultimate grouping, we see in mountain chains and systems one of the most important features in the physical machinery of the globe. Rising and falling—here in easy undulations, there in steep peaks and ridges—here in abrupt crags, and there in gentle slopes—they produce a diversity of surface eminently fitted for diversity of vegetable and animal life. Presenting their high ridges to the moisture-laden currents of the atmosphere, they serve as so many points of condensation, producing clouds, mists,

and showers, that temper the heat in the lower regions, and refresh and nourish their vegetation. Elevated into regions of perpetual snow, in hot climates they cool the breezes that descend from their heights, while their snows and glaciers become perennial storehouses, which, under the summer sun, yield a copious supply to the streams and rivers of the thirsty lowlands. From their geological structure and formation, they are necessarily the chief repositories of the precious minerals and metals; and even such deposits as occur in the sands and gravels of their streams have been worn and washed from their disintegrated veins. Their healthy, life-bracing heights have ever been the notable nursery-grounds of active, courageous, and independent races; while their snow-clad ridges become boundaries and barriers to nationalism, as well as to the dispersion of plants and animals, more impassable even than the breadths and depths of the ocean.

Table-lands or Plateaux.


64. Next in importance, in the vertical relief of the land, are those elevated expanses known as *table-lands* and *plateaux*. A table-land, as the name suggests, is a flat elevated surface; but this idea of flatness must be received only in a comparative sense, for the surface, though plain-like on the whole, is usually diversified by minor undulations and irregularities. Being in effect broad mountain-masses, these plateaux form the *gathering-grounds* and sources of some of the noblest rivers, while their elevation confers on them a climate and a vegetable and animal life distinct from that of the surrounding lowlands. They are inseparably associated with the mountain systems, most of these systems not rising in narrow ridges from low-lying plains, but towering above broad elevated regions, which seem to be formed and supported by their bases. Thus, on turning to the map of Asia, it will be seen that all the great rivers flow north, south, east, and west from the central region, which consists in reality of a succession of lofty terraces or plateaux. First, we have the table-land of Iran or Persia (including large tracts of Beloochistan, Afghanistan, and Bokhara) rising from 2300 to 3500 feet above the sea-level, upwards of 300,000 square miles in area, and presenting throughout a riverless, parched, and desolate region; next in altitude, the great sandy and rainless desert of Gobi, rising from 4000 to 6000 feet, and occupying an area of nearly 400,000 square miles; then rising on either side of this, towards the centre, the plateaux of Dzungaria and Upper Tartary, less arid and more varied in surface; and lastly, the still loftier plateau of Thibet—the highest inhabited region in the world, with an elevation of from 11,000 to 15,000 feet, and an area of 166,000 square miles. Besides these great central uplands, there are the lateral and more isolated plateaux of the Deccan, rising from 1600 to 2000

feet in Hyderabad, to 4000 feet and upwards in Mysore; of Arabia the sandy and arid, varying from 3000 to 6000 feet high, and spreading over an area of more than 700,000 square miles; of Armenia (7000 feet), supported by the Taurus and Anti-Taurus, and extending from the Dardanelles to the Caspian; and, lastly, that of Ust Urt between the Caspian and Aral Seas. With the exception of the Deccan, and the mountain-platforms of Armenia and Thibet, the whole of these table-lands, from Arabia on the west to Gobi or Shamo on the east, belong to one great belt of arid and rainless country, sandy and shingly in soil, desert in character, and evidently belonging to the same geological age and formation.

65. In Europe, less elevated and more broken up by seas, we have a smaller development of table-lands, and these generally of limited area, and in the southern division of the continent. The most notable is that of Castile in Spain, having an elevation of from 2000 to 2300 feet, and traversed by hilly ridges that give great irregularity and diversity to its surface. There is next the less defined upland of Switzerland, from 3000 to 4000 feet in elevation; and trending north-eastward, in the same direction, the lower plateaux of Bavaria and Bohemia, the latter having an elevation only of 900 or 950 feet above the sea. The so-called plateaux of Auvergne and of the Scandinavian and Balkan chains, may be regarded as mere mountain-flats too limited in extent to possess any physical feature, or to exercise any influence distinct from those of their associated ranges.—Of Africa we have too scanty information to speak with certainty as to the plateaux and mountains that may rise within its interior; but we know that much of the Sahara or Great Desert is of a flat elevated nature (from 1500 to 2000 and 4000 feet), and that inland from the coasts of Congo and IZango the country assumes the character of a lofty table-land. There is also the plateau of Abyssinia (Amhara), 8000 feet above the sea-level, supported and traversed by several clustering mountain-ridges, which form the gathering-grounds of the Atbara and Blue Nile; and, lastly, the *karoos* or terrace-plains of Southern Africa, rising stage by stage towards the interior to an elevation of 2000 feet—carpeted with grass during the rainy season, and parched and barren for the rest of the year.

66. In the New World the great superficial contrast is less between mountain and table-land or plain and elevated upland, than between the gigantic mountain barrier that walls the Pacific from one extremity of the continent to the other, and the low broad plains that stretch eastward from its base towards the Atlantic. Nevertheless, in South America, the chain of the Andes presents several table-flats of vast elevation—the most remarkable being that of Bolivia, a great table-land 120,000 square miles in extent, rising from 11,000 to 12,500 feet above the sea, and early the seat of a busy and wealthy civilisation. Much also of the interior of Brazil partakes of the table-land character, having a mean eleva-

tion of 3000 feet, and traversed by the sierras (par. 62) that give feature to that fertile and tropical country.—In North America, from the Russian territory on the north to Mexico on the south, there occur a series of elevated uplands, upborne, as it were, by the parallel ranges of the Rocky Mountains. One of the most remarkable of these plateaux is that comprising the highlands of Oregon and the saline desert or inland basin of Utah, whose elevation is from 4000 to 5000 feet, and the waters of which, having no outlet, form a series of salt lakes, one of which (Utah) is of considerable extent, and almost saturated with salt. The most decided of the American table-lands, however, is that of Mexico, not more remarkable for its elevation than for its persistent extent. “On the eastern and western coasts,” says M. Balbi, “are low countries, from which, on journeying into the interior, you immediately begin to ascend, climbing, to all appearance, a succession of lofty mountains. But the whole country is thus in fact raised into the air from 4000 to 6000 and 8000 feet.



The conformation of the country has most important moral and physical results ; for, while it gives to the table-land, on which the population is chiefly concentrated, a mild, temperate, and healthy climate, unknown in the burning and deadly tracts of low country into which a day's journey may carry the traveller, it also shuts out the former from an easy communication with the sea, and thus deprives it of a ready access to a market for its agricultural productions.” As with the Mexican table-land so in fact with all others of any decided elevation. A distant island in the ocean is not more separated from its contiguous continent, or more strongly marked by its own physical peculiarities, than a high mountain-walled tract raised several thousand feet into the atmosphere is characterised by a climate and vegetable and animal productions unknown in the regions that surround it.

RECAPITULATION.

In the preceding chapter attention has been directed to the more elevated portions of the Land, as consisting of mountains and table-lands. These mountains have been arranged by geographers into *chains* and *ranges* ; and these ranges, again, into

groups and systems that occupy the same contiguous area, and apparently belong to the same series of geological operations. The continuity of mountain-chains has been observed from the earliest times, and hence the antiquity of the names by which many of them are distinguished ; but their classification into groups and systems is of recent date, and liable to correction and amendment as clearer and more correct views are obtained respecting the forces on which their formation and elevation depend. In the mean time, and for the sake of reference, the mountains of Europe have been arranged into the British, Iberian, Alpine, Scandinavian, Uralian, and Caucasian systems ; those of Asia into the Western, South-Eastern, Eastern, and North-Eastern ; and those of Africa into the Atlas, Abyssinian, Guinea, Cape, and Eastern systems. In the New World the mountains of North America are usually arranged into the Eastern or Atlantic system, and the Western or Pacific ; while those of South America are distinguished as the systems of the Andes, of Parimè, and of Brazil. As to the table-lands, the more important and better known in the Old World are those of Castile, Switzerland, Bavaria, and Bohemia in Europe ; and of Armenia, Arabia, Persia, Tartary, Mongolia, Thibet, and the Deccan in Asia. In the New World, those of Bolivia and Brazil are the most notable in South America ; while in North America the only similar tracts deserving of notice are the table-land of Mexico and the desert uplands of Utah and Oregon. In whatever form the highlands of the globe appear, whether as linear mountain-chains or as broad-spreading plateaux, they exercise most important influences on climate, and consequently on the distribution of plants and animals. In the torrid zone they afford the climate and produce of temperate regions, and in temperate zones they assume the characteristics of polar latitudes ; while everywhere they are the great gathering-grounds of glacier, stream, and river—dispensing their stores to the thirsty lowlands in moderated but never-failing supplies.

VII.

THE LAND—ITS LOWLANDS.

Plains and Deserts.

67. As the higher and more irregular portions of the earth's surface consist of mountains and table-lands, so the lower and more level consist of plains and valleys. The one set of features counterbalances as it were the other, and thus contributes to that variety of aspect so pleasing in the landscape, and so indispensable to diversity in its vegetable and animal productions. Though the term *plain* is usually applied to level expanses of no great elevation, and is apt to be associated with ideas of verdure and fertility; yet several of the great plains of the world are considerably above the sea-level, and present every variety of surface, from green grassy flats to deserts of shingle and loose shifting sand. As mountains were the results of volcanic upheaval, continued through indefinite ages, so plains and valleys are the undisturbed portions of the earth's crust, and, in most instances, represent the beds of former seas, and the silted-up sites of lakes and estuaries. Not only does their general contour convey this impression, but their soil and subsoil usually reveal their origin, and point to a time when large expanses of ocean occupied the areas of the present plains, and shallow estuaries and chains of lakes the sites of our alluvial valleys. Bearing in mind this origin, it will help to explain certain appearances of soil and surface, and enable us to account for certain distributions of plants and animals that might otherwise remain inexplicable.

68. In treating of the low level tracts of the Land, the terms *plain* and *valley* are sufficiently general and well understood, and are, therefore, the most frequently employed in geographical description. There are others, however, which refer either to some peculiarity of surface and condition, or are of local origin, and these it may be useful at this stage to explain. Thus, the term *prairie*, though simply the French word for "meadow," is usually applied in a technical sense to the open, slightly undulating, grassy plains of North America; *llanos* are the river-plains of tropical South America, alternately covered with rank vegeta-

tion, and reduced to a desert by periodical droughts ; *selvas* (Lat. *silva*, a wood) the higher tracts of the same region densely covered with forest-growth ; and *pampas*, the treeless but grassy plains of the Parana and La Plata. The term *steppes* is applied to the plains of Northern Asia, generally covered with long rough grass, but also partially wooded, and not unfrequently shingly and desert ; *tundras*, to the boggy flats of Siberia and Northern Russia ; and *tarai*, to the belt of unwholesome jungle that lies between the plain of Hindoostan and the Himalayas. *Sahara* is the long-established and familiar name for the great arid and sandy desert of Northern Africa ; while *karoo* is applied to the open flats in the southern region of the same continent, which are hard and arid in the dry season, but carpeted with grasses and flowers during the periodical rains. In Britain, the terms *dale* and *vale* are usually applied to minor river-plains ; *strath*, in Scotland, to any wide stretch of generally flat-lying land ; and *carse*, to those level alluvial flats that occur in connection with existing estuaries, and which have evidently been reclaimed from their waters either by the ordinary process of silting, or by partial upheaval of the land. The term *delta* is also largely applied to the alluvial land formed at the mouth or rather *mouaths* of a river, such as that of the Nile, which first received this name from the resemblance that the triangular space enclosed by its two main mouths bears to the Greek letter Δ or delta. Other terms than the above are still more local and restricted in their application, and will be better explained in the text as they happen to occur.



Nilotic Delta.

69. The principal plain in the Old World is that usually known to geographers as the *Great Northern Plain*. It may be roughly sketched as commencing with the shores of Holland on the west, and extending eastwards through Prussia, Poland, Russia, and Siberia, without any very marked interruption save the intersecting range of the Uralian Mountains. In width it stretches from the shores of the Arctic Ocean almost to the base of the Carpathians in Europe, and to the table-land of Persia and the flanks of the Altai Mountains in Asia—thus occupying between 4,000,000 and 5,000,000 square miles, or nearly one-third of the entire area of these continents. While it slowly rises from the Arctic shores towards the interior, as may be seen by the courses

of all the great northern rivers, it may be traversed from east to west (if we except the Uralian range) without changing the level more than three or four hundred feet. In Europe, this vast expanse is usually subdivided into the *Germanic plain* on the west, and the *Sarmatian plain* on the east; while in Asia it comprises the *Steppes of Kirghis, Ishim, and Baraba*, on the west, and the *Siberian plain* on the north and east. Each of these subdivisions is necessarily characterised by its own peculiarities of soil, situation, and climate. Thus the Germanic section includes the low-lying polders and morasses of Holland, and the sandy boulder-strewn plains of northern Germany and Prussia—partly under cultivation, and partly occupied by heath and open pasture. The Sarmatian section, on the other hand, consists of cold, swampy, and partially-wooded flats in the north; of moderately temperate, fertile, richly-wooded, and undulating tracts in the middle; and of indifferent grassy steppes and saline deserts in the south. The steppes of Kirghis, Ishim, and Baraba present, as the name implies, wide monotonous tracts covered with rough grass and shrubs during a brief season, but soon converted into arid deserts by the drought of summer, and into bleak, shelterless wastes by the storms of winter. The Siberian plain, as might be expected from its extent, is of a more varied character—consisting of low-lying tundras or earthy peat-mosses, of broad, undulating steppes, and partially-wooded uplands. In summer the tundras are thawed to a small depth, the steppes are scantily covered with grass and mosses, and the banks of the great rivers and uplands are green with the birch and pine; but during the long winter frost and snow reign supreme, and the whole plain is one dreary and inhospitable wilderness.

70. The secondary plains of the Old World, though of minor extent, are not without their decided influences on the physical and vital phenomena of their respective continents. Among those in Europe may be noticed the *plain of France*, comprising the conjoined low grounds of the Seine, Loire, and Garonne; the basin-shaped *plain of Hungary*, watered by the Theiss and Danube, and sometimes spoken of as the "European Pampas;" the *plain of Wallachia* on the lower Danube; the *steppe of Astrakhan*, lying along the Caspian and Volga; and the still more restricted plains of Ireland, Denmark, Andalusia, and Lombardy. Among the secondary plains in Asia may be noticed those of China, Hindoostan, Turan, and the Euphrates. The great *river-plain of China*, said to occupy nearly 200,000 square miles, is alluvial throughout, and still on the increase by the silting up of the Yellow Sea, is crossed in every direction and irrigated by canals, and is the fertile seat of one of the oldest, most industrious, and most numerous populations on the face of the globe. The *plain of Hindoostan*, stretching from the base of the Himalaya to the Deccan, and from the Indus to the Ganges, is also low-lying and alluvial, abundantly fertile under irrigation, and only

light and sandy in the district of "the Thur," on the lower Indus. The *plain or steppe of Turan* extends along the southern shores of Lake Aral westward to the Caspian, and is fertilised by the waters of the Syr and Amoo ; while the historic *plain of Mesopotamia* (Gr. *mesos* and *potamos*, between or in the middle of the rivers) stretches between the Euphrates and the Tigris southwards to the Persian Gulf.

71. In Africa, with the exception of the Sahara or Great Desert, which may be described as a vast expanse of arid sands and shingle, with an occasional oasis of life and verdure, we know of no continuous plains like those of Europe and Asia, though several minor flats and deltas give diversity to its generally unbroken and monotonous seaboard. The surface of the Sahara is not, however, as at one time supposed, a uniform plain of burning and drifting sands, but presents considerable diversity both in composition and altitude. Some portions are low and flat ; others sink, it is said, even below the sea-level ; while many rise in bare, broad plateaux 1000, 2000, and 3000 feet above the surrounding surface. Immense tracts are shingly and saline ; others consist of loose drifting sand ; occasional patches are rocky, and scantily covered with thorny scrub ; while, at distant intervals over the arid waste, some spring or surface retention of water gives birth to an *oasis* or islet of vegetation. This great desert, stretching from Morocco on the west to the valley of the Nile on the east, is prolonged, as it were, eastward and northward through Arabia, Persia, and Tartary, to Mongolia, where it terminates in the equally arid and sandy desert of Shamo—the whole tract from the Sahara to Shamo pointing at once to similarity of conditions and sameness of geological origin. The minor low-grounds of Africa, sufficiently known to geographers, are those on the lower courses of the Senegal and Gambia—Senegambia ; the swampy and jungly delta of the Niger ; and the fertile, periodically inundated delta and valley of the Nile. Respecting Australia—the only other region in the Old World sufficiently extensive to admit of plain-lands—we are yet too scantily informed to offer any definite opinion as to the nature of the interior, though recent discovery would seem to indicate that, instead of a vast scrubby and waterless plain, there is much practicable country—grassy flats, gum forests, and river-creeks, marking, at intervals, the routes of the adventurous explorers.

72. Turning next to the New World, we find the plains on a much more conspicuous and decided scale—stretching more continuously, and rising less above the level of the ocean. In North America, the *Great Central Plain*, lying between the Rocky Mountains on the west and the Alleghanies on the east, extends, it may be said, from the Arctic Ocean to the Gulf of Mexico, a distance of nearly 3000 miles. Throughout this long course it is only once interrupted by the gently swelling prairie-grounds (1500 or 1600 feet in height) that turn the courses of the Red and

other rivers to the north, and those of the Missouri and Mississippi to the south. Varying, in wide expanses, from 100 to 600 and 800 feet in elevation, it presents considerable diversity of surface—swampy marsh, grassy prairie, forest-land, and barren ground. The most notable feature in the surface of this great central plain, which occupies an area of nearly 3,000,000 square miles, and has been characterised by Humboldt as “an almost continuous region of savannahs and prairies,” is undoubtedly that of the open, slightly undulating, and grassy portions denominated *prairies*. These prairies are of vast extent; some are rolling, others are flat and level in surface; many of them are treeless, and covered only with luxuriant grass and flowers; towards the south some tracts verge into shrubby woodland, while in the extreme north the soil is largely swampy and desert. The only other notable lowland in North America is the *Atlantic plain*, lying between the Alleghanies and the Atlantic—a fertile district of little elevation, but often flat and swampy along the seaboard.

73. In South America we are presented with three well-marked and distinctive river-plains—viz., those of the Orinocco, the Amazon, and La Plata. The first is one of the lowest and most level tracts in the world, rising not more than 200 feet at the distance of 500 miles from the sea, and throughout that course marked by swamps, periodically overflowed grass-flats (*llanos*), and tropical forest-growths. The llanos occupy an area of about 160,000 square miles, which is almost wholly inundated during the rainy season, but shortly afterwards is so densely covered with luxuriant grasses that it is known to the natives as the “sea of grass.” During the ensuing tropical droughts these llanos become parched and withered; hence the frequent conflagrations to which they are subject; and hence also, to a great extent, the perpetuation of their treeless character. The second or Amazonian plain, still more extensive and unique, is characterised by similar forest-growths (*selvas*) and grassy plains, and is the largest river-basin in the world, occupying an area of about 1,500,000 square miles. Richly alluvial in soil, periodically inundated, and under the influences of a tropical sun, these *selvas* present the rankest luxuriance of primeval forest-growth, and are in many places accessible only by the river-courses that traverse their areas. The third, comprising the contiguous basins of the Uruguay, Parana, La Plata, and Colorado, embraces an area of not less than 880,000 square miles, and is characterised chiefly by its deep alluvial soil, broad thistly flats, and grassy pastures, known as *pampas*. Stretching from the flanks of the Andes to the shores of Buenos Ayres, and thence southwards to the deserts of Patagonia, these pampas differ considerably in character, being flat and thistly towards the coast, slightly undulating and grassy towards the interior, and full of bogs and swamps and scrubby ridges as we approach the Andes. The whole region is treeless; verdant during the rains, but withered

and parched during the dry season. The desert terrace-land or steppes of Patagonia, extending southward from the Rio Colorado to the extremity of the continent, is a sterile, uninviting region—the soil shingly, and strewn with boulders, the grass stunted, and the climate cold and tempestuous.

Valleys and Minor Depressions.

74. Besides the great plains of the world, there are numerous valleys and minor low-lying tracts that exercise a decided influence on the soil, climate, vegetables, and animals of the countries in which they occur. These are the *dales* and *vales* of existing streams, the broader *basins* of lakes and rivers, and the *straths*, *carsees*, and *deltas* of our estuaries. Whether they have been formed by subsidences of the earth's crust, by the silting-up and drainage of lakes and morasses, or by the slow erosive power of rivers (valleys of erosion), their characters are much alike, and the purposes they subserve identical. Their low-lying situations, their tempered climates, their rich soils, and well-watered surfaces, have ever made them the grand nurseries of vegetable and animal growth; and this fertility and amenity has ever attracted the human race, and rendered them the main theatres of industry and civilisation. To particularise the lesser valleys—the *dales* and *vales* that give amenity and richness to the land surface—would be merely to enumerate the rivers and streams that flow through the different continents. Every stream has, in some part or other of its course, its strip or patch of valley-ground, and these occur of every extent, from the meadow of a few acres to the dale of many leagues, at all altitudes, from the polder and fen protected from the tides by embankments, to the valley high among the mountains, and in every variety of surface—morass, silt, sand, gravel; from the warp of the latest flood to the green sward of a thousand centuries. In civilised countries these low-lands are the principal seats of culture and husbandry; while in semi-civilised regions they constitute the pasture-lands of the nomadic shepherd and herdsman.

75. Besides the *dales* and *vales* and valleys properly so called, there are also the *deltas* of many rivers—low-lying tracts still in course of formation, and which, from their swampy and partially inundated character, can scarcely be said to belong to the domain of the land. To this category belongs much of the *deltas* of the Niger and Nile, the Indus and Ganges, the Mississippi, and the Orinocco and Amazon. The older and higher portions of those *deltas* have long since been converted into fertile alluvial plains, but the lower portions consist largely of mud-flats, sandbanks, and lagoons, partially covered with jungle-growth during the dry season, yet inundated as far as the eye can reach during the periodical overflows of their rivers. In course of time

the existing mud-flats and sandbanks will be converted into alluvial land, and other banks and shoals will arise as the sediments of the rivers are carried farther forward into the area of the ocean. Besides the river-deltas, there are also considerable low-lying tracts of marine silt and sand-drift formed, and still forming, along many of the more sheltered bays and recesses of the ocean. These, like the *links* of our own islands, the *landes* of France, the *dunes* of Holland and Denmark, the *fens* of England, and the *swamps* of Florida, are gradually on the increase, and in process of time assume a flat or plain-like surface.

76. As there are plains and valleys at considerable elevations, and others again scarcely raised above the overflow of the tides, so there are several areas depressed even beneath the general level of the ocean. Laying aside some tracts of the Sahara or great desert of Africa, which are said to sink beneath the sea-level, the most remarkable of these ascertained depressions are the Aralo-Caspian basin and the trough of the Dead Sea. The former of these, in which are situated the Caspian and Aral seas, is a depression of 162,000 square miles in extent, all considerably below the ocean—the surface of the Caspian, the lowest portion of the cavity, being actually 83 feet beneath the level of the Black Sea. The latter, though a mere trough in comparison with the former, is still more remarkable for its depression, the Dead Sea being no less than 1312 feet below the Mediterranean, from which it is separated by the mountain-range of the Lebanon. Were it not that the river-supply of these cavities is fully counterbalanced by evaporation, they would, in process of time, become filled with water, and their surplus find an outlet to the ocean—the Aral and Caspian to the Euxine, and the Dead Sea to the Gulf of Akaba in the Red Sea, with which at some former period it was evidently connected by the dry stony valleys or “wadies” that now lie between.

RECAPITULATION.

In the two preceding chapters we have endeavoured to present an outline of the superficial features of the land, as composed of mountains and table-lands, of plains and valleys, and the relations they bear to each other in the scheme of Physical Geography. Whatever be the geological law that regulates the successive upheaval and submergence of large tracts of the earth's crust, we see in the arrangement of the present continents the more violent effects of the earthquake and volcano in producing abrupt and mountainous irregularities, and the more gradual efforts of air and water in moulding into uniformity of surface its plains and valleys. Whatever be the nature of the vulcanism

acting from within, we see that it exerts itself in certain definite lines and centres, and these lines and centres give rise to mountain chains and groups, and these again give contour and configuration to the terrestrial areas. In the Old World, as already stated, the main axis of elevation is from east to west; hence the greatest length of the continent in this direction, and hence also the determination of the rivers and river-plains in northerly and southerly courses from this axis. In the New World, again, the main line of elevation is from south to north; hence also the corresponding direction of that continent, and the opposite courses of its principal rivers. As with the main continental masses, so with the minor spurs and peninsulas—their direction of greatest length being invariably regulated by the direction of their hills. Geology thus becomes the evident groundwork of Geography; and on reference, for instance, to the Geological Map of Europe (Plate II. of Johnston's 'Physical Atlas'), it will be seen that all the granitic, trappean, and volcanic outbursts are but expressions, in other terms, for the extent and directions of the mountains of that continent. As the great plains are but the undisturbed portions of the continents, they will naturally take the same main direction as the mountain-ranges; just as the table-lands, which are upraised by the mountains, will be situated among their groups or along their axes. On the other hand, the rivers, obeying the laws of descent from the opposite sides of these axes, will have their valleys and dales running less or more in cross courses, thus giving additional features of diversity to the land. In the Old World the principal mountain-chains are those of the Pyrenees, Alps, Carpathians, Caucasus, Himalayas, Thian-Shan, and Altai, which trend in one main direction from the Atlantic on the west to Behring Strait on the east; and this great irregular axis is flanked on the north by the plains of central Europe and Siberia, and on the south by the minor plains of Hindostan and China. In the same manner, in the New World, the Rocky Mountains and Alleghanies alternate with the central and Atlantic plains of North America; while in South America the great lowland may be said to stretch between the Andes and the Cordilleras of Brazil, the only exception being the river-valley of the lower Amazon. Mountains and table-lands, plains and valleys, are thus but the counterparts of each other; and rugged and inhospitable as the former may seem, the latter would be but thirsty deserts were it not for the clouds and rains, the streams and rivers, that are generated among their summits, and descend in perennial supplies from their glens and recesses. Ethnologi-

cally, as mountain regions have ever been the nursing-fields of hardy, brave, and independent races, so plains and valleys have ever become the sheltered and fertile seats of settled industry and civilisation. The plains of China and Hindostan, of the Tigris, the Euphrates, and the Nile, were the early and populous abodes of mankind in the eastern hemisphere, just as in the western the valleys of the Missouri and the Mississippi were the chosen grounds of primitive mound-building races. And as in former ages, so even now the principal sites and centres of industry are to be found in the river-plains of the Old and New Worlds—the causes that induced the early shepherd settlers being equally operative on their agricultural, city-dwelling, mechanical, and commercial descendants.

VIII.

THE WATER—ITS OCEANS AND SEAS.

Their Area and Configuration.

77. HAVING considered the various conditions of the Land—its area, configuration, highlands, and lowlands—we now turn to those of the Water, as exhibited in its oceans and seas, their areas, depths, composition, tides, currents, and kindred phenomena. And here it may be observed of WATER, which forms so important an element in the constitution of the globe, that, chemically speaking, it is the *protoxide of hydrogen*, consisting of two volumes of hydrogen and one of oxygen, or of eight parts of oxygen to one of hydrogen by weight—88.9 oxygen and 11.1 hydrogen. When pure and at ordinary temperature, it is fluid and amorphous, without taste or smell, colourless in small quantities, but in large masses of a peculiar bluish-green or blue. The specific gravity of pure or distilled water, at 62° Fahr., is assumed at 1.000, and is taken as the standard of gravity for all other bodies; but sea-water varies, according to locality and the depth from which it is taken, from 1.027 to 1.029. When heated to the temperature of 212° Fahr. at the level of the sea, and under the ordinary pressure of the atmosphere, water boils, and is converted into steam; and this *boiling point* (as it is termed) becomes less as we ascend above the sea-level. At 40° Fahr. water is at its minimum volume, expanding as it rises above that temperature, till it is wholly converted into vapour, and also, as it falls below it, till at 32° for fresh, and 28½° for salt water, it is converted into *ice*—a transparent, brittle, crystalline solid, which floats on the surface. Water, as found on the earth, is never absolutely pure, but contains more or less of various substances, as atmospheric air, carbonic acid, nitrogen gas; silica, alumina, and salts (carbonates, sulphates, nitrates, phosphates) of lime, magnesia, soda, potash, protoxide of iron, manganese; or chlorides and fluorides of their metallic bases; and in the sea and some saline springs, also iodine and bromine. Like all other fluids whose particles are free to arrange themselves, water at rest always assumes a level surface, and this surface, in the case

of the ocean, corresponds with and forms part of the circumference of the globe. As an agent in nature, it is indispensable to the life of plants and animals, it enters into the composition of all bodies, whether organic or inorganic, and in the form of rain, streams, rivers, waves, tides, and currents, is the great modifier and remodeller of the geological aspects of the globe.

78. It has already been noticed (par. 36), that though encircling the globe on every side, and spreading over nearly three-fourths of its surface, the great "world of waters" is more or less configured into certain expanses which are termed *oceans*, and thus we have on the west of the Old World, and between it and the new, the *Atlantic Ocean*; while on the west of the New World, and between it and the Old, spreads out the still vaster area of the *Pacific*. These divisions become apparent on the most cursory inspection of the map of the world—the former lying like an irregular valley between the two continental land-masses, and communicating freely with the arctic and antarctic waters; the latter narrowed to a mere strait on the north, but spreading out towards the south over nearly half the globe, and ultimately losing its individuality in the undefined expanse of antarctic waters. Besides the Atlantic and Pacific, the *Arctic* and *Antarctic* constitute well-recognised though imperfectly known oceans; while between Africa and Australasia stretches the familiar and much-traversed area of the *Indian Ocean*. In treating of these great oceanic expanses, various names and subdivisions are employed by navigators, but for all practical purposes in physical Geography the terms North and South Atlantic, North and South Pacific, Arctic, Antarctic, and Indian Oceans, are sufficiently explicit and comprehensive. Or, looking upon the waters that extend southwards from the extreme points of Africa, Australia, and South America, as one great united mass, the term *Southern Ocean* will often be found to be convenient, and not inappropriate.

79. Taking the Atlantic Ocean as extending from the arctic to the antarctic circle, its length is upwards of 9000 miles; its breadth varies from 900 to 4000 miles, being only 900 between Norway and Greenland, 1700 between Sierra Leone and Brazil, and 4100 between Morocco and Florida; and its computed area is about 25,000,000 square miles. This vast expanse is little interrupted by islands; in its northern section it is irregular in form, and throws several important branches into the land; but in the southern its form is regular, and its shores continuous. Towards the north it is partly enclosed by the rocky coasts of Greenland, Iceland, and Norway; but towards the south it is quite open, and merges broadly into the Antarctic Ocean. The leading branches are Baffin and Hudson Bays, the Gulf of St Lawrence, Bay of Fundy, Gulf of Mexico, and the Caribbean Sea, on the west or American side; and on the east or Old World side, the North Sea, Baltic Sea, English Channel, Bay of Biscay, the

Mediterranean Sea, and the Gulf of Guinea. All, or nearly all, of these recesses occur in the northern division of the Atlantic; hence the greater interest of this section to the geographer, naturalist, and navigator. Of the minor seas (some of which are ice-locked for a considerable portion of the year, and others encumbered by reefs and shoals), the most important, physically and vitally, is the Mediterranean, whose shores formed the early nurseries of civilisation and commerce, and whose waters are still the highway of communication, not merely between the three continents—Europe, Asia, and Africa—that encircle its shores, but between these and every other portion of the globe.

80. The Pacific Ocean, though less important as a highway of commerce, occupies nearly twice the expanse of the Atlantic—its greatest breadth being 12,000 miles, and its computed area about 50,000,000 square miles. Unlike the Atlantic, it is almost entirely shut out from communication with the Arctic Ocean—the only passage of connection being that of Behring Strait, not more than 36 miles in width, with a maximum depth of 25 fathoms; but, like the Atlantic, it also opens out towards the south, and merges undefinedly into the Antarctic. It is thickly studded with islands and clusters of islands, and these physically and vitally constitute one of its most distinctive features. Its leading branches are the Sea of Kamtchatka, Sea of Okhotsk, Sea of Japan, Yellow Sea, and Chinese Sea, on the west or Asiatic side; while on the east or American side, the Gulf of California and the small bay of Panama are the only indentations that break the uniformity of its coasts. The most important of these minor seas are those of Japan and China, whose shores have been the seat of an early and peculiar civilisation, and whose waters have long been traversed by the ships of every other nation.

81. The Indian Ocean, stretching between Africa and Australia on the one hand, and between Asia and the Southern Ocean on the other, is upwards of 4000 miles in breadth, and is computed to have an area of about 17,000,000 square miles. If we except the Indian Archipelago, which forms its boundary rather than belongs to it, it is encumbered by few islands; and it also penetrates the land by few branches—the Red Sea, Arabian Sea, Persian Gulf, and the Bay of Bengal, being the only minor seas—and these all on its northern or Asiatic boundary. The most important of these minor branches are the Red Sea and the Bay of Bengal—the former early and intimately connected with the history of man, and the latter, the leading highway of modern commerce to the varied wealth of India.

82. The Arctic and Antarctic Oceans, from their circumpolar situations, are largely blocked up with ice, and consequently but imperfectly known to geography. The Arctic forms, as it were, a circular basin, bounded in general by the northern coasts of Europe, Asia, and America, which remarkably conform to the parallel of 72°, and having an area roughly estimated at 4,000,000

square miles. It penetrates northern Europe by the White Sea and Sea of Kara, and northern Asia by the Gulf of Obi and a few small inlets, and northward from these shores seems interrupted by comparatively few islands. The northern shores of America, however, present so many islands and ice-locked inlets, that it has, up to the present moment, been impossible to determine whether land or water continues northward and surrounds the pole. The Antarctic Ocean, on the other hand, is open on all sides to the Pacific, Atlantic, and Indian Oceans, which thus insensibly merge themselves into the great Southern Ocean. So far as navigators have ventured to approach the southern pole, various islands and shores have been observed which would favour the idea of a circumpolar continent; but whether land, sea, or an ice-bound archipelago, occupies the immediate region of the pole, is likely long to remain an undetermined problem. Altogether, the Antarctic is a cold, boisterous, and unapproachable ocean—its ice extending 10° nearer the equator than that of the Arctic, and offering few of those inducements that have stimulated repeated research in the Northern Ocean.

83. Such are the areas occupied by the waters of the ocean—areas and subdivisions which are not only necessary to intelligible description, but which are marked in reality by different conditions and characteristics in nature. Position on the globe decides their temperature; area and configuration determine their tides and currents; and the sum of these physical conditions regulates the nature and distribution of their plants and animals. Any change, therefore, either in position, area, or configuration, would be attended by a corresponding change of conditions, and any such alteration would affect all the consequences, physical and vital, that depend on external conditions. As they exist, the North and South Atlantic, situated under different latitudes, enjoy different temperatures; while the north, by virtue of its greater irregularity of form, and numerous ramifications into the land, exhibits a much more varied display of vegetable and animal life. The same also holds good of the North and South Pacific, with these important modifications, that the North Pacific, compared with the North Atlantic, is almost excluded from Arctic influences, while the numerous islands of the South Pacific occasion conditions, physical and vital, peculiarly its own. The Indian Ocean, surrounded on three sides by land, and situated, for the most part, in the torrid zone, presents peculiarities unknown in other subdivisions; while the Arctic and Antarctic, receiving the minimum of solar heat, are ice-locked for the greater part of the year, and have little in common with the other sections of the ocean. As already stated, it is on these primary relations of position and configuration that the different temperatures, tides and currents of the various oceanic subdivisions, depend; and it is entirely owing to these conditions that the

specific life of one sea or ocean differs from the life of all other seas and oceans. And yet it must be borne in mind that, though diversely situated and characterised, there is still the most intimate connection and interchange between their waters—the colder ever flowing towards the warmer, and the warmer towards the colder, so that in this respect they constitute in reality one great and indivisible “world of waters.”

Composition, Density, Depth, Temperature.

84. This great ocean, in all its areas and ramifications, is characterised by a greater or less degree of *saltiness*—this saltiness arising from the presence of certain substances held in chemical solution in its waters. These substances are chloride of sodium (common salt) and sulphates of magnesia and lime, together with minor and varying proportions of salts of potash and ammonia, iodides and bromides of sodium, carbonate of lime, silica, &c.—amounting in all from $3\frac{1}{2}$ to $4\frac{1}{2}$ grains to the hundred of water. According to M. Regnault, the following is the mean of several analyses of sea-water :—

WATER,		96.470
SALINE INGREDIENTS, 3.505	{ Chloride of sodium, . . .	2.700
	{ Chloride of magnesium, . .	.360
	{ Chloride of potassium, . .	.070
	{ Sulphate of lime,140
	{ Sulphate of magnesia, . .	.230
	{ Carbonate of lime,003
Loss (including iodides, silica, &c.),	{ Bromide of magnesium, . .	.002
		.025
		<hr/> 100.000

The preceding ingredients may vary in different seas, and according to the locality whence, and the season when, the water is taken, but only to the extent of a fractional percentage—the incessant circulation and intermingling of the ocean's waters by waves, tides, and currents, producing a uniformity, or all but uniformity, in its saline composition. It has been found, however, that the waters of the Southern Ocean are slightly saltier than those of the Northern ; that the greatest saltiness takes place along the parallels of 22° north and 17° south, or in the courses of the trade-winds, which absorb and carry off an excess of evaporation towards the equator, where it descends in freshening rain-falls ; and that from these limits of maximum saltiness there is a slight progressive diminution towards either pole. Though communicating freely (by currents and counter-currents) with the ocean, the majority of inland seas are less salt than the ocean, in consequence of the influx of rivers into their circumscribed areas ; but some, like the Red Sea, receiving no rivers, and subjected to

active evaporation, have their saltness slightly in excess. As a general rule, inland seas receiving numerous rivers, and from their situation subjected to little evaporation, will be fresher than the ocean (*e. g.*, the Baltic); while others also receiving rivers, but subjected to a more active evaporation (as the Mediterranean), will have their saltness somewhat in excess. Though the saltness of the sea be pretty uniform at great depths, still at the surface, owing to the admixture of rain, river, and iceberg water, it is not quite so salt; and this freshness will increase, of course, according to proximity to the mouths of the entering rivers. It has also been ascertained that water from the surface contains less air than that from depths, and the difference may equal one-hundredth of the volume of water. Another noticeable property of salt water is, that it is less sensitive, if we may so speak, to cold than fresh water—the latter freezing, as is well known, at 32° , while sea-water is not converted into ice till the thermometer sinks to $28\frac{1}{2}^{\circ}$ Fahr. It is likewise less vaporisable than fresh water—that is, a given extent of salt-water surface gives off less vapour during the same time, and under the same conditions, than an equal extent of fresh-water surface. Such composition and properties are no doubt all-essential in the economy of nature. Shell-fish, crustacea, coral-zoophytes, and other creatures, derive the calcareous matter of their structures from the salts of the ocean; fishes breathe the aerated waters of the sheltered and undisturbed depths; and both plants and animals obtain conditions of existence which absolutely pure water would fail to supply. By its lower freezing point a larger amount of surface is ever kept open and accessible; and by its slower evaporation a less amount of moisture is borne from its greater expanse to the comparatively smaller surface of the land.

85. It will be seen that no notice has been taken in the preceding paragraph of ingredients—sand, mud, and organic debris—that may be *mechanically suspended* in the waters of the ocean. These are purely local and accidental, depending on river-floods, tidal-currents, waves, storms, and other commotions. When the commotion subsides, the waters regain their transparency; and altogether, unless along wasting shores, in tidal estuaries and river-embouchures, there is really very little matter mechanically suspended in the waters of the ocean. On the other hand, the ingredients held in *chemical solution* are all but constant and universal. The water that evaporates from the ocean is all but absolutely pure; it falls on the land in mist and rain and snow; percolates the soil and rocks; and returns again to the ocean, carrying with it the saline substances it has dissolved from the rocky strata. The ocean is thus the great equalised repository of all that is borne from the continents; and there they would accumulate, were it not for the beautiful counterpoise that is ever kept up by the requirements of plants and animals, as well as by the intervention of new

chemical arrangements among its multifarious sediments. So far as Geology can determine by a study of the marine life (shell-fish, corals, foraminifera, &c.) of former ages, the composition of the ocean seems to have been much the same as it is now; and thus, in all our reasonings, we may regard its saline contents as having long arrived at a state of equilibrium and fixity. Even if there were a slight excess at any one period, that excess would be merely temporary, as those incessant mutations of sea and land, involving the formation of new limestones, magnesian limestones, rock-salts, and the like, are ever taking up the surplus and restoring the equilibrium.

86. *The mean specific gravity of sea-water*, as compared with absolutely pure water at 62° Fahr., is found to be 1.0275 — an amount that corresponds to a percentage of 3.505 of saline ingredients. The salter, therefore, that water is, the greater its gravity; and hence the fresh-water of rivers, of melting icebergs, &c., will float for many miles on the surface of the sea before the two fluids be thoroughly diffused and commingled. It is owing to this that potable water has been *skimmed* from the surface several miles from the mouths of large and rapid rivers; and it is also for this reason of unequal densities that currents are established in different parts of the ocean—the heavier ever seeking to establish its equilibrium. As already mentioned, water acquires its minimum volume, or greatest density, at a temperature of 40°, and becomes lighter and lighter either as it rises above or falls below this temperature. Owing to this property a perpetual interchange or circulation is kept up among the waters of the ocean, *horizontally* from colder to warmer, and from warmer to colder regions; and *vertically* from surface to depths, and from depths to surface. Again, water being slightly compressible, it follows that at great depths in the ocean the water will be denser than at the surface, and consequently what takes place near the shore will be impossible at extreme depths. According to experiment, water at the depth of 1000 feet is compressed $\frac{1}{115}$ th of its bulk; and at this ratio the pressure at the depth of one mile would be equivalent to 160 atmospheres, or 2320 lbs. on the square inch; while at the depth of 4000 fathoms, or about $4\frac{1}{2}$ miles, it would amount to 750 atmospheres! At vast depths, therefore, it is generally supposed that vegetable and animal life, as known to us, could not possibly exist; and though some recent soundings in the north seas at the depth of 1260 fathoms would seem to oppose this opinion, yet the paucity and uncertainty of these trials leave the question still in doubt; and we may, in the mean time, adhere to the general belief that the extreme depressions of the ocean, like the extreme elevations of the land, are barren and lifeless solitudes.

87. Touching the *depth of the ocean*, it has been already observed, that as the dry land rises variously and irregularly above the level of the ocean, so the bottom of the ocean sinks variously

and irregularly beneath its waters. All that has been learned from the soundings of navigators establishes the fact, that there are shallow shoals and banks, deeper flats and plateaux, and still deeper troughs and valleys ; and that were the whole laid dry, we should have presented to us much the same kind of inequalities as are presented by the surface of the land. In fact, the bed of the ocean is but the submerged surface of former lands, and unless, perhaps, in the instance of coral-reefs and submarine volcanoes, there is no foundation for the belief that the troughs and precipices of the ocean are sharper and more abrupt than those of the dry land. If the inequalities of the land are worn and rounded by meteoric agency, and masked by a covering of soil, so the inequalities of the ocean-bed were worn down before submergence, and have since been masked, except in the courses of tides and currents, by a still deeper covering of sediment and drifted debris. As a general fact, it may be stated that, where the land slopes gradually towards the ocean, the waters also deepen gradually ; and, on the other hand, where the land descends abruptly, the sea deepens, in like manner, suddenly and abruptly. In fact, this is only the natural consequence of slope and counter-slope, and a great relation to which there is scarcely an exception. The northern plains of Russia and Siberia, for example, slope gradually into the shallow bed of the Arctic Sea, just as the abrupt terminations of South America, Africa, and Australia, dip suddenly into the deeper waters of the Southern Ocean. The level plains of China spread gently outwards into the shallow waters of the Yellow Sea ; the low shores of eastern England, in like manner, slope slowly into the comparatively shallow basin of the North Sea ; while, on the other hand, the precipitous coasts of Norway dip suddenly down into a corresponding depth of water. This fact, that low lands are generally bordered by shallow seas and high lands by deeper water, affords no idea, however, of the depths of distant and central expanses, and for these we must either appeal to theoretical deduction or to actual observation.

88. So far as experiment is concerned, comparatively little is known of the absolute depth of the ocean ; and even where deep soundings have been made, there has been great liability to error, partly from the imperfection of the apparatus employed, and partly from the chance of the line being deflected from the perpendicular by the force of under-currents. The common notion that the extreme depths of the sea correspond to the extreme heights of the land—that is, as the highest mountains rise little above five miles, so the greatest depths sink little below that amount—has no foundation in fact, there being no necessary connection between the two phenomena. The *mean elevation* of all the land—continents and islands, mountains and plains—has been estimated by Humboldt at somewhat less than 1000 feet ; and the *mean depth* of the ocean has been calculated by Laplace,

from tidal waves and kindred phenomena, to be at least 21,000 feet, or about four English miles. We know, however, that a very large proportion of the ocean is comparatively shallow, and not a tithe of this depth, and therefore, to make up the mean, some other portions must be proportionally deeper, and to the extent, it may be, of eight or ten miles. Indeed, soundings (no doubt open to question) have been made in the South Atlantic, both by British and American navigators, varying from 27,000 to 46,000 feet; and soundings perfectly reliable have been taken in the North Atlantic, off the bank of Newfoundland, to the depth of 25,000 feet; while from calculations on the velocity of tidal waves, which are found to proceed according to the depth of the channel, it has been estimated that the extreme depths of the same ocean are about 50,000 feet, or more than nine miles. Altogether, and according to our present information, we may admit a mean depth of four miles for the ocean, and reliable soundings to the extent of five miles; but believe, partly on experiment, and partly on theoretical grounds, that several portions sink to the depth of eight or ten miles. It may also be stated in general terms that the Atlantic, averaging from three to five miles, is, on the whole, deeper than the Pacific; that great depths (from four to six miles) have been determined in the Indian and Southern Oceans; that the Antarctic becomes shallower as we approach the pole; and that the Arctic, of moderate depth, is characterised by great irregularity and diversity. With regard to the minor seas, the greatest ascertained depth in the Mediterranean is about 13,000 feet; in the Red Sea, 6300 feet; in the Baltic, 840 feet; in the Caribbean Sea, 14,000 feet; and in the Gulf of Mexico, about 8000 feet.

89. Respecting the *temperature of the ocean*, few reliable or sufficiently extended observations have yet been made, either as regards its various areas or its successive depths. We know, however, that it is more equable than that of the land, and that, though the superficial portions are colder in summer than the surrounding atmosphere of any contiguous terrestrial district, they are in winter always several degrees higher—thus exercising the function of a great storehouse of heat for modifying and equalising the climates of the adjacent land. The surface temperature is necessarily highest along the equator, or rather along a belt, varying from 2° to 8°, on either side of the equator, and then gradually diminishes as we approach either pole. Along this equatorial zone temperatures have been found ranging from 78° to 85°—higher exceptional temperatures (87° and 88°) having occasionally been taken in parts of the Indian Ocean and the Gulf of Mexico. Though varying in *surface temperature* according to latitude—from 80° at the equator to perpetual ice towards either pole—it has yet been found that at very great depths the ocean preserves a uniform *mean temperature* of about 39½°. Thus, according to the experiments of Sir John Ross, the circle of the mean temperature of the ocean in the southern

hemisphere lies between the 56th and 59th parallels of latitude ; along which belt the uniform temperature of $39\frac{1}{2}^{\circ}$ has been found to prevail at all depths, from the surface downwards. To the south of this line, owing to the absence of solar heat, the surface depths are colder, and the mean of $39\frac{1}{2}^{\circ}$ is not reached in the 70th parallel till we descend to the depth of 4500 feet, beneath which to the greatest depths the temperature is uniformly at $39\frac{1}{2}^{\circ}$, while the surface temperature is only 30° . To the north of the line of mean temperature, in consequence of the absorption of the sun's heat, the surface depths are warmer ; and in the 45th parallel the mean of $39\frac{1}{2}^{\circ}$ is not reached till we descend 3600 feet ; while at the equator we have to descend 7200 feet before the same mean is obtained, and then at all depths below this it maintains the unvarying mean temperature of $39\frac{1}{2}^{\circ}$, though the surface is at 80° . Presuming that a similar order prevails in the northern hemisphere, we have thus three great regions of oceanic temperature—an equatorial and two polar, the former characterised by warm water at the surface, and the latter by cold. Such are the facts, so far as ascertained, respecting the general temperature of the ocean ; but inland seas and currents may be colder or warmer according to the position they occupy and the direction from whence they proceed. The surface temperature of the Gulf of Mexico, for example, is several degrees warmer (86° and 88° have been named) than the main Atlantic under the same latitude ; the waters of the Gulf Stream are also several degrees higher in a large portion of its course than those through which it flows ; while the Arctic Current, on the other hand, is considerably colder.

90. Besides the preceding conditions of saltness, density, depth, and temperature, there are also those of *colour* and *luminosity*, usually adverted to by navigators and geographers. In small quantities water is generally regarded as colourless, but that of the ocean assumes different hues, and this altogether independent of the colours of the sky which may be mirrored on its surface. Thus, in the open ocean, shallow water is indicated by a *green tint* of different degrees, while profound depths are characterised by an *indigo blue*. Whether this arises from the greater density of the mass, or from some peculiarity of its saline constitution, is not yet satisfactorily known ; but the fact remains, and “sea-green waves” and “dark-blue oceans” are something more than mere poetic fancies. Of course, in some localities there may be accidental or even permanent discolourations, arising from the entrance of river-water, from peculiarity of bottom, or from the presence of countless myriads of vegetable and animal organisms ; hence the application of such terms as Red, Black, White, Yellow, Green, and Vermilion to certain seas and areas of the ocean. In general, however, the ocean water is clear and limpid, and, under favourable circumstances, objects are reported to have been seen at a depth of 300 and 400 feet, or about half the dia-

tance to which the sun's light is supposed to penetrate into the abyss of waters. The phenomenon of luminosity or phosphorescence is less general, perhaps, and seems to depend in a great degree on locality, season of the year, and state of the weather. Luminous animalcules and creatures of various kinds appear to be the proximate cause of the phenomenon, which becomes more apparent in still, dark nights, and where the surface of the water is disturbed by the stroke of an oar, or the friction of a passing keel.

Waves, Tides, Currents.

91. Like other fluids whose particles are free to yield to every impulse, the waters of the ocean are subject to several movements, the more important of which are waves, tides, and currents. Waves are produced by winds, and occasionally by earthquake commotion, tides by the attractions of the moon and sun, and currents chiefly by that incessant tendency which waters of different densities and temperatures have to assume a state of equilibrium. And first of WAVES, whose main characteristics are magnitude, velocity, and force of impact. Waves occur in every part of the ocean, and wherever the wind blows,—the aerial current drifting the surface waters along with it, and producing undulations, which increase, according to the power of the propelling force, from the gentlest ripple to billows 30 or 40 feet in height. In deep and open seas a continuous wind produces merely an undulation or up-and-down movement of the surface waters; and this commotion, even in the case of a wave a quarter of a mile in breadth and 40 feet high, is not sensibly felt at a depth of 220 fathoms. But in obstructed and shallow seas the lower part of the advancing undulation is retarded by frictional contact with the bottom—the upper portion advances with headlong motion, and ultimately breaks with forcible impact on the opposing shore, which is worn and abraded by the backward and forward motion of the surf. Such is the beginning, course, and termination of ordinary waves,—at first a mere ripple; as the gale increases, a long roll and swell in the deep sea; and ultimately a cresting and dash of breakers on the shelving shore. Occasionally, however, the wind shifts, and sets in waves from opposite directions, and these crossing and commingling produce a violent commotion, even far out at sea, and in the deepest waters. The aspects and characters of waves are known to sailors by many different names; the ruffle or ripple under a rising breeze being spoken of as a *catspaw*, the long undulation as a *swell* or *billow*, the shorter undulations as they approach the shore as *rollers* and *breakers*, and the broken water along shore as *surf*. The big heavy waves that occasionally set in when there is no wind (having been produced by storms far out at sea), are said to form a *ground-swell*; the commotion produced by


cross-waves forms a *chopping sea*, and in a less degree a *jabble* or *cross-lipper*; but these and similar terms belong more to nautical technicality than to the generalities of Physical Geography.

92. Generally speaking, the magnitude of *wind-waves* has been greatly exaggerated, partly from the difficulty of making correct observations, and partly from the impression of dread produced on the mind of the observer. The greatest waves known are said to be those off the Cape of Good Hope, where, under the influence of a north-west gale, they have been found to exceed 40 feet in height. Off Cape Horn they have been measured at 32 feet from trough to crest; and in the North Atlantic, waves from 20 to 25 feet are by no means uncommon. In our own seas, however, they rarely exceed 8 or 10 feet; and all accounts of their running "mountains high" must be received as mere poetical exaggerations. In the case of *earthquake-waves* the conditions are altogether different; and as the whole mass of water is then thrown into commotion by sudden and abrupt risings, fallings, and whirlings, waves, or rather walls of water, 60 or 80 feet high, may be thrown with tremendous impetus upon the land. In the Lisbon earthquake of 1755, the destructive wave that rolled in upon the coasts of Portugal was estimated at 60 feet; and in the Simoda (Japan) earthquake of 1854, three huge waves, at intervals of a few minutes, rushed into the bay, destroying the native craft, and completely submerging the town of Simoda.

93. The velocity of waves depends primarily, of course, upon the power and continuance of the wind, but is greatly modified by, and bears an ascertainable relation to, their magnitude and the depth of the water over which they travel. Thus it has been calculated by Professor Airy that a wave 100 feet in breadth and in water 100 feet deep travels at the rate of about 15 miles per hour; one 1000 feet broad and in water 1000 feet deep, at the rate of 48 miles; whereas another, 10,000 feet in breadth and in water 10,000 feet deep, will sweep forward with a velocity of not less than 154 miles per hour. The force with which a wave strikes against any opposing barrier depends, in like manner, upon its bulk and velocity, and in the case of huge waves this impact is enormous. From experiments made at lighthouses and breakwaters their effective pressure has been estimated as high as 6000 lb. per square foot: and one has only to observe the breaches occasionally made in sea-walls, and the distance to which blocks of stone, several tons in weight, have been hurled forward, to be convinced of their great propulsive power. Of course the force with which a wave simply *strikes* is not to be altogether estimated by its propulsive power, for substances submerged in water lose a certain portion of their weight, which greatly facilitates their displacement and transport.

94. The next, and perhaps the most important and persistent of oceanic movements, is that of the *TIDES*—a term applied to


the periodic rising and falling of the waters, occasioned chiefly by the attraction of the moon, but partly also by that of the sun. In obedience to the universal law that "every particle in nature attracts every other particle with a force inversely as the square of the distance," the earth is attracted by the sun and moon, but more by the latter in proportion to its greater proximity. Land and water alike experience this attraction, but the particles of the latter being free to move among themselves, the mass of the ocean is drawn out beyond its normal circumference towards the attracting bodies. Had the earth been immovable as regards the sun and moon, this bulging out of the waters would have been stationary; but as she turns on her axis, meridian after meridian is brought directly opposite to the attracting force, and thus the rising of the waters becomes a great *tidal wave* or *flow* that travels round the globe. The moon, we have said, exercises the greater attraction, but when the sun and moon are in conjunction, or in opposition (that is, at new and full moon), the sum of the two attractions will cause the greatest possible rise, known as *spring-tides*; and when the moon is in her quadratures (that is, at her first and last quarters), the sun's attraction, acting in a different direction, will diminish the lunar tide, and then we will have the least rise, or *neap-tides*. The following diagram may assist the comprehension of these phenomena:—



Here t being the nearest point of the Earth's surface to the moon, the waters at that part are most attracted towards that luminary, and of course rise highest; while on the opposite side at t' the earth is drawn, as it were, away from them, and they stand out nearly at the same height as those at t . But as the waters rise simultaneously at t & t' they are drawn away from e & e' , and as the earth turns round, each point on its surface will necessarily have two high-waters and two low-waters per day. In other words, the sea *flows* or rises as often as the moon in her apparent circuit passes the meridian, both the arc above and the arc below the horizon, and *ebbs* or falls as often as she passes the horizon, east and west. The solar day, however, being only 24 hours, and the lunar (owing to the moon's monthly course round the earth) being 24 hours, 54 minutes, it requires rather more than a rotation of the earth to bring the same meridian to the same position, relatively to the moon, as it had the preceding day. In other words, it requires more than 24 hours to bring the moon round to its vertical position over any given place, and thus the tides of one day are always about an hour later than they were on the preceding day. Again, had the moon been the sole attract-

ing body, the tides would have risen always to the same height ; but the sun, exerting a simultaneous attraction either along with or against that of the moon, creates an alternate maximum and minimum of flow. Thus when the sun is at S and S' his attraction is combined with that of the moon, and a higher tide is the result. When at S the darkened side of the moon will necessarily be towards the earth, and it is new moon ; and when at S' the illuminated face of the moon will be towards the earth, and it is full moon ; so that the higher or *spring-tides* take place alternately at new and full moon. On the other hand, when the moon is 90° from the sun's place (that is, when she is in her first and last quarters, or half moon), his attraction, being exerted at right angles, counteracts that of the moon, and the result is the lower or *neap-tides*. The greatest tides occur, of course, when the luminaries are nearest and pass most vertically to the place of observation ; and as each tide has only about six hours to *flow* and about six hours to *ebb*, the highest must necessarily be the *swiftest* and the lowest the *slowest*.

95. Such, in general terms, is the theory of the tides ; and had the surface of the globe been entirely covered with water, the tidal wave would have been regular and continuous from meridian to meridian, and, as a consequence, highest in the region of the equator, and gradually falling away towards either pole. But the continuity of the ocean being interrupted by land, and this land lying in a great measure meridionally, as well as being irregular in outline, and consisting in many parts of islands, the course of the tidal flow is obstructed and deflected into various courses. Under the present arrangement of sea and land, these courses are, however, sufficiently persistent ; and thus their directions, times, velocities, and heights, can be determined with accuracy for the purposes of navigation. The Southern Ocean, encircling the globe, and being comparatively uninterrupted by land, may be regarded as the area in which the tidal wave receives its great primary impulse. It is thence carried forward, deflecting itself northward into the Indian, Atlantic, and Pacific Oceans, where, uniting with the minor tide-waves generated in these expanses, it flows, rises, and subdivides, according to the outline of the coasts, the depth of water, and the obstruction of islands. Notwithstanding the complications arising from these causes, there is still great regularity in the bi-diurnal flow and ebb of the tides ; and by noting the times at which the same high-water reaches different parts of the coast, a series of lines connecting these points may be laid down so as to indicate the course of the tidal wave with great precision. Such a series of lines are termed *co-tidal lines*, or lines of simultaneous tide, and mark the progress of the summit of high-water from its origin in the Southern Ocean to its remotest ramifications in northern waters. We say northern waters, for though the primary and normal direction of the tidal wave is from east to west,



in obedience to the apparent course of the sun and moon, yet, on entering the troughs of the Indian and Atlantic Oceans, it is compelled to assume a northerly course in accordance with the configuration of these seas. Thus, the new or full moon high-water that passes Van Diemen's Land every morning at twelve, takes twelve hours to reach Ceylon, and thirteen to reach the Cape of Good Hope; in another twelve hours it has passed up the Atlantic, and arrived at Newfoundland; at the end of the third twelve it has rounded the north of Scotland, and is opposite to Aberdeen; at the fourth twelve, or at midnight of the second day, it is opposite the mouth of the Thames; and it is "not till the morning of the third day that this wave fills the channel of the Thames, and wafts the merchandise of the world to the quays of the port of London." (See Map of *Co-tidal Lines*.)

96. The tides, we have said, may be regarded as taking their rise in the uninterrupted expanse of the Southern Ocean. As the wave proceeds westward, it is deflected northward broadly into the Indian Ocean, rapidly and deeply into the larger channel of the Atlantic, and slowly and feebly into the Pacific, where its course is obstructed by numerous islands and coral-reefs. The *velocity of the tidal wave* depends primarily on the conformation and depth of the ocean—proceeding with the greatest rapidity where the ocean is freest and deepest. As the co-tidal lines are laid down at hourly distances, they afford a pretty correct estimate of the tidal velocity—the wider the lines (that is, the greater the distance travelled over in one hour) the greater the speed, and the closer the lines the slower the rate of progress. In the free depths of the Southern Ocean, this velocity may equal 1000 miles per hour, while in restricted seas like the North Sea the rate is scarcely a twentieth of that amount. As the tidal wave differs from a wind-wave in not being a mere undulation, but a wave of translation, its *height* in any sea will depend mainly on the configuration of the shore, the form of bottom, and the direction in which it is propelled. In the open expanse of the Southern Ocean, the tidal wave rarely exceeds five or six feet, and in the Indian and Atlantic Oceans, perhaps eight or ten; but in bays and gulfs opening broadly to its course, and narrowing towards their interior recesses (such as the Bay of Bengal, our own Bristol Channel, and the American Bay of Fundy), it may rise to 20, 30, or, under favourable circumstances, even to 50 or 60 feet in height. And where such seas terminate in river-estuaries, the wave, still converging, forms a high head or wall of water, termed a *Bore*, which ascends the river with sudden and destructive impetuosity. Such are the tidal bores of the Tientang, which are said to extend across the river 30 feet high, and ascend with a velocity of 25 miles per hour, of the Hooghly 20 and 25 feet high, of the Garonne 10 and 12 feet, the Severn 9 feet, the Amazon 12 and 13 feet, and other rivers whose gradually narrowing estuaries are exposed to the concentrated incidence

of the tidal wave. On the other hand, inland seas and gulfs, whose openings are narrow, and lie transversely to the course of the tidal wave (as the Mediterranean and Baltic), experience little or no rise, and are next to tideless. Not only is the primary wave that sets up the Atlantic excluded from such seas, but their own areas are too limited to admit of the formation of any perceptible tide-wave of their own, and thus, for all practical purposes, strictly inland seas and lakes may be considered as tideless.

97. Unlike an ordinary wind-wave whose undulations are imperceptible at the depth of 1000 or 1200 feet, and unable to disturb from its repose the smallest grain of sand, the progressive motion of the tide-wave affects the whole mass of waters from the surface down to their greatest depths. The mechanical effects of the wind-waves are felt chiefly along the shore-line in their battering, wearing, and abrading action; the tide-waves, on the other hand, act more like currents, which, in their ebb and flow, are ever transferring, arranging, and re-assorting the sediments of the ocean. The wind-wave may be stilled, and the surface of the ocean be as smooth as a mirror; while it is driven from one coast it impinges on another; or the wave from one direction may counteract that from another: but the ebb and flow of the tidal wave is regular and incessant, and in its direction, time, height, velocity, and power, will continue the same while sea and land remain undisturbed in their present relations.

98. We now proceed to what are termed the CURRENTS of the ocean—movements which, like great rivers, are ever transferring the waters from one region to another. They form, in fact, the circulatory medium by which the ocean is maintained in a state of equilibrium, and depend primarily upon the unequal temperatures and densities of different zones of the ocean and the unequal evaporation sustained by these zones, and secondarily upon the rotation of the earth, which modifies the directions imparted by these primary causes. Wherever we have waters of different temperatures, or, what is the same thing, of different densities, there the lighter will ascend and the heavier descend; and wherever a deficiency takes place through evaporation, there the waters will flow in from the adjacent parts to make up the deficiency. But difference of density may also arise from different degrees of saltness, and wherever the salter water subsides and flows off as an under-current to some fresher region, there at the same time will the fresher and lighter flow in from above to restore the equilibrium. Understanding these facts, and bearing in mind that the continuity of the ocean is interrupted by continents and islands, reefs and shoals, and further disturbed by winds and tides, it will readily be seen why its currents should assume different characters and courses. In fact, the currents and counter-currents of the ocean are extremely complicated, and though the courses and causes of some of the main streams

are intelligible enough, there remains very much to be done in this department of Hydrography.

99. It is usual to arrange the ascertained currents into *constant, periodical, and variable*—the constant being those arising from the combined influences of unequal temperatures and densities in the waters of the ocean, the rotation of the earth, and the trade-winds; the periodical by the tides, the monsoons, and the sea and land breezes in tropical countries; and the variable, such as may be produced by local peculiarities in the tides and winds, the melting of ice in polar regions, and other similar causes. It is also customary to speak of *drift-currents* and *deep-sea currents*—the former due to the long-continued agency of the wind, and only affecting the waters to a trifling depth, the latter arising from the great primary causes of temperature and density above alluded to, and extending their influence hundreds of fathoms beneath the surface. In like manner, it may be useful to note the distinction between a *marine*, or upper current, and a *sub-marine*, or under-current; between a *current* flowing one way and a *counter-current* coming from an opposite direction; and between the mode of naming winds and water-currents—the former being named after the direction *from* which they blow, as a “west wind,” that is, one blowing from the west, and the latter after the direction *to* which they are flowing, as an “easterly current,” that is, one flowing towards the east.

100. The constant and deep-sea currents being the more important, it is to these and these alone that we would here direct attention. As already mentioned, the heat of the torrid zone diminishes the specific gravity of the equatorial waters, and occasions a greater evaporation there than in any other region, and as a consequence, the waters of the polar regions, being heavier, set in as an under-current towards the equator to restore the equilibrium. Here then we have two great primary currents setting in from north and south respectively; but as they proceed towards the equator they come stage by stage into latitudes where the earth's circumference rotates with greater velocity, and as they cannot at once partake of this increased momentum, they fall behind, as it were, and gradually assume a westerly course, in which their velocity is augmented by the influence of the trade-winds. They become, in fact, a combined equatorial current, where, growing warmer and warmer, they ascend to the surface, and are partly evaporated and partly flow over in warmer and lighter surface currents northwards and southwards to either pole, again to become colder and again to find their way to the equator in incessant circulation. From these four primary flows—the two from the poles towards the equator and the two from the equator towards the poles—arises the great circulatory system of the ocean, which is modified and broken up into a number of minor currents by configuration of coast, form of bottom, unequal reception of heat by different areas, the influx of rivers, and other

kindred causes. These various currents, having different directions, volumes, velocities, and temperatures, will be better understood, perhaps, by being arranged under the three great oceans—Atlantic, Indian, and Pacific—in which they respectively occur.

101. The principal and better known currents of the Atlantic are the Equatorial, the Guiana, the Brazil, the Gulf Stream, the Guinea, and the Arctic. Besides these, there are some minor drifts and branches, the courses of which will be better understood by reference to the map (p. 98) than by any amount of description. The *Equatorial*, as the name implies, manifests itself chiefly in the region of the equator, and flows across the ocean from the African towards the American continent. When more than half-way across it shows a tendency to bifurcate into a north-west branch and a south-west branch, and this tendency increases till within 300 or 400 miles of Cape St Roque, when it fairly divides—sending one main stream northwards by the coasts of Guiana into the Caribbean Sea; and another, somewhat feebler, southwards along the shores of Brazil. The length of the Equatorial current, from the coasts of Africa to the Caribbean Sea, is about 4000 miles; its breadth at its commencement 160, and where it divides 450 miles; its velocity is from 20 to 60 miles a-day; and its average temperature about 75° Fahr., or from 4° to 6° under that of the ocean through which it flows. Its southern, or *Brazil* branch, flows at a distance of about 250 miles from the coast (the intermediate space being occupied by variable currents), and at the rate of 25 miles per day; a rate, however, that is sensibly diminished by the cross stream from the Plata, and which gradually declines till the current ultimately dies away in the Strait of Magellan. The north-west branch of the Equatorial, spreading out as it proceeds and gradually diminishing in speed, ultimately merges into the drift of the north-east trade-winds; while the *Guiana* section proceeds unimpeded to make the circuit of the Caribbean Sea and the Gulf of Mexico.

102. As they make the circuit of the Caribbean Sea and the Gulf of Mexico, these western branches of the Equatorial current acquire more heat, a greater degree of saltness, and that intense blue colour so characteristic of briny waters. With these new acquisitions they leave the Mexican Gulf, and, pressing through the narrow channel of Florida, become the celebrated *Gulf Stream*—of all the Atlantic currents the most wonderful in its character and the most important in its results. Doubling the Cape of Florida, this Gulf Stream (that is, stream from the Gulf of Mexico) flows north-east in a line almost parallel to the American coast; touches the southern borders of the banks of Newfoundland; and thence, with increasing width and diffusion, proceeds across the Atlantic, till, in the region of the Azores, it spreads out into two great branches—one curving southwards towards the equator, and the other flowing northwards, imping-



ing in its course against the western coasts of Europe, and ultimately losing itself in the waters of the Arctic Ocean. The length of this great ocean-river from its commencement to the Azores is 3000 miles, and its greatest breadth about 120 miles. When it leaves the Strait of Florida its velocity is about 4 miles an hour; off Cape Hatteras, in North Carolina, it is reduced to 3 miles; on the Newfoundland banks it is further reduced to $1\frac{1}{2}$ miles; and this gradual abatement of force continues with its diffusion across the Atlantic. A similar decrease takes place in its temperature, the maximum of which in the Strait of Florida is 86° , or 9° above that of the ocean in the same latitude. Off Newfoundland, in winter, it is said to be from 25° to 30° above the water through which it flows; in mid-ocean, from 8° to 10° ; nor is the heat wholly lost when it impinges against the shores of Western Europe. The Gulf Stream is thus, in reality, a great thermal ocean-river, incessantly flowing from warmer to colder regions, diffusing warmth and moisture along its course, and tempering the climates of countries that lie within its influence. Compressed, as it were, at its commencement between two areas of colder water, its deep-blue warm current rises in convexity above the surrounding ocean; but as it proceeds it cools, becomes diffused, assumes the ordinary level, and partakes of the greener hue of northern waters. It sets a limit to the southward flow and chilling influences of the Arctic iceberg, that melts away in its warm stream, and becomes at the same time the great natural barrier between the Life of the Northern and Southern Atlantic.

103. The Equatorial current, flowing westward from Africa to the Gulf of Mexico, and the Gulf Stream, flowing from that gulf eastward to the Azores, and thence curving southwards, produce a great whirl, as it were, in the Atlantic, in the centre of which there is the still water of the *Sargasso*, or *Grassy Sea*. This area, so called from the vast accumulation of the *Sargassum* (*Fuens*) *natans*, and other floating sea-weeds, swarms with a Life, vegetable and animal, peculiarly its own, and presents one of the most remarkable features in the geography of the Atlantic.—Following the southward flow from the Azores, and the south-easterly flow from the south coast of Ireland, against which a minor current (*Kennel's*) from the Bay of Biscay impinges and recoils, there arises a new and increased stream, which holds onwards to the African continent. Of this current, the eastern portion sets in through the Strait of Gibraltar to the Mediterranean, while the western trends southward, and becomes the *North-African and Guinea Current*. This stream, from 160 to 180 miles broad, hugs closely the coast, and for a long part of its course flows in contact with, but counter to, the equatorial current, with whose waters it is supposed ultimately to mingle in the South Atlantic.

104. The next and last great current of the Atlantic to which

our limits will permit us to advert, is the *Arctic*, or main cold-water stream from the north. Setting strongly down the eastern coast of Greenland, it partly doubles Cape Farewell, where, augmented by the Davis Strait Current, it holds southward to Newfoundland. Arriving at Newfoundland, it sends a branch through the Strait of Belle Isle to the St Lawrence, while the main portion continues its course till it meets the Gulf Stream. Here it divides, one portion flowing southward to the Caribbean Sea, which it enters as an under current; the other, flowing south-west, forms the United States *Counter-current*. "The Arctic Current thus replaces the warm water sent through the Gulf Stream, and modifies the climate of Central America and the Gulf of Mexico, which, but for this beautiful and benign system of aqueous circulation, would be one of the hottest and most pestilential in the world."

105. In the more limited and land-locked area of the Indian Ocean, the constant currents are few, and the periodical numerous and important. The *Equatorial* is less defined than in the Atlantic, and consists of a westward tendency of the tropical waters towards the coasts of Africa, where, divided by the large island of Madagascar, one branch sets down with considerable force, and forms the *Mozambique Current*, while another, broader but feebler, trends southward, again to be united with that from the Mozambique Channel. The combined stream, now from 90 to 100 miles broad, with a temperature 7° or 8° above that of the ocean, and with a velocity of 60 or 80 miles a-day, sets in towards the Cape of Good Hope, and constitutes the *Agulhas* or *Cape Current*. Of this current, one main branch, doubling the Cape, flows northward into the Atlantic as far as St Helena; where, meeting the Guinea Current, it is deflected westward, and merges into the Equatorial of that ocean; whereas another portion, obstructed by the Agulhas bank, is turned back, and, combining with the connecting flow from the Southern Ocean, constitutes the important *Counter Current* of the Indian Ocean. We say "important" counter current, for it flows along the direct route to Australia, and with a velocity (after leaving the Agulhas bank) of nearly 50 miles a-day—a motion which, though gradually declining, is still sensibly felt at more than midway between the Cape and Tasmania. North of the equator, the currents and surface-drifts of the Indian Ocean are regulated by the monsoons, and vary, of course, with the seasons; while in the Red Sea, Arabian Sea, Persian Gulf, and Bay of Bengal, they set in and out with the local winds, and often change in a very complicated and capricious manner.

106. The currents of the Pacific are co-extensive with its greater area, but are less decided in their courses in consequence of the numerous obstructions presented by its reefs and islands. They are altogether less known than those of the Atlantic and Indian oceans, and their investigation is greatly complicated by the fre-

quency of counter-currents, surface-drifts, and local gyrations. Being all but excluded from the Arctic Ocean, it is to the great interchange between the cold waters of the Antarctic and the warmer waters of the equator that we must look for the primary impulse of its currents. Beginning with the Antarctic, to which it opens broadly, we have first the *Drift Current* of that ocean setting in towards the north-north-east, then north-east, and lastly towards the east-north-east till it nears the coasts of Chili, where it divides—sending one branch south along the coast to form the *Cape Horn Current*, and another northwards to form the *Peruvian Current*, so remarkable for its cold stream along a coast of torrid temperature. This Peruvian, Chili, or “Humboldt’s Current” (as it is sometimes called after its first investigator), has a temperature along more than 400 miles of its course 12° or 14° below that of the surrounding atmosphere, and 8° or 10° below that of the ocean through which it flows. As it holds on to the north it gradually becomes warmer and inclines to the west, till in 20 degrees south latitude it turns fairly to the westward and merges into the great *Equatorial Current* of the Pacific. This vast current, occupying the entire torrid zone, and aided in its westerly flow by the trade-winds and the tidal wave, sweeps boldly across from South America on the one hand to the Indian Archipelago on the other. As it proceeds, it sends off a few minor streams to the north and south, which become *counter-currents*, but the main mass holds onward to the Indian Islands, where it is broken up into several sections, and becomes further complicated by the *monsoon drifts* of that region. One main portion, however, sets southward with its warm waters along the coasts of Australia under the name of the *New South Wales Current*, and enters the Antarctic Ocean; while another, still more decided, trends northward of the Philippine Islands, and becomes the well-known *Japan Current*. This last current, like the Gulf Stream of the Atlantic, which it greatly resembles in its course and character, carries the warm waters of the equator to the Northern Pacific; but having but a partial opening into the Arctic Sea by Behring Strait, it sweeps round the Aleutian Isles and shores of Russian America, and returns again, with diminished temperature, to the torrid zone. As it proceeds along the shores of Oregon and California it begins to bifurcate, one branch trending westward into the ocean and another holding still southward and merging into the waters of the *Mexican Current*. This current, which flows along the coasts of Mexico and Central America, is an alternating rather than a constant one—depending on the monsoons of these coasts, and setting south-eastward during winter and north-eastward during the opposite half of the year. Besides the preceding, other minor currents have been noticed in the Pacific, as the *Carolinian Monsoon Current*—an alternating flow depending on the influence of the Indian and Chinese monsoons; the *Okhotsk Current*

—an easterly set of the waters of that sea, arising from the south-east and easterly winds that prevail there during the summer ; the *North Equatorial Counter-Current* ; and *Mentor's* (after the Prussian surveying vessel of that name) or the *South Equatorial Counter-Current*. These, however, and others of less note, are but partially known ; the numerous islands and reefs of the Pacific not only interrupting the regular flow of the major currents, but rendering the minor ones more complicated and less decided in their directions.

107. Such are the principal *constant* currents of the ocean—the *periodical* and *variable* being too multifarious and local for the limits of a general outline. In whatever character they appear, these currents are all-essential to the equilibrium and uniformity of the ocean. Excessive evaporation in the torrid zone is instantly counterbalanced by an influx from the temperate and frigid. The heated waters of the equator are modified by the colder currents from the poles ; while the warmer and lighter waters of the equator flow over to temper the rigours of the polar seas. The salter and denser water of one area subsides and flows off as an under-current to some fresher region ; while the fresher and lighter flows in from above to supply the deficiency. The heavier ever descends and the lighter ascends ; and thus, vertically as well as horizontally, an incessant circulatory and equalising system is established throughout the expanses of the ocean. The colder ever flowing to the warmer and the warmer to the colder, their influence also extends to that of the lands against which they impinge, and thus additional warmth and moisture is borne to one country and refreshing coolness to another. Climate, and consequently terrestrial life, vegetable and animal, are modified by these currents ; and in all likelihood they play equally important parts in the distribution and arrangement of the life of the ocean. Some coursing in shallow streams, and others pressing forward as deep and impetuous rivers, they must exercise considerable influence in transporting and assorting the sediments and debris of the ocean—here laying down long belts of uniform sediment, and there ploughing out deeper troughs and valleys—here arresting the progress of icebergs, that drop their burden of boulders and gravel on one zone, and there fostering the growth of animalcules and zoophytes (foraminifera, corals, &c.), that drop their exuviae and rear their structures in another zone equally definite and restricted. Besides these great natural functions they often subserve the purposes of navigation and commerce ; and though the steam-ship has rendered the mariner less dependent on winds and tides and currents, he still skilfully seeks to avail himself of the favouring stream of one current and to avoid the opposing influence of another.

RECAPITULATION.

In the preceding chapter attention has been directed to the main natural divisions of the Ocean ; to the composition, density, depth, and temperature of their waters ; and to the waves, tides, and currents by which they are respectively traversed. The divisions of the ocean into Pacific, Atlantic, Indian, Arctic, and Antarctic areas, are sufficient for the ordinary purposes of description, and the nomenclature of their various ramifications after some discoverer, position, adjacent country, colour, or other peculiarity, has been long established in general geography. Their further subdivision into *zones*, *belts*, *regions*, and the like, comes under the consideration of Climatology rather than of Hydrography ; and the technicalities (bays, gulfs, straits, creeks, &c.), by which their minor sections are distinguished, have been already noticed in par. 37. Their dimensions are matters merely of measurement and calculation ; their composition, depth, temperature, and the like, are subjects for experiment and observation. Occupying different positions on the earth's surface, and having different configurations, the oceans and seas, properly so called, are necessarily characterised by different physical and vital conditions. The Pacific, almost shut out from the influences of the Arctic Ocean, is in a different condition from the Atlantic, which communicates freely with the north ; while the Indian Ocean, encircled on three sides by land, and lying largely in the torrid zone, presents external conditions differing widely from either. All three open broadly to the great Southern Ocean from which they receive the primary impulse of their tides and currents ; but while the Indian and Atlantic are deep and free, the Pacific is largely obstructed by reefs and islands. It should also be borne in mind that the main oceans, like the great continents, lie meridionally or at right angles to the equator, and are thus prevented, by the intervention of the land, from receiving in their winds, tides, and currents, the normal effect of the earth's daily rotation. Though differing slightly in different areas, the composition of sea-water is, on the whole, very equable, and $3\frac{1}{2}$ per cent of saline ingredients may be set down as the general average. Along the courses of the trade-winds, in circumscribed seas, and other similar areas subjected to active evaporation, this percentage may be slightly increased ; while in others it may be correspondingly decreased by periodical

rain-falls, by the melting of polar ice, or by the influx of large and rapid rivers. The average density of sea-water, as compared with pure water at 62° Fahr., is 1.0275, and this density, corresponding as it does with an average saltness of $3\frac{1}{2}$ per cent, will decrease where the water is fresher and increase where it becomes saltier than the general average. This saltness and density renders the ocean less vaporisable than fresh water, and also keeps it longer from freezing—the freezing point of fresh water being 32° Fahr., while that of sea-water is 28 $\frac{1}{2}$ °. This composition is all-essential to the wants of its plants and animals, and though the rivers are incessantly carrying in fresh accessions of saline matter, the equilibrium is ever maintained by these wants as well as by the chemical interchanges that take place among its sediments. The average depth of the ocean is estimated at 4 miles, and reliable soundings have been taken at nearly 5 miles; but, as a large portion is much under the general average, it is highly probable that some of its recesses may sink to the depth of 8 or 10 miles. So far as we know, the ocean bed has its deeps and shoals, its pits and precipices, its troughs and ridges, very much like the surface of the dry land. In fact, it is but the submerged surface of former lands; and no better conception can be formed of its irregularities than by standing on some lofty mountain, and supposing the hills and valleys, the glens and gorges, the plains and plateaux that lie beneath, to be covered with water. The surface-temperature of the ocean varies of course with the latitude, being highest at the equator, and gradually decreasing towards either pole. In the torrid zone temperatures varying from 78° to 88° have been noted, and from these maxima it declines, stage by stage, to the perpetual ice of the polar regions. The mean temperature has been ascertained to be 39 $\frac{1}{2}$ ° Fahr., or nearly that of water at its maximum density; and between the 56th and 57th parallels (S. lat.) this temperature has been found to be uniform from the surface to the greatest depths. Towards the poles, however, the surface becomes colder, and the mean of 39 $\frac{1}{2}$ ° in the 70th parallel is not found till at the depth of 750 fathoms; while towards the equator the surface grows warmer, and the mean of 39 $\frac{1}{2}$ ° is not reached till at the depth of 1200 fathoms. The motions to which the waters of the ocean are generally subjected are waves, tides, and currents. Waves are produced by the friction and impact of the winds, and occasionally by earthquake commotions. Wind-waves vary from a mere ripple of the surface to billows 10, 20, 30, and 40 feet in height; earthquake-waves have been known to rise higher, and,

rolling in with greater impetus, they are more destructive. Tides are produced by the attraction of the moon and sun acting on the mobile waters of the ocean. In consequence of the earth's rotation there are two tides a-day ; and owing to the moon's revolution, and her position as regards the sun, there are two spring-tides and two neap-tides a-month. The moon being the nearer luminary, her attraction or tide-wave compared with that of the sun is as 100 to 38 ; and from this cause also the difference between spring-tide and neap-tide in any locality is as 7 to 3. The currents of the ocean depend chiefly upon the unequal temperatures and densities of its different regions—the colder and warmer, the denser and lighter, the salter and fresher, incessantly tending to interchange and equilibrium. These currents are either constant, periodical, or variable—the constant depending on the primary and established relations of the earth ; the periodical on monsoons, tides, and sea and land breezes which occur in certain seas at stated seasons ; and the variable on local winds, melting of ice, river floods, and other similar contingencies.

IX.

THE WATER—ITS RIVERS AND LAKES.

108. HAVING directed attention to the Water, as manifested in the larger expanses of oceans and seas, we now turn to its minor exhibitions in springs and streams, rivers and river-systems, lakes and lacustrine areas. In the former instance the waters were truly oceanic, and characterised by their *saltiness*; in the present instance they are more terrestrial, if we may so speak, and characterised, for the most part, by their *freshness*. They belong, however, to the same great aquatic system. The vapour elaborated from the ocean ascends invisibly and diffuses itself through the atmosphere, where, subjected to colder currents, it is condensed, and becomes visible in clouds, mists, and fogs. Receiving further condensation through greater cold or electrical agency, it returns again to the earth in the form of dew, rain, hail, sleet, and snow. When in the atmosphere water belongs to the domain of Meteorology; when it returns to the earth it comes once more within the province of the Hydrographer, who tracks it in spring, stream, and river, to the ocean. The circulation of water is incessant: now in the ocean, now in the atmosphere, now in the tissues of plants and animals, now in the crust of the earth, now coursing its surface, and anon in the ocean—again to repeat the same circuit, and this without intermission while the present relations of the universe endure.

Springs and Streams.

109. The water that falls on the surface of the land is partly and mainly carried off in runnels, partly supplies the wants of plants and animals, and partly sinks into the soil and rocky strata. Percolating the rocks, it collects in chinks, fissures, and other cavities, or, meeting in with some impervious bed, it accumulates in the porous strata above, and there forms extensive sheets of subterranean water. In virtue of the great law of gravity, water is ever tending to lower levels, and thus what gathers in the higher portions of the crust finds its way, in process of time, through rents and crevices, and springing forth

holds on its course to the valleys below. Where there are no impervious beds in the higher lands to intercept its downward tendency, it often percolates to vast depths, and far below the surface of the adjacent valleys; but sooner or later it is arrested by some obstructing stratum, and there, in obedience to hydrostatic pressure, it bursts forth through the nearest outlet, and rises to the surface with a force proportionate to the height and volume of the accumulated waters. In whatever manner water may accumulate within the crust, or find its way again to the surface, the outlets are known as *springs*; and these springs occur in all countries and at all levels—some issuing with force even from the bed of the ocean. It is usual to speak of them as surface springs and deep springs, as perennial, intermittent, hot, cold, mineral, and the like—characteristics, the consideration of which belongs more, perhaps, to Geology than to Physical Geography.

110. It may be noticed, however, that *surface springs* are those which issue from superficial beds of sand, gravel, and the like, and being immediately dependent for their supplies on the amount of rainfall, are often very feeble or altogether dry in summer, though gushing forth copiously during winter. *Deep-seated springs*, on the other hand, uninfluenced by summer droughts or winter rains, flow steadily at all times—though, generally speaking, the great majority of springs are more or less affected by the seasons. *Perennial springs*, as the name implies, are those that flow year after year without signs of abatement. They are evidently deep-seated, and many that were celebrated by the ancients gush copiously now as they did thousands of years ago. *Intermittent*, on the other hand, are those that well forth at one period, and at another stop suddenly, and seemingly in a capricious manner. They are no doubt connected with subterranean reservoirs, whose waters alternately sink beneath and rise above their outlets; and this may be occasioned by sudden droughts and rainfalls, by the expansion and escape of steam and gases, by the ebb and flow of tides, and other kindred causes. Surface springs generally vary in temperature with the seasons: deep-seated ones, being beyond these influences, are equable at all times. It was stated in par. 16 that the earth's crust, at the depth of 80 or 90 feet, is unaffected by summer's heat or winter's cold; but that below this depth the temperature went on increasing at the rate of 1° Fahr. for every 50 or 60 feet of descent. Springs whose sources are above this invariable stratum will therefore fluctuate in temperature according to the season of the year; those that ascend from beneath it will gradually increase in heat—the deeper being the hotter. *Cold springs* generally flow from chilled mountain sources; *hot springs*, on the other hand, arise either from vast depths, or are situated in the neighbourhood of volcanic action. Thermal and hot springs (hot even beyond the boiling point of water, and accompanied by violent jets of steam) occur abundantly, and on a gigantic scale,

in Iceland, the Azores, Central Asia, New Zealand, the Andes, and other volcanic regions; they occur also in Britain, the Pyrenees, Germany, Austria, and other countries far removed from any centre of volcanic activity. Like ordinary springs, hot springs are either perennial or intermittent — perennial, like those of southern France and the Pyrenees, that have flowed without abatement since the time of the Romans; and intermittent, like the Geysers ("roarers") of Iceland and New Zealand.

111. The waters that percolate the earth being of different temperatures, and often containing carbonic acid and other gases, act chemically, less or more, on the rocks through which they pass. In this way they become *mineral springs*—that is, hold in solution various mineral ingredients, which are either deposited along their courses on land, or borne onward to the ocean. Some, for example, are saline, or contain salt; some chalybeate, or contain iron (*chalybs*, iron); some siliceous, or contain flint (*silex*); some calcareous, or contain lime (*calx*); while others give off sulphurous vapours, or are impregnated with the salts of various minerals and metals. Such are the sulphuretted waters of Harrogate, the brine springs of Cheshire, the borax springs of Tuscany, the travertine or lime-depositing waters of the Anio, the siliceous hot springs of Iceland and the Azores, and thousands of others that occur in almost every country. To the same category also belong escapes of steam, hot mud, and the like, so frequent in volcanic districts; and in the same way may be ranked discharges of gases, naphtha, petroleum, and so forth, which generally come to the surface in connection with water, and are seemingly dependent on its percolating and hydrostatic power.

112. Insignificant as springs may individually appear, they perform in the aggregate most important functions in the economy of nature. If the waters, as they fall from the atmosphere, were immediately carried off by surface runnels, the earth would at one time be flooded, and at another parched by destructive droughts. Sinking, however, into the soil, and diffused through its mass by capillary attraction, percolating the crust, and accumulating in its fissures and porous strata, this water is stored up, as it were, for future use, and given out by springs in moderate and continuous supplies. These springs are the great vivifiers of external nature, supplying one of the main wants of animal life, and clothing even the desert around them with verdure and blossom. They are also the fountainheads of our streams and rivers, whose channels, but for this incessant discharge from the earth, would often be reduced to a mere succession of stagnant pools, instead of flowing, as they now do, with never-failing currents. Ever silently seeking their way down through the rocky strata, these springs dissolve some minute portion, and carry it to the ocean, there to compensate for the abstraction of mineral ingredients by the plants and animals that inhabit its waters. Acting chemically, this internal permeation of water is evidently

connected with the formation of mineral crystals and metalliferous veins, and also with that greater metamorphism of rock-masses by which strata of one nature are in process of time converted into strata of another and very different nature.

113. *Streams*, as explained in par. 38, are formed by the union of several springs, and by the union of one stream with another, till, increasing in breadth and volume, they assume the character of rivers. When mainly dependent on springs (though augmented by rainfalls), they are said to be *permanent*; but when arising solely from rainfalls, melting of snow, and the like, they are *temporary*, and often of short duration. Permanent streams are usually distinguished, according to their dimensions, as *rills*, *streamlets*, *brooks*, *rivulets*, &c. (each country having its own expressive term for such distinctions), and their currents form *ravines*, *glens*, *gorges*, *valleys*, and other descriptive excavations. Temporary streams, on the other hand, are little noticed save in regions subjected to periodical rains, their dry courses being known as *nullahs* in India, *wadies* in Arabia, *creeks* in Australia, and so on in other localities. Streams occur in all countries, but fewest and least, of course, in rainless latitudes. Their great headquarters are the slopes of mountain-chains and the outer margins of valleys, and there they impress on the country geographical features and geological relations peculiarly their own. Coursing down the mountain-sides with headlong speed, they form ravines, gorges, and waterfalls—their excavating power depending primarily upon their velocity; and, secondarily, on the nature of the materials to be eroded. The aggregate amount of debris brought down by streams and torrents is enormous; and one has only to cast his eye over some lofty mountain-slope to see how deeply it is scarred and seamed and furrowed by these restless and resistless agencies. In fact, the great diversity of mountain scenery depends on this erosive power of water—all the hills and highlands depending less for their present outlines on the igneous forces by which they were upheaved, than on the meteoric and aqueous agencies to which they have been subsequently subjected.

114. Streams, like springs, subserve important purposes in the economy of nature. They constitute a great network of drainage for the land—carrying off superfluous moisture, and presenting its accumulations in bogs, swamps, and morasses. They are at once the carriers and distributors of water—dispensing in measured supplies the heavy rainfalls, melting snow and ice of the highlands, to the less rainy regions of the plains below. Geologically, they are incessant workers of change—here cutting out ravines, glens, and gorges—there filling up lakes with the eroded material; here wearing deeper channels for the further drainage of swampy lowlands, and there assisting in that universal transport of sediments to the ocean, which is destined to become the strata of future lands. Man, too, ever early locates himself on their

banks, employing their perennial supplies in the wants of his home, in the irrigation of his fields, or as a cheap and effective agent in the turning of his machinery.

Rivers and River-Systems.

115. As streams are formed by the union of springs and minor runnels, so rivers are formed by the union of streams. In general they have their origin in some notable spring, lake, morass, or melting glacier, far up among the mountains. This infant current, augmented by the accession of other springs, becomes a main stream; and this stream, by the influx of other *affluent* or *tributary* streams, very soon assumes the dimensions of a *RIVER*. A river is thus said to have its *head* or *source* in some higher region; to flow along a certain *course*; to wear out for itself a *bed* or *channel*, whose margins form its *banks*; and ultimately to discharge its waters in some lake, sea, or estuary by a *mouth* or *mouths*. In other words, every river has a *rise*, *flow*, and *fall*, which form the essential features of its existence. Where the ground, in the ultimate part of a river-course, is low and level, the accumulating waters often discharge themselves by several branches, and form a *delta* (par. 68), as in the Nile, Niger, Indus, Ganges, &c.; and such deltas may be either marine, lacustrine, or fluvial, according as the rivers flow into the sea, a lake, or another larger river. Where the tides and river-currents contend, there frequently accumulates across the entrance a spit of sand or gravel, known as a *bar*; and these bars are serious impediments to the navigation of some of the finest rivers. As all rivers flow from higher to lower levels, the terms *up* and *down* have reference to this natural condition; and as a spectator descends with the current, the *right bank* lies on his right hand, and the *left bank* on his left. When he ascends, however, this order is reversed—the right bank being then on his left hand, and the left bank on his right.

116. The whole extent of country drained by a river and its tributaries is spoken of as the *basin* of that river (basin of the Thames, basin of the Tay, &c.); and this naturally, from the hollow or valley-shaped aspect which such districts usually assume. As streams unite to form rivers, so several river-basins may descend towards the larger depression of some inland sea; and thus geographers speak of the “basin of the Baltic,” the “basin of the Mediterranean,” and, carrying the idea still farther, even of the “basin of the Atlantic.” As the great oceans form the ultimate receptacles for all the rivers (with one or two exceptions), the river-basins that descend to any ocean constitute what is termed the *river-system* of that ocean; and thus we have the Arctic System, the Atlantic System, the Pacific System, and the System of the Indian Ocean—together with the Aralo-Caspian

System and the minor basins of Utah, the Mexican table-land, and the plateau of Bolivia, whose streams are strictly continental, and confined to these areas. The line or ridge that separates one basin or system from another basin or system is termed the *watershed* of these basins (or *watershed*, if we affect a German derivation)—all the springs and streamlets “shedding or parting off,” as if from the ridge of a house, to their respective areas. Though hills and mountains generally form the boundaries between river-basins, it does not follow that watersheds should always consist of elevated ground—a few feet of fall being sufficient to determine the current in either direction. The low ridge that thus separates the streams of different basins is spoken of as a *portage* (Fr.), from the fact that goods and boats are frequently carried across from one stream to another; and thus, it may be, completing the passage of water from one side of a continent to another.

117. Such are the usual *terms* employed by geographers in treating of rivers; the *characteristics* of the rivers themselves—that is, their rise, course, length, volume, depth, velocity, capability of navigation, and the like—require more detailed consideration. As to their *rise*, it has been already stated that it generally takes place in some upland spring, lake, or morass, and not unfrequently in the melting terminus of some descending glacier. Many of these springs, especially those issuing from chasms and caverns in limestone districts, are often of considerable magnitude, being, in fact, full-flowing streams when they make their appearance at the surface. Mountain lakes, being the recipients of springs and other runnels, become in like manner copious sources—the streams that flow from them being already rivers in miniature. The same may be said of morasses and high swampy tracts, which, being saturated with springs and rainfall, discharge at once regular and continuous streams. Glaciers and melting snows, though less frequent as sources, are important regulators of supply in the regions where they occur—their waters flowing freest when the summer's heat has most diminished the lowland streams. Whatever their origin—and it must be confessed that the exact sources of many of our larger rivers are unexplored and undetermined—their *courses* are primarily directed by the general slopes of the continents. In the Old World, the great slopes being north and south, the principal rivers flow in these directions, just as in the New World their courses are easterly and westerly, in obedience to the relief of that continent. But while this is true, and necessarily so, as regards the main slopes, it should be borne in mind that there are many causes tending to modify the courses and directions of rivers. Among these, for example, may be noted the strike of the secondary hills and spurs, which are usually at right angles to the main chains, the occurrence of lines of fracture in the crust, the passage from one formation to another whose strata

lie in a different direction, and the alternation of softer and harder beds—the current readily excavating its channel in the line of the former, but obstructed and turned aside in another direction by the latter. All these, and many other circumstances, concur to give streams and rivers very irregular courses—winding and bending, turning and returning, but still guided in the main by the primary slope of the region. As to the *length* of their courses, that will depend primarily upon the extent of the region through which they flow—the larger area giving scope to the longest river; and, secondarily, upon the bendings and windings to which they may be subjected. The direct distance between the source and outlet of a river may not exceed two hundred miles, and yet its absolute course, through all its doublings and windings—its *development*, as it is termed—may amount to a thousand. This absolute length gives no true idea, however, of the importance of a river, either in nature or to navigation—its value mainly depending on the permanence of its volume, its depth, and velocity.

118. The *volume*, size, or magnitude of a river—that is, the quantity of water which its channel contains—depends on many collateral circumstances, but chiefly on the extent of country drained by its affluents. There may occasionally be some peculiarity of basin as regards rainfall, melting of snow, and the like; but, generally speaking, the greater the area over which the tributaries ramify, the greater the quantity of water brought to the main trunk or channel. In temperate latitudes this volume is somewhat lessened in summer and increased in winter; but on the whole the supply is pretty equable and permanent. Sudden excesses, arising from heavy rainfalls, rapid melting of snow in spring, and the like, are merely temporary, producing what are termed *floods*, *freshets*, and *debacles*. In tropical countries, on the other hand, where the rains fall and the snows melt only at stated seasons, these floods assume the character of regular and *periodical inundations*. And where the rain falls in the low countries at one period, and the far-off mountain-snows melt at another, there may be two such inundations occurring with wonderful regularity both in time and amount of overflow. Again, where the sources of a river lie near the equator and its outlet in the temperate zone (as in the case of the Nile), a considerable time must elapse between the equatorial rainfall and the deltaic inundation; and the length of this lapse will depend partly on the windings of the river, and partly on the slope and freedom of the channel. In the preceding instances, whether subjected to irregular floods or periodical inundations, the river itself is always more or less *permanent*; but in some regions, as Australia and South Africa, many of their so-called rivers are merely *temporary*, being roaring impetuous torrents during the rains, and a succession of stagnant pools and dry shingly reaches during the season of drought.

119. The *velocity* of a river depends mainly on the slope or declivity down which it flows, partly on the nature of its channel, according as this may be straight or winding, deep and narrow, or wide and obstructed by rocks and shoals, and partly also on the amount or volume of its current. As all rivers descend from higher to lower regions, and many of them from very elevated sources, they may be said to have an upper, middle, and lower course—the *upper* being characterised by rapidity of stream through gorge and glen and waterfall; the *middle* by less velocity through rapids and cataracts; and the *lower* by a quiet steady flow through level and alluvial plains. Other things being equal, the deeper the river, the more rapid the current—the greatest velocity being in the centre of the stream, and a little below the surface, where there is least retardation from friction on the sides and bottom of the channel. The velocity of rivers is an all-important consideration in their geographical, geological, and commercial relations. The more rapid their currents, the less the irrigating effect, and the greater the geological effect on the countries through which they flow; and in proportion as their velocity increases, so their fitness for the purposes of navigation is diminished. Formerly a descent of more than 1 foot in 200 was considered unnavigable; and though the power of steam has enabled man to contend with higher velocities, it is still the velocity of current more than the depth of water that renders a river unavailable as a means of internal communication. Geologically, the cutting as well as transporting power of rivers is greatly aided by the rapidity of their currents; hence the effect of mountain torrents compared with the quiet, sluggish flow of the lowland river. It has been calculated, for example, that a velocity of 3 inches per second will tear up fine clay, that 6 inches will lift fine sand, 8 inches sand as coarse as linseed, and 12 inches fine gravel; while it requires a velocity of 24 inches per second to roll along rounded pebbles an inch in diameter, and 36 inches per second to sweep angular stones of the size of a hen's egg. During periodical rains and land-floods the currents of rivers often greatly exceed this velocity; hence the tearing up of old deposits of gravel, the sweeping away of bridges, and the transport of blocks many tons in weight—an operation greatly facilitated by the circumstance that stones of ordinary specific gravity (from 2.5 to 2.8) lose more than a third of their weight when immersed in water.

120. The *depth* of rivers is as various as the circumstances under which they occur, though, generally speaking, the greater the volume of water, the deeper the channel of excavation. In their upper and middle courses, the stream, being for the most part through steep rocky excavations, is of no great depth; but in the lower and slower portion the waters accumulate and deepen as they proceed. The *outlet*, or *embouchure* (Fr.), as it is frequently termed, depends altogether upon the conformation of the country.

Some rivers discharge themselves at once and by a single mouth into the ocean; others into an estuary, and are consequently affected for a certain length by the flux and influx of the tides; while others, again, creep sluggishly along, branching and bifurcating through their swampy deltas, till they find their way into the sea by several main mouths. Seeing that the conditions of rivers are so exceedingly varied, their *facilities for navigation* will depend in each case on the nature of the outlet as accessible from sea and free from bars and banks, on depth and volume of water, and on the velocity of the current. Length of course and volume of current may be obstructed by rocks and shoals; depth and volume may be rendered unavailable by shallow bars and shifting sandbanks.

121. It will be seen from what has been stated respecting the characteristics of rivers, that each has its own individuality, and that this individuality depends in the main on the geographical position, extent, and superficial configuration of its basin. To give an account of these basins would be to describe in detail the rivers that traverse the surface of the globe—a subject which extends beyond our limits, and belongs more especially to hydrography and general geography. All that our outline will permit is the arrangement of the basins into *SYSTEMS*, and a notice of some of the more remarkable rivers that belong to each system. As all the affluent streams of a river belong to one basin, so all the river-basins that descend to the same ocean-basin—whether directly, as the Niger to the Atlantic, or intermediately through some inland sea, as the Nile to the Atlantic through the Mediterranean—belong to the same river-system. In this way we have four great *oceanic river-systems*—the Arctic, Atlantic, Pacific, and Indian; and one or two minor *continental ones*, as the Aralo-Caspian, the Utah, the Mexican, and the Bolivian. To the main characteristics of these respective systems, and the leading rivers that compose them, the attention of the student is now directed.

122. The *Arctic System*, as the name implies, embraces those rivers which, obeying the northern slope of the Old and New World continents, discharge themselves into the basin of the Arctic Sea. Lying for the most part within the limit of constantly-frozen ground (see Atlas, Plate XII.), and flowing through notably level plains, there is a degree of sameness in the conditions of these waters which renders the Arctic more homogeneous in its character than any other river-system. For convenience, however, it may be divided into three sections—Asiatic, European, and American—according as the rivers belong to these respective areas. 1. The Asiatic comprehends the great Siberian rivers—the Kolyma, Indigirka, Lena, Olenek, Yenesei, and Obi, which, rising in the Altai, Yablonoi, and Stanovoi mountains, receive numerous tributaries in the upper and middle stages of their course, wind slowly through the plains, and ultimately dis-

charge themselves by wide estuarial mouths into the Arctic Sea. The largest of these are the Lena, Yenesei, and Obi, which, had they discharged themselves into temperate or tropical seas, would have taken rank with the noblest rivers on the globe. The Lena has a development or winding course of 2400 miles, receives several important tributaries (Vitim, Olekma, and Aldan), and is computed to drain an area of 594,000 square miles. It is a deep sluggish stream in the lower part of its course, and finds its way into the sea between high banks of frozen mud and sand, in which are imbedded the remains of mammoth, rhinoceros, and other huge mammals, which in former ages (upper tertiary), and under different geographical conditions, inhabited the plains of Northern Asia. The Yenesei is still larger, having a development of 2800 miles, and a basin of not less than 784,000 square miles. It receives several large affluents—the Great and Little Kem, the Angara from Lake Baikal, and the Upper and Lower Tongouska, most of which, in their upper and middle stages, are impetuous torrents, interrupted by rapids and cataracts. The Obi, or Irtish-Obi, as it may be termed, from the equal importance of its two main branches, is the most notable of these rivers, having a development by either branch of 2400 miles, and a conjoint basin embracing nearly one-third of the entire area of Siberia. As already mentioned, the main affluent of the Obi is the Irtish, and the Irtish receives in turn the streams of the Tobol and Ishim. One leading characteristic of all the Siberian rivers is, that though rapid and sufficiently diversified in the upper stages of their courses, they become sluggish and monotonous on entering the low boggy plains. Frozen, like the ocean into which they enter, for many months of the year, they are unfit for navigation; while, on the melting of the snows in their upper and more southerly sources, the swollen floods, finding an insufficient outlet by the level, ice-locked, and more northerly mouths, overspread the country in lakes and morasses, which render its surface still more dreary and inhospitable. 2. The European section of the Arctic river-system embraces the Petchora, which falls directly into the ocean, and the Mezen, Dwina, and Onega, which enter it indirectly through the White Sea. With the exception of the Dwina, on which Archangel is situated, and which drains an area of 106,000 square miles, none of these rivers are of much importance, being usually ice-locked from September to June, and otherwise flowing through a flat and uninviting region. 3. The American section comprises those streams and rivers which, obeying the northern slope of that continent, find their way through a labyrinth of lakes and swamps, and ultimately fall into the Arctic Sea. The more noticeable of these are the Great Fish, the Coppermine, the Mackenzie, and the Colville. Comparatively little is known of the sources and ramifications of these rivers, of which the Mackenzie is the largest and most important. This great river,

which drains an area of 441,000 square miles, is formed by the union of several streams that rise in the eastern slopes of the Rocky Mountains. The most important of these are the Athabasca and Peace rivers, which, after passing through Lake Athabasca, unite to form the Slave river; and this, after entering the Slave Lake, reissues as the Mackenzie. Like the rivers of Siberia, those of Arctic North America flow through low frozen and inhospitable swamps, and enter the sea by wide mouths; but, unlike those of Siberia, they are connected with a labyrinth of lakes, a feature peculiarly characteristic of the northern section of the American continent.

123. The *Atlantic System*, as the name implies, embraces all those rivers that find their way directly or intermediately into the great basin of the Atlantic from the slopes of the adjacent continents. It necessarily arranges itself into four sections—the European, African, North American, and South American, according as the rivers descend from either of these continents. 1. In the European section we have certain rivers that flow into the Atlantic directly, and others that enter it indirectly through the Baltic and Mediterranean. The chief of those flowing into it directly are the Elbe, Weser, and Rhine, from the western slopes of the Germanic plain; the Seine, Loire, and Garonne from France; and the Duero, Tagus, Guadiana, and Guadalquivir from the Spanish peninsula. The Rhine—the “beautiful Rhine” of the Germans—springs from Alpine glaciers at an elevation of 7650 feet, descends rapidly to Lake Constance (1250 feet), thence through the falls of Laufen to Basel (800 feet), and thence with a navigable course to the North Sea, which it enters by the largest of European deltas. The Rhine has thus a well-marked upper and lower course, a development of 600 miles, and a drainage of 65,000 square miles; the Loire an absolute course of 520 miles, and a drainage of 33,000 square miles; while the Tagus, the third in importance, has a length of 400 miles, and a drainage of 21,000 square miles. The principal streams that discharge themselves through the intermediate basin of the Baltic are the Neva, Niemen, Vistula, and Oder—all more or less obeying a north-westerly slope, and, like their recipient, ice-locked for a considerable portion of the year. Of these the Vistula has the greatest development and drainage—its length being 520 miles, and its basin 56,000 square miles. The waters that fall from the European side into the basin of the Mediterranean are—the Ebro and Rhone into the main sea; the Po into the Gulf of Venice; and the Danube, Dneiper, Dneister, and Don into the Euxine Sea. The Rhone rises among the Alps at an elevation of 5500 feet, and, having an actual course of only 500 miles, is necessarily one of the most rapid rivers in Europe. It has a drainage of 23,000 square miles, discharges itself by two main mouths, and, when in high flood, projects its current with such force that its fresh waters can be skimmed from the surface several miles out at sea.

The waters that enter the Black Sea branch of the Mediterranean basin obey the great southern and south-eastern slope of the European continent. Of these the Don, Dnieper, and Dniester have comparatively gentle currents, and all in their lower courses flow through flat swampy plains. The Danube, the most important of the suite, rises in the Black Forest, at an elevation of 2850 feet, has a winding course of 1496 miles, and an area of 234,080 square miles. Originating in the union of several mountain streams, it is first known as the Donau or Danube in the Duchy of Baden, from which it runs through an Alpine country to Ulm, and thence to Passau it traverses the plain of Bavaria. From Passau to Vienna it runs through a second hilly region, and the remainder of its course is generally through a flat country, except on approaching the rocky defile of the "Iron Gate," till its embouchure by three main mouths into the Euxine Sea. The Danube, with its navigable tributaries, the Theiss, Drave, and Save, form important channels of internal communication for Eastern Europe—an importance that has been greatly enhanced by the adoption of steamers fitted to the peculiarities of their currents. Such are the principal rivers of Europe that fall directly or indirectly into the basin of the Atlantic. Lying in the temperate zone, and having their sources at no great distance from the ocean, their volumes are wonderfully regular and persistent, being merely liable to occasional floodings from excess of rainfall, or from sudden meltings of the snow in early spring. The extent and configuration of the continent prevents the formation of large and long rivers; but as the rapids and waterfalls are confined to their upper stages, the lower portions of their courses are in most instances easy of access and more or less navigable. Whether viewed as channels of drainage and irrigation, or as means of internal communication, these rivers constitute an essential feature in the physical geography of Europe. Flowing from the central region in every direction, they equalise the water-supply more than in any other continent, confer greater beauty and amenity on its surface, and afford at once a perennial supply and an available mode of intercommunication to its busy populations.

124. The African section of the Atlantic system embraces all those rivers which flow from the western and northern slopes of that continent—the former discharging themselves directly into the ocean, the latter indirectly through the basin of the Mediterranean. Of those entering directly into the Atlantic, the Senegal, Gambia, Niger, Congo, Coanza, and Orange are the more important, though still but very partially known in their middle and upper courses. The Senegal and Gambia, which rise in the Kong Mountains, and have courses from 600 to 850 miles in length, form the conjoint basins of Senegambia, and are navigable to some extent. The Niger, the noblest of these western rivers, is supposed to have its sources in the

northern slopes of the Kong Mountains, and, after a circuitous course of more than 1500 miles, during which it receives the Chadda and other large tributaries, enters by several navigable mouths the Gulf of Guinea. The Niger (Joliba or Quorra) drains an immense but unknown extent of tropical Africa; has a navigable middle course of many hundred miles, varying from one to six miles in width, and running at the rate of five and eight miles an hour; and possesses a low pestilential delta of 32,000 square miles, alternately choked with the rankest jungle-growth and overspread by inundation, which attains its height about the middle of August. The Congo and Coanza are so far navigable, but are unknown in the interior, though supposed to be connected with that maze of rivers now known to occupy the central table-land. The Orange or Gariep is unnavigable, being alternately a large impetuous torrent during the rains, and a shallow shingly stream during the season of drought. The most important, and indeed the only African river that discharges itself into the Mediterranean, is the Nile—the most classical, and in many respects the most remarkable river in the world. Rising in the equatorial highlands of Africa as the Bahr-el-Abiad, or White Nile, it receives after a considerable course the Bahr-el-Azrek, or Blue Nile, from the Galla country, and after a further track is augmented by the Atbara from Abyssinia—the conjoint stream flowing downward through Dongola, Nubia, and Egypt (a distance of 1200 miles), without receiving any additional accession of waters. The total length from its unknown source, between 3° and 4° N. lat., is 2240 miles, and its supposed drainage-basin not less than 520,000 square miles. Comparatively little is known of its upper course, or of its source, which, in all likelihood, lies in some mountain lake or swamp near the equator; its middle course through Nubia is marked by a succession of low rapids or “cataracts,” nine or ten in number; and, its lower course having a fall of only two inches a-mile, the current flows gently through the plains of Egypt till its final discharge into the Mediterranean by the two main mouths which form its delta. This delta commences 90 miles from the sea, and has a coast-line of 187 miles between its main mouths, the Rosetta and Damietta. In the plain of Egypt, which is from 2 to 18 miles in width, the current of the Nile, when not in flood, is about $2\frac{1}{2}$ miles an hour, but in its delta branches the flow is almost imperceptible. The most remarkable feature in the Nile is the regularity of its annual inundation, which arises, in all likelihood, from rainfalls dependent on the south-east trade-winds of the Indian Ocean, and on the melting of snows among the equatorial ranges near its sources. In its upper branches the river begins to rise in April, but at Cairo the flood is not perceptible till towards the summer solstice. It then continues to rise for nearly a hundred days, and remains at its greatest height till the middle of October, when it begins

to subside, and reaches its lowest point in April and May. So far as we have evidence, this inundation has remained unchanged for the last 4000 years. Its height in Upper Egypt is from 30 to 35 feet, at Cairo from 20 to 24, and in the northern part of the delta, where it spreads out over a wider area, it seldom exceeds 4 or 5 feet. The fine black slime or mud deposited by this inundation has been the unfailing source of wealth and fertility to Egypt, and from its accumulation, in like manner, has arisen the formation and increase of the delta.

125. The rivers of South America constitute by far the most unique and gigantic section of the Atlantic System. Indeed, all the rivers of that continent obey its great eastern slope towards the Atlantic—the streams that flow down the abrupt and rainless counterslope towards the Pacific being mere runnels fed by the melting snows of the Andes. In this case, we have at once area for development, supply from heavy rainfalls engendered by the moisture-laden winds of the Atlantic, as well as from melting snows and perennial springs, and breadth of volume arising from the flatness of the surface over which these rivers flow. Beginning at the south, the rivers of Patagonia are of little importance, being described as “of small magnitude, with few or no affluents, and making straight across the dry shingly terraces of that sterile region.” North of this the great river-system commences, embracing the basins of the Plata, the San Francisco, the Paranahyba, the Tocantins, the Amazon, Orinoco, and Magdalena. The Rio de la Plata, whose drainage is estimated at 886,000 square miles, receives its affluents from very distant and different regions—the Salado, Vermejo, and Pilcomayo from the deserts and salinas of the Lower Andes—the Paraguay and Parana, two noble rivers, from the wooded slopes and verdant plains of the central regions—and the Uruguay from the low sierras and valleys of Southern Brazil. Some of these affluents, like the Paraguay and Parana, are navigable for hundreds of miles inland, and, on the whole, have deep and accessible channels, but are subject to periodical and destructive inundations. Their united waters meet in the estuary of the Plata—a freshwater sea, 180 miles long, and 120 broad at its entrance; but shallow and loaded with mud, whose discoloration is perceptible far out at sea. The San Francisco, having a length of 1400 miles, and an estimated drainage of 187,000 square miles; the Paranahyba, having a drainage of 115,000 square miles; and the Tocantins, or Para, of 284,000 square miles, are strictly Brazilian rivers, rising among, and flowing between, the sierras of that country. Little is known of their sources, but they descend with considerable currents till they enter the lowlands that fringe the Atlantic, from which they are navigable for long distances inland. The Amazon, by far the largest river in the world—having a length of 3500 miles from its remotest feeder (the Apurimac) in the Andes to its union with the ocean, and

an estimated drainage of 1,512,000 square miles—is that which gives life and character to the lowlands of tropical America. Entering the ocean by an estuary 200 miles long and 130 broad, its freshening influence is felt several hundred miles out at sea; while its main stream, though much impeded by mud-banks and islands, is navigable for 2000 miles inland. Its principal affluents on the right are the Xingu, Tapajos, Madeira, Purus, and Ucayali, which flow from the sierras of Brazil and the central region; and on the left the Rio Negro, the Japura, and Putumayo, which descend from the mountains of Parimé and the northern Andes. Many of these tributaries are largely navigable, and, though treated as mere feeders of the Amazon, would be esteemed important rivers in the geography of other countries. The head-waters of the Amazon drain the entire circle of the middle Andes, almost connecting its stream by a low portage with the Paraguay on the south, and actually uniting it by the Casiquiare with the channel of the Orinocco on the north. Like other tropical rivers, the Amazon is subject to periodical inundation—its waters beginning to rise in December, being at their greatest height in March, and lowest in July and August. When in flood, its waters overspread an immense extent of country, which thus becomes alternately a freshwater sea, and *selvas*, teeming with the rankest tropical vegetation. The Orinocco—the last of South American rivers to which our space will permit us to advert—has a basin of nearly 300,000 square miles, and a navigable course at all seasons of more than 1000 miles. Rising in the mountains of Parimé, it takes a circuitous course northwards and then eastwards, receiving on the one hand several important tributaries from the Andes, and on the other a number of smaller streams from the ranges of the Parimé. Increased by these affluents—the Guaviare, Meta, and Atures on the left, and the Paraqua and Caroni on the right—it ultimately finds its way by a perfect labyrinth of mouths into the basin of the Atlantic. Its annual floods take place with great regularity—commencing about the end of March, and decreasing by the end of August. During high flood, thousands of square miles of the low flat basin are inundated to a depth of 30 feet—these flats, alternately flooded and covered with long rank grass, constituting the celebrated *llanos* of the Orinocco.

126. The river-systems of North America are governed by the disposition of the Rocky Mountains much in the same way as those of South America are governed by the Andes. The counter-slope of the Rocky Mountains, being less abrupt than that of the Andes, gives development to two or three streams of some importance on the Pacific side, but in the main the great river-basins are directed towards the Atlantic. The central plain that stretches from the Gulf of Mexico to the Arctic Sea being divided by a low watershed of 1300 or 1500 feet, a few

rivers, as the Mackenzie and Coppermine, trend northward to the Arctic basin, but all the larger and more important lie south of this watershed, and belong to the Atlantic system. Of these the Rio del Norte and the Mississippi fall into the Gulf of Mexico; the St Lawrence directly into the Atlantic; and the Saskatchewan and Churchill into Hudson Bay. The Rio del Norte, which rises among the sierras of New Mexico, has its upper course through a desert region, in which it receives few affluents; but as it descends into the lower grounds it gains considerable accessions, and becomes a natural boundary between the United States and New Mexico. Its drainage area is estimated at 180,000 square miles. The Mississippi ("father of waters") is by far the largest of North American rivers. Rising in the small lake of Itasca, on the verge of the middle table-land, at an elevation of 1500 or 1600 feet, it runs southward through the great central plain, receiving numerous accessions from either side, and, after a winding course of 3160 miles, falls through a swampy delta into the Gulf of Mexico. Its main tributaries on the right or Rocky Mountain side, in which they take their rise, are the Missouri—a river even longer, larger, and having more affluents, than the Mississippi itself—the Arkansas, having also many affluents, and the Red River; while on the left its chief tributary is the Ohio, with its numerous feeders from the western slopes of the Alleghanies. Between the Rocky Mountains on the west, and the Appalachians on the east, lies, therefore, the great conjoint basin of the Missouri, Mississippi, and Ohio, having an area of more than a million square miles; containing every description of soil and scenery—prairie, barren, and woodland; watered by innumerable streams and several navigable rivers; and possessing every variety of climate, from the coldly-temperate highlands of Nebraska to the sub-tropical warmth of the Gulf of Mexico. The Mississippi, we have said, has a winding course of 3160 miles, but if the Missouri be taken as the main stream, its length is 4260 miles; while that of the Arkansas is 2000, and the Ohio more than 1000 miles. The breadth of the Missouri and Mississippi at their confluence is about half a mile, and the main stream downwards is very little more. Though gentle, the current is by no means sluggish; and thus, during floods, from the melting of the snow in the higher latitudes, the rivers sweep downwards immense quantities of mud, driftwood, and other debris. The Missouri is said to be navigable "from the Great Falls in the Rocky Mountains to the sea, a distance of 4000 miles; the Mississippi, from those of St Anthony, 2240; while the Ohio, being connected by a system of canals with Lake Erie, and thence to Lake Ontario, carries out a water-communication between the Gulfs of Mexico and St Lawrence." The Delta of the Mississippi, which projects itself forward into the Gulf of Mexico, is a low and unhealthy region, nearly 14,000 square miles in area, and full of lagoons, creeks, and pestilential marshes.

The St Lawrence, which forms the great water highway to Canada, takes its rise, under the name of the St Louis, far west in the central lake-region of North America. After uniting the Lakes Superior, Huron, Erie, and Ontario, it issues from the last by the name of the Iroquois, and, alternately contracting and expanding into lake-like reaches, it is known as the St Lawrence at Montreal, from whence it flows to Quebec, and thence by its long estuary into the Atlantic. Its drainage basin is estimated at 297,000 square miles, of which 94,000 or more is covered by freshwater lakes. Its estuary is 350 miles long, and 80 broad at its mouth, but, from its northward trend, is frozen and ice-locked during the winter. The Saskatchewan and Churchill take their rise, in like manner, among the labyrinth of lakes that characterises the extreme northern plain of America, and, after flowing from lake to lake with irregular and tortuous courses, find their way to the North Atlantic through Hudson Bay. The basin of the Saskatchewan is estimated at 300,000 square miles, and that of the Churchill at 73,000, but a large portion of both is occupied by lakes and frozen morasses.

127. Encompassed as the *Pacific* is by the continent of America on the one hand, and by that of Asia on the other, its *river-system* is altogether peculiar, and disproportionate to the encircling areas. On the American side, from Tierra del Fuego to the Gulf of California, there is not a single stream of importance: and from that point northward to Behring Strait the only rivers of note are the Colorado, rising among the sierras of New Mexico, draining, with its tributaries, an arid and rocky area of 170,000 square miles, and falling into the Gulf of Mexico; the Columbia, with its main tributary the Lewis, descending with impetuous current over waterfalls and cataracts from the Rocky Mountains, traversing the Oregon territory, and draining an area computed at 194,000 square miles; and the Frazer, an equally rapid stream, rising among the ranges of the Rocky Mountains, and watering the wild but picturesque country of British Columbia. Abutting as the Andes and mountains of Mexico do upon the very shores of the Pacific, there is no space for the development of rivers; and it is only where the Rocky Mountains bend inland, and are flanked by minor hill-ranges, that the Colorado, Columbia, and Frazer make their appearance. On the Asiatic side of the Pacific the case is altogether different, the mountains from which the rivers descend being not only far inland, and thus affording ample area for development, but being, moreover, the recipients of abundant rain and snow-fall. We have thus, descending the eastern slopes of Asia, the Amoor in Chinese Tartary, the Hoang-Ho, Yang-tse-Kiang, and Tche-Kiang in China, and the Menam and Mekong in Cochin-China—all first-class rivers—besides a vast number of affluents and minor streams that add to the fertility and importance of that region. The Amoor (Tunguse, "Great Water" of the Manchooes, Sagalhién,

or "Black Water") rises high in the Daurian mountains, receives several large tributaries in its descent, and, after a course of 2400 miles, enters as a navigable river into the land-locked Gulf of Tartary, which opens on the one hand into the Sea of Okhotsk, and on the other into the Sea of Japan. It drains a computed area of 583,000 square miles; and, from its navigable capabilities, safety of entrance, and relative position to Japan and Russian Siberia, is likely to become of considerable commercial importance. The great rivers of China—the Hoang-Ho and Yang-tse-Kiang—take their rise among the Kihan-Schan and Kuen-lun mountains, that buttress the plateau of Tartary, and after making their descent between the minor ranges of the Yun-ling and Peling, wind with slow and steady flow through the plain of China into the Yellow Sea. The course of the Hoang-Ho, or Yellow River (so called from the colour of its waters), is estimated at 2300 linear, and its basin at 537,000 square miles; while the course of the Yang-tse-Kiang is 2900 linear, and its drainage area not less than 548,000 square miles. Though their main mouths are wide apart, yet they may be said to fall into the same delta—their broad navigable streams being united throughout their lower courses by innumerable canals and natural channels. Indeed the whole country, from the southernmost branch of the Yang-tse, north to the Pei-ho, or White River, which enters the Pe-che-le Gulf, is one alluvial flat, intersected by channels, canals, and embankments, and liable to be flooded and broken in upon by shiftings of the river-courses, and other similar changes. The burden of sediment which these rivers carry down to the shallow basin of the Yellow Sea is immense; and though frequently shifting their channels, and encumbered by sand-banks, they are navigable by vessels of considerable burden for 600 or 800 miles, and by smaller craft to perhaps double that extent. The Mekong is a large but little known river, rising in the mountains of Eastern Assam, flowing with a long course between the mountain-ranges of Cochinchina, draining, along with the Menam, an estimated area of 216,000 square miles, and falling into the Gulf of Siam or Cambodia by a many-mouthed delta.

128. *The river-system of the Indian Ocean* has three separate and independent sections—the Asiatic, the Australasian, and the African, according as the rivers flow from either of these regions. 1. The Asiatic embraces the Martaban and Irawady in Burmah; the Brahmapootra, Ganges, and Indus in India; and the Tigris and Euphrates in Persia and Asia Minor. The Martaban, Irawady, and other rivers of Further India, occupying the narrow valleys which lie between the parallel hill-ranges of that region, flow southward in long and comparatively straight courses. Very little is known of their sources or affluents; but the Irawady, which enters the Gulf of Martaban by a many-branching delta, is navigable for 400 or 500 miles, and though encumbered in its lower course by mud-banks and islands, is a

channel of vast importance to the Burmese empire. During the season of inundation it spreads to a breadth of 3 or 4 miles, runs at the rate of 4 or 5 miles an hour, and is heavily laden with sedimentary debris. There is, indeed, a great similarity among all the rivers of the Indo-Chinese peninsula, whether flowing like the Menam and Mekong into the Pacific, or like the Martaban and Irawady into the Indian Ocean: they have long straight courses, flow with considerable current through a country rich in tropical vegetation, and enter the sea by several mouths through low and gradually increasing deltas. The Ganges, which rises at an elevation of 13,000 or 14,000 feet among the glaciers of the Himalaya, descends rapidly to the plain of India, and there flowing with gentle current it receives numerous affluents from the southern slopes of its parent mountains on the one hand, and from the northern slopes of the Vindhya high grounds on the other, till it ultimately falls through a many-branched delta into the Bay of Bengal. The Brahmapootra "offspring of Brahma") is formed by the union of two main streams, the Dzangho-chur, which has its origin in the northern slopes of the Himalayas and flows eastward till its union with the Brahmapootra proper, which, descending from the distant recesses of the Thibetan mountains, runs southward towards Assam. After their junction the united waters, under the name of the Brahmapootra, cut transversely the eastern Himalayas, and then, flowing south and westward with a volume considerably exceeding that of the Ganges, enter the delta of that river about 40 miles above the sea, but still maintain the individuality of their own channel. The conjoint area drained by these rivers and their affluents (several of which are larger than the Rhine) is estimated at 432,000 square miles, and their respective lengths at 1680 miles. Like other tropical rivers, the Ganges and Brahmapootra are subject to annual inundations—the floods commencing in April, attaining their maximum about the middle of August, and continuing till October. They begin first to swell from the melting of the snows among the mountains, but before this influence has reached the low grounds these are widely under water from the periodical rainfalls. The quantity of water arising from these two causes and brought down by the Ganges and Brahmapootra is enormous, overspreading the plains for hundreds of square miles, and freshening more or less the whole upper area of the Bay of Bengal. The amount of sediment brought down by these rivers when in flood is also immense; and the whole delta, 200 miles in length and 180 broad at its base, with all its channels, creeks, lagoons, and mud-islands (Sunderbunds), is clearly the offspring of this debris. Though liable to sudden shiftings during inundations, several of the branches of the Gangetic delta are navigable at all seasons for vessels of large draught; while the main stream can be ascended by smaller craft to the foot of the Himalayas. Though possessing a large

volume, the Brahmapootra, from the rapidity of its current and the obstacles in its channel, is of less importance as a means of internal communication. The Indus, the third great river of India, takes its rise in the highlands of Thibet, and after a precipitous course through the western Himalayas, descends into the plains, where it is augmented by the streams of the well-known Punjab, or Five Rivers. These affluents, like the main stream, are the offspring of the Himalayas, and drain a large expanse of upland as well as fertile lowland. After their union near the southern extremity of the Suleiman hills, the Indus receives no more tributaries of any note, but flows with gentle current through a somewhat arid country for nearly 300 miles, and then forks into a delta 80 miles long by more than 100 broad at its base, discharging itself by many mouths, only three or four of which are navigable. The annual floods commence with the melting of the Himalayan snows in April, attain their greatest height in July, and terminate in September. The winding course of the river is nearly 2000 miles, and its estimated drainage 312,000 square miles; but owing to the shallow and shifting nature of its mouths it is navigable only by vessels of comparatively small burden. Lying open to the great tidal wave of the Indian Ocean, it has, like the Ganges, a rapid and dangerous *bore*, which ascends the main channels to a distance of 60 or 70 miles.

129. Australia, owing to its geological conformation, is singularly destitute of rivers—those descending the eastern counter-slope of its hills into the Pacific being mere streams, while those flowing from their western slopes seem to lose themselves in the great central and yet unexplored plain. The Murray, with its tributaries the Darling, Lachlan, and Murrumbidgee, are the only known rivers of note, and these, though roaring torrents during the rains, are often reduced to a mere chain of ponds and creeks in the dry season. Of the African rivers that enter the Indian Ocean geography knows too little as yet to offer anything like reliable description. So far as known, the Zambesi, now under exploration by Dr Livingstone, is the largest and most important, apparently draining an immense extent of inland country, which spreads out in plain-like expanses, reticulated by lakes and streams that flow during the periodical rains, but are desiccated and partially obliterated during the season of drought. On the whole, recent investigation seems to point to the interior of Central Africa as a plateau-shaped plain, deriving its main water supply from rains, which form, for the most part, only temporary streams and lakes—the surplus being carried off by evaporation rather than by a regular system of stream and river-drainage.

130. Such is a brief outline of the chief river-systems of the world—the Arctic, Atlantic, Pacific, and Indian. Though the

basins of which they are composed have their own individual characteristics, there is still a great similarity, and necessarily so, among the various members of a system. Thus the rivers of the Arctic system run more or less meridionally, flow from *warmer to colder* latitudes, and all their embouchures are for a large portion of the year ice-locked and impervious to navigation. By the melting of the snows in their upper courses they are flooded before the thaw has reached their lower waters, and thus the plains through which they pass are largely occupied by lakes and morasses. The rivers that descend the southern slopes of Asia, Europe, and North America, flow also in meridional courses, but pass, on the contrary, from *colder to warmer* latitudes, and consequently their embouchures are open at all seasons. Those of them that belong to the Atlantic system derive their floods chiefly from the melting of the snows along their upper courses; while the inundations of those belonging to the Indian system depend partly on the melting of the snows and partly on the periodical rainfalls in their lower courses. With the exception of the Rio del Norte and the Mississippi on the one hand, and the rivers that enter the Mediterranean on the other, all the basins of the Atlantic system, whether from the American, European, or African continents, run in latitudinal courses; and thus, from their sources to their embouchures, each passes along the same climatological zone. These and other great conditions must at once present themselves to the mind of the inquiring student, and produce the conviction that the idea of *river-systems* has something more to recommend it than the mere convenience of artificial arrangement.

131. Besides the oceanic systems there are what have been termed *Continental Systems*—a few inland basins cut off from all connection with the ocean, and having the equilibrium of their waters maintained by evaporation and absorption. The chief of these, as already mentioned, are the Aralo-Caspian basin in Asia; the Utah and Mexican in North America; the basin of the Andes plateau in South America; and in all likelihood, when better known, some similar depressions in Australia and Central Africa. On referring to the map it will be seen that a number of the streams of Central Asia—the Asiatic plateaux—run inland to lakes or lose themselves in the sand and shingle of the desert. This is the “Basin of Continental Streams” in Asia, and includes the Lob, Balkash, Helmund, and other lake-hollows, with their tributary streams; the Aral, with its main feeders the Syr and Amoo; and the Caspian, with its rivers the Oural and Volga. Of these the most important is the Volga, which rises on the Valdai slopes at an elevation of 550 feet, and, after a long and winding course of 2400 miles, during which it receives many important tributaries (the Kama, Viatka, &c.), falls by two main mouths into the Caspian Sea, which is 82 feet below the level of the Euxine. The Volga is the largest river in Europe, has a fall

of only 633 feet during a course of 2400 miles, drains an estimated area of 397,640 square miles, and constitutes the great internal water-way of Russia. The seas and lakes of this basin, as indeed of all other continental basins, having no escape for their waters save by evaporation, must necessarily be more or less salt; and hence the saline peculiarities of the Caspian, Aral, Balkash, Dead Sea, Lake of Utah, and others occurring in these so-called "basins of continental streams." Basins of this kind necessarily occur in rainless flats and plateaux; for were the amount of rainfall exceeding the evaporation, the sheets of water would accumulate till they overflowed their margins, and found an outlet towards the nearest oceanic area.

Lakes and Lacustrine Areas.

132. *Lake* is the general term for any considerable body of standing water surrounded by land, and not directly connected with the sea or any of its branches. Lakes are thus strictly terrestrial expanses, and belong to the land as much as the streams and rivers that channel its surface. Wherever there is a depression of that surface beneath the surrounding country or the bed of the nearest river, there water will accumulate (unless carried off by evaporation) till it rises above the lowest part of the enclosing margin and flows off by some river-channel. In this way we may have sheets of all sizes, from the merest pools to lakes occupying many thousands of square miles. Lakes generally consist of *fresh water*; but some, like the Caspian and Aral, having no river of discharge, are *salt*, while others, having only partial outlets, are *brackish*. They occur in all regions, but most abundantly on mountain table-lands and plateaux, and on the lower reaches of great plains. Wherever there is a want of declivity, or any obstruction to the natural flow of the surface-waters, there lakes will accumulate, and this in proportion to the extent of country, the magnitude of the obstruction, and the amount of rain and snow-fall. In consequence of less evaporation, they occur also more abundantly in low than in high latitudes; and hence their frequency and magnitude in the northern regions of America, Europe, and Asia, as compared with other areas. It is usual to arrange lakes into *four* kinds:—1. Those that have neither outlet nor inlet—subterranean springs and rain supplying the water, and evaporation carrying off the excess. These are generally of small dimensions—lowland pools, mountain tarns, and not unfrequently the craters of extinct volcanoes. 2. Those which have an outlet, but receive no running water, being fed by springs rising from their bottoms and rocky margins. Lakes of this class are also small, and for the most part situated in upland districts. 3. Those which, like the Caspian and Aral, receive streams of running water, but have no visible outlet—the

balance of level being maintained by evaporation. Such lakes are more or less impregnated with saline matter; and this saltiness must be still on the increase. 4. Those that both receive and discharge streams of running water, and which form alike the most numerous and most extensive in both hemispheres.

133. To whatever class they belong, lakes form essential elements of diversity in the landscape, and perform important functions in the economy of nature. Exposing considerable surfaces to evaporation, they serve to temper the aridity of their respective districts, at the same time that they act as so many reservoirs, in which the superabundant supplies of winter are stored up for the increased requirements of summer. In many instances they act as checks to the too rapid discharge of rivers—retaining for perennial supply what would otherwise be run off in a few days, and restraining, moreover, the destructive flood which is brought to rest in their placid areas. Occurring so frequently in the course of rivers, they act as settling pools for the debris and sediment of their waters—the streams they discharge being pure and pellucid, whilst those they receive may be turbid and laden with impurities. In this way they get gradually silted up, and form rich alluvial tracts, the while that their outlet-currents are deepening their channels and forming the means of a more efficient drainage. In this way lakes become important agents in the surface-modification of the land; and one has only to cast his eye over the fertile dales and vales of long-established regions to perceive how much of these areas was at one time a mere succession of lakes and morasses. Biologically, too, these fresh-water sheets become the habitats of a peculiar flora and fauna—thus extending the range of Life, and affording conditions of existence that no other habitat could supply.

134. In Europe there are two main lake regions—a highland and a lowland; the former embracing the picturesque lakes of Britain, Switzerland, and Italy; the latter the numerous sheets that stud the Baltic provinces of Russia, Prussia, and Sweden. The highland lakes generally occupy the deep narrow troughs of mountain-glens, and from their situation and adjuncts are celebrated for their scenery; the lowland, on the other hand, spread over areas little depressed beneath the surrounding country, and are for the most part tame and unattractive. The waters of Highland or Alpine lakes are usually retained by rocky barriers, through which the outflowing stream is gradually deepening its channel, and thus their dimensions become less and less as the drainage outlet becomes deeper. On the contrary, the drainage of lowland sheets is often slow and sluggish; hence a large portion of their areas is little else than morass, which the accumulation of aquatic vegetation and sedimentary matter is year after year converting into alluvial land. With two or three exceptions (Platten See, Niesidler See, &c.), the European lakes, strictly so called, consist of fresh water; they have all or nearly

all, streams of ingress and egress ; vary in size from a few hundred square yards to several thousand square miles ; rise slightly with the rains of winter and fall with the droughts of summer. The more notable of those in the northern lowland region are Lakes Ladoga 6330 square miles, Onega 3280, Saima 2000, Peipus 1250, and Enara 1200 square miles, in Russia ; and Lakes Wener 2130, Wetter 840, and Maelar 760 square miles, in Sweden—the latter being only three or four feet above the level of the Baltic. The larger Alpine lakes are on the Swiss side ; Geneva or Leman, 240 square miles, at an elevation of 1230 feet ; Constance or the Boden See, 228 square miles, at an altitude of 1250 feet ; Neufchatel, 114 square miles and 1440 feet high ; Lucerne, 98 square miles and 1430 feet high ; and Zurich, 74 square miles, at an elevation of 1330 feet : while on the Italian side there are Garda 182, Maggiore 150, and Como 66 square miles, at elevations respectively of 320, 680, and 685 feet. In the British Isles, Loughs Neagh, Corrib, Erne, and Loch Lomond, respectively 150, 63, 56, and 45 square miles in area, and 48, 31, 130, and 22 feet in elevation, are the largest, though others of smaller dimensions are better known for their picturesque scenery and associations.

135. In Asia, the lakes occupy three main and distinctive regions—the mountains and plateaux of the central highlands, the alluvial lowlands of China and Siberia, and the depressed areas of the Aralo-Caspian and Dead Sea. Unlike those of Europe, many of the Asiatic lakes are salt or brackish, and this peculiarity belongs more especially to those having no river of outlet. Respecting these salt or brackish sheets, accounts differ considerably as to the amount of saline matter held in solution in their waters—a difference that may arise either from taking the specimens at different places, at different seasons, or from errors in analyses. Whatever the source of error, there can be no doubt that where new saline supplies are continually carried in by rivers and springs, and none carried off by running water, the percentage must be gradually on the increase. Of the *saline* lakes the more important and better known are the Caspian (slightly fresher than the ocean), 82 feet beneath the level of the Black Sea, having an area of 130,000 square miles and an extreme depth of 960 feet ; the Aral, 36 feet above sea-level, and having an area of 26,000 square miles ; Balkash or Tengiz, having an area of 7000 square miles ; Urumiah (25 per cent of salts), 1800 ; Van, 1600 ; Zurrah, 1600 ; Tongri-nor, 1800 ; Koko-nor, 1500 ; Lop, 1300 ; and the Dead Sea, 360 square miles, with a depression of not less than 1312 feet beneath the level of the Mediterranean, and containing from 25 to 26 per cent of saline ingredients. Of the *fresh-water* sheets the largest is Baikal (the “Holy Sea” of the Russians) in Siberian Tartary, with an area of 14,800 square miles, and at an altitude of 1535 feet ; and Ton-Ting, Poo-Yang, and Tai-Hou in China, having respectively areas of 2000, 800, and 700 square miles, and all valuable aids to the internal communication, com-

merce, and agriculture of that peculiar country. Besides these may also be noted Zai-zan in Mongolia, 1000 square miles; Bookanor in Tibet, 1000; Erivan in Armenia, 500; and Tiberias in Syria, 76 square miles, at a depression of 328 feet beneath the Mediterranean.

136. In Africa there are several lakes less or more known to geographers; but while the internal configuration of that continent remains undetermined, it is impossible to arrange them into anything like systems or areas. Between the Atlas range and the Sahara, and fed by streams from the former, there exists a series of minor lakes, of which the largest (Loudeah, &c.) lies in Tunis; in the central and hilly region of Nigritia the existence of sheets of considerable magnitude (Tchad, Dibbie, Fittie, &c.) has been so far determined; and in the southern table-land, several shallow and apparently fluctuating sheets (Ngami, Nyanja, Taganyika, &c.) have been partially explored. Among the longer known lakes may be mentioned Dembea or Tzana in Abyssinia, at an elevation of 6270 feet, and embracing an area of 1400 square miles; the deltoid areas of Menzaleh and Mareotis in Egypt; and the salt-water lake of Assal in Abyssinia, extending to 30 square miles, and lying in a depression 570 feet beneath the level of the Red Sea. With regard to Australia, we know that certain areas covered with water during the rains are but marshes or mud-pools during the season of drought, and that, along many portions of the sandy coasts, there exist saline and brackish lagoons; but we are yet too slenderly acquainted with the interior to speak with certainty as to the existence of *lakes* in the true sense of the term either fresh-water or saline.

137. In South America there are comparatively few lakes, and these, for reference sake, may be arranged into three distinctive regions—namely, the mountain-glens and depressions of the higher Andes; the low plains that border the eastern flanks of the Andes; and the still lower river-plains that trend toward the Atlantic. Of the Andian lakes the most remarkable is Titicaca, situated at an elevation of 12,847 feet, embracing an area of 3800 square miles, and stamping peculiar features on the scenery, life, and civilisation of the surrounding district. It is fed by several streams from the Andes, but discharges itself by a single outlet, the Desaguadero, which after a short course disappears, partly by evaporation and partly by absorption into the sandy and arid soil through which it flows. Of the lakes that skirt the Andes on the east, many of them are of temporary nature, being extensive sheets during the rains, but mere marshes and salt pools (*salinas*) during the period of drought. In the lower plains of the Plata, Amazon, and Orinocco there exist many creeks, lagoons, and stagnant areas that appear in the dry season as lakes, and during the rains as one vast area of inundation.

138. In North America the lakes may be said to affect only two main districts—a highland and a lowland; the former the

troughs and depressions of the Rocky Mountains from Panama on the south to Columbia on the north, and the latter the northern section of the great central plain from the southern boundaries of Canada to the Arctic Ocean. In the former area the more important sheets are Nicaragua, 3500 square miles, at an elevation of 128 feet, Maragua, 450 square miles, and Yojoa, 150 square miles, in Central America; Chapala in Mexico, 1000 square miles; and in the United States, Utah, 150 square miles, and the Great Salt Lake, 1800 square miles, at an elevation of 4200 feet. Of the lowland section the more important and extensive are the great "fresh-water seas" that form the lacustrine chain of the St Lawrence. These are Superior, with an area of 32,000 square miles, an altitude of 628 feet, and a depth of 900 feet; Michigan, 24,000 square miles, altitude 574 feet, and depth 1000 feet; Huron, 20,000 square miles, altitude 574, and depth 1000 feet; Erie, 9600 square miles, altitude 565, and depth 84; and Ontario, 6300 square miles, altitude 232, and depth 500 feet. Besides these, there is the immense lacustrine net-work of the extreme north, forming with that of Canada a lake-region unsurpassed by any other on the globe. Of the larger of these boreal, shallow, marshy, and long-frozen lakes we may enumerate Great Bear, 19,000 square miles; Great Slave, 12,000; Winnipeg, 9000; Winnipegosis, 3000; Athabasca, 3000; Deer, 2400; Manitoba, 21,000; Wollaston, 1900; and Lake of the Woods, 1200 square miles.

RECAPITULATION.

In the preceding chapter attention has been directed to what may be termed the *terrestrial* waters of the Globe—that is, to the springs, streams, rivers, and lakes that occur on the land in contradistinction to the great united mass which constitutes the Ocean. Though separable in this way, it is still the same vast volume of waters we have to deal with, its appearance on the land being only local and temporary. The vapour exhaled from the ocean is conveyed to the land, where, condensed into rains and snows, it falls and forms springs and streams, and these by their union give birth to rivers and lakes. But the stream and river are ever hurrying to the ocean again—again to be evaporated as *fresh water*, again to fall as rain and snow, and again to course the dry land, and to carry with them a portion of those ingredients which, accumulating within certain limits, constitute the peculiar *saltiness* of the ocean. The rain and snow that percolate the rocky crust reappear at the surface as *springs*, which, according to their situation, may be superficial or deep-seated; accord-

ing to their supply, temporary, intermittent, or perennial ; according to their temperature, cold or thermal ; and according to the rocky material through which they pass, either pure or mineral—that is, impregnated less or more with saline, calcareous, siliceous, chalybeate, and other mineral and metallic substances. By their union, springs and surface rivulets form *streams*, and these streams occurring most abundantly in hilly regions unite, one with another, as they descend to the plains, and there ultimately form *rivers*, whose continually augmented volumes constitute one of the most important features in the hydrography of the globe. The highest or farthest-off spring to which we can trace a river constitutes its *source* or *rise*, and from this source, through all its windings till it finally falls into the sea by one or by many mouths, lies its *course*. It is usual to divide river-courses into upper, middle, and lower stages—the upper being characterised by ravines and waterfalls, the middle by rapids and cataracts, and the lower by a broad gentle current which often finds its way into the ocean by several branches through a low-lying and marshy delta. The track worn or hollowed out by running water constitutes its *channel*, and the land that forms the margins of this channel is known as its *banks*—that on the right hand as we descend the current being termed the right bank, the left hand the left. The streams that enter any river or main-current from either side form its *affluents* or *tributaries*, and the entire expanse of country drained by a river and its tributaries is said to be the *basin* of that river. An ideal line, connecting the ultimate sources of all the tributaries of a river, forms the *watershed* of its basin ; and generally speaking, the ridge from which the streams of a country flow in opposite directions is the watershed of that region. As all the streams of a river-basin fall to the lower level of that basin, so several river-basins may descend to the still broader basin of some sea or ocean, and all such basins discharging themselves into the same sea constitute a *river-system*. The river-systems are necessarily coincident with the great oceans into which they flow ; and thus we have the Arctic, the Atlantic, Pacific, and Indian systems, together with a few inland systems whose streams flow towards certain depressed areas altogether cut off from connection with the ocean. These are the Aralo-Caspian basin, the Utah, the Mexican, the Bolivian, &c., whose waters flow inwards to certain lakes, and are thence carried off by evaporation. Where the slopes of a country are pretty uniform, the surface waters are at once carried off by streams and rivers ; but where any unusual depression occurs, there they accumulate and form

lakes and morasses. Most of these lakes both receive and discharge running water; but some of a small size, being fed by springs and reduced by evaporation, neither receive nor discharge running water; while others again receive streams and rivers but have no outlet—their surplus waters being carried off by evaporation. The former set of lakes are always fresh; the latter, like the Caspian, Aral, and Dead Sea, always less or more brackish and saline.

X.

CLIMATOLOGY—THE ATMOSPHERE.

Nature and Constitution of the Atmosphere.

139. HAVING, in the preceding chapters, directed attention to the leading features of the Land and Water—the mountains, table-lands, and plains of the one, and the seas, lakes, and rivers of the other—we now proceed to consider the main relations of the Atmosphere or aerial envelope (Gr. *atmos*, vapour, and *sphaira*, a sphere) by which the whole is surrounded. What is the nature and composition of this envelope, what its capacity for heat and moisture, what are the motions to which, as an elastic fluid, it is subjected, and generally, what are its conditions as the great medium of climate (heat, cold, winds, rains, &c.) on which the Life of the globe is so intimately dependent? This varied subject of climatology belongs more especially to the science of Meteorology; but as an integral portion of the globe, and bearing on everything—organic and inorganic—that presents itself at the surface, the atmosphere and its principal phenomena become important themes to the student of Physical Geography.

140. As already stated in paragraph 17, the atmosphere surrounding the globe, and partaking in all its motions of revolution and rotation, is an aëriform fluid consisting of nitrogen and oxygen gases—79 parts of the former to 21 of the latter by volume, together with a small and variable percentage (.05 to .10) of carbonic acid gas and other impurities. These gases are merely in a state of admixture or diffusion, as is also the invisible vapour which less or more is ever present in the atmosphere. This special composition is alike indispensable to vegetable and animal life. Both animals and plants breathe, if we may so speak, the air; but while carbonic acid is exhaled by the former, it is absorbed and assimilated by the latter, which in turn exhale oxygen for the requirements of the animal kingdom. By this means the balance of composition is harmoniously maintained, any local variations being too unimportant to affect the general result. Being an elastic or compressible medium, the air nearest the sea-level, pressed upon by all the superincumbent mass, is denser

than that at considerable elevations; and by calculating the rate at which this rarity takes place, it is estimated that, at the height of 45 or 50 miles, the atmosphere becomes so rare or attenuated as to be inappreciable. What the *absolute* height or extension may be is still an undetermined problem—90 miles, 212 miles, and even greater altitudes being given, according to the principles on which the calculations are founded.

141. The mean pressure of the atmosphere taken at the sea-level is usually reckoned at 15 lb. avoirdupois (14.7304 lb.) upon the square inch—a pressure which, being equal on all sides, is insensible to living structures, the only inconvenience being experienced at great altitudes (5 or 6 miles), where the air becomes too rare and light to maintain respiration, or to retain sufficient heat for the functions of vitality. This aerial pressure is balanced by a column of mercury 30 inches in height; hence the construction of the *barometer*, and hence also the term *barometric pressure* as applied to the oscillations of the atmosphere, which sometimes rise above and at other times fall below this normal standard. The pressure or weight diminishing as we ascend, the barometric column will fall in proportion, and this fall (correction being made for varying effects of temperature) becomes a convenient measure of altitude. Thus at the sea-level, near the foot of Chimborazo, Humboldt found that the barometer stood exactly at 30 inches, while at the elevation of 19,332 feet, to which he ascended, it fell to 14 $\frac{3}{4}$ inches. Again, under the normal pressure of the atmosphere, water at the level of the sea boils at the temperature of 212° Fahrenheit; but as the pressure becomes less, ebullition takes place sooner, and thus, as we ascend, the boiling point diminishes, and may in like manner be taken as a measure of altitude. Of course, water at 212° and water at 190° (the boiling point at 13,000 feet in the Andes) will produce very different effects, absolute temperature and not mere ebullition or bubbling-up under diminished pressure being the test of its heat.

142. The atmosphere being the medium through which the sun's heat is conveyed to and from the earth, the lower and denser strata absorb the greatest amount and are necessarily the warmer; hence, as a general rule, temperature decreases with the height to which we ascend from the surface, but not uniformly, as has been established by the recent balloon ascents of Mr Glaisher. As air is expanded by heat and contracted by cold, the warmer and lighter volumes will ascend, and the colder and denser rush in from all sides to supply the vacancy. In this way currents or *winds* are originated in the atmosphere, and on these currents depend in a great measure the essentials of climatic diversity. As a property of the gaseous or aëriform nature of the atmosphere, the vapour or minute watery vesicles, generated from the earth's waters by the heat of the sun, ascend and are diffused through its mass in an invisible state. This humidity of the

atmosphere also decreases with the height, and this in so rapid a ratio that at the altitude of 5 or 6 miles it is all but inappreciable. These vapours, invisible at first, become visible on being condensed by colder currents; and hence arise fogs, mists, rains, snows, and other kindred phenomena. The effective portion of the atmosphere, so far as its heat and moisture are concerned, lies, therefore, within a few thousand feet of the earth's surface, all above the limit of 5 or 6 miles being inoperative or nearly so in the production and regulation of climate. As the atmosphere is the medium through which the sun's heat is conveyed to and disseminated over the earth, so also it is the medium of his light-giving rays. Heat and light are alike indispensable to plants and animals, and, from the peculiar constitution of the atmosphere, as regards its varying density, moisture, &c., both are reflected and diffused so as to become most available to vegetable and animal life.

143. Had the globe presented either one uniform surface of land or one uniform surface of water to the sun's rays, the interchanges of heat and moisture between the colder and warmer, the dryer and moister regions of the atmosphere, would have been regular and continuous. But the earth's surface being composed of land and water in unequal proportions, and this land lying not only in irregular masses but presenting also irregular surfaces of highland and lowland, the movements of the atmosphere are so broken up and complicated that the winds and rains, the heat and cold, are rendered extremely variable—so variable, indeed, that Meteorology consists as yet more of collections of facts and approximations to results than of ascertained and definable laws. Under these circumstances we shall glance merely at the leading features of the atmosphere—its heat, moisture, and motions—as bearing more directly on the *climate*, or weather-conditions, if we may so express it, of the earth's various regions.

Its Heat and Moisture.

144. In dealing with the subject of climate, the superficial or atmospheric temperature of the globe may be viewed as wholly dependent on the heat of the sun—the amount conveyed from within (paragraph 16) or radiated from the moon and other external sources being by far too feeble to have any perceptible influence. As is well known, the solar rays exercise the greatest effect within the tropics, where they fall vertically or nearly so on the earth's surface, and that this effect gradually declines as we proceed through the temperate zones towards either pole. In other words, the higher the angle at which the sun's rays strike the earth the greater their heating power, and the lower the angle the less the amount of heat imparted. We feel the truth of this every summer day, when the sun is high a-head during

the scorching noontide, and when his beams fall at a lower angle during the cooler afternoon. At whatever inclination they fall, there is always, as before explained, a certain amount of heat lost or extinguished in passing through the atmosphere; and this amount for nearly vertical incidence within the tropics has been estimated at one-third; for inclinations of less than 25 degrees at fully one-half; while at inclinations of 5 degrees scarcely one-twentieth of the solar heat is supposed to reach the surface. In this way $23^{\circ}44'40''$ on either side of the equator forms a belt within which the solar heat received is equal to what is received by all the rest of the world; and hence the learner will perceive the applicability of the terms "torrid," "temperate," and "frigid" zones. Keeping out of view all minor modifications produced by the peculiar disposition of sea and land, as well as by the sun's alternate passage to the northern and southern hemisphere along the course of the ecliptic, it may be received as the first great axiom, that the superficial temperature of the globe is highest under the vertical incidence of the sun's rays near the equator; and that it gradually becomes lower and lower as we proceed towards either pole, and this in proportion to the obliquity at which the heat-giving rays fall upon the surface.

145. The heat that falls on the land being partly absorbed and partly radiated into the atmosphere, the lower aerial strata, or those nearest the influence of this radiation, will be warmer than those at higher elevations. The rate at which this diminution takes place depends on many correlative circumstances—dryness, moisture, currents &c.; but as a general rule, the decrease may be regarded as equal to 1° of Fahr. for every 300 or 350 feet of ascent. In this way the lower plains of a region may be teeming with vegetation while the higher mountains are enveloped in perpetual snow. We have thus two great causes affecting the superficial temperature of the globe—distance from the equator, and elevation above the level of the sea; so that at any given latitude it is much the same whether we travel towards the nearer pole or ascend a lofty mountain. In both we will meet with a gradually decreasing temperature, in both we will meet with a similar change in animal and vegetable life, and in both we will ultimately reach a limit of perpetual ice and snow. In the case of elevation, this limit is known as the *snow-line*, at which the air in summer attains the temperature of freezing water, and though slightly modified by surrounding circumstances, it varies primarily with the latitude—ascending from 2000 feet within the polar circle to 15,000 or 16,000 feet in the vicinity of the equator. In the case of horizontal extension, the *line of constantly frozen ground* differs in either hemisphere—that in the northern (see Map of Isotherms) assuming an irregular course, in consequence of the unequal distribution of sea and land, and that of the southern being less determinable in consequence of the greater expanse of Antarctic Ocean.

146. Another cause of variation in atmospheric temperature is the unequal reception and radiation of the sun's heat by the respective surfaces of land and water. The heat falling on the land is partly absorbed and conducted downwards into the soil, and partly radiated into the atmosphere. The amount absorbed will differ according to the colour, slope, and character of the land (dry, firm, and dark-coloured soils being most favourable for absorption); and according also as it is naked or shut out from the sun's rays by a covering of vegetation. The amount radiated differs with almost every passing condition of the atmosphere—radiation going rapidly forward under a still, clear sky, while a cloudy atmosphere, or even a passing mist, serves to retard it. The greater quantity of heat absorbed by day is given off by night; and the amount accumulated during summer is returned to the atmosphere during winter. The depth thus affected by summer's heat or winter's cold will differ according to latitude, soil, and situation; but, on the whole, it may be safely asserted that beyond 60 or 80 feet the temperature of the earth's crust is altogether unaffected by external influences. Indeed, for all practical purposes the superficial temperature of the land, as affecting climate, may be regarded as confined to a very few feet beneath the surface—the slow conducting power of the soil and capillary attraction of moisture from below combining to limit its downward tendency.

147. On the other hand, the heat that falls on the water is much more readily absorbed but more slowly radiated, so that, being diffused by waves and currents, it penetrates to a considerable depth, and accumulates within the mass of the ocean. The heat that falls on any zone of the land is absorbed and radiated wholly within that zone; whereas that falling on any zone of the ocean is diffused and interchanged between every other zone. As the ocean radiates its heat more slowly than the land, and exposes more than thrice the surface to the sun's rays, it becomes, as it were, a great storehouse of heat for the exigencies of the land. It is for this reason that islands and seaboards possess a milder and more equable climate than the interior of continents—being warmer during winter and cooler during summer. The British Isles, with their comparatively cool summers and mild variable winters, are thus said to possess an *insular climate*; while the interior of Germany, with its excessively hot summers and cold winters, is said, on the contrary, to possess a *continental climate*. In tropical latitudes, unaffected, of course, by winter, the interior of continents is always much warmer than the sea-coasts and adjacent islands. It is also for this reason that the northern and southern hemispheres, having unequal distributions of sea and land, are differently affected in their zones of climate—the greater extent of land in the northern imparting to it, as it were, a continental climate; while the greater expanse of ocean in the southern confers on its similar latitudes more of an insular climate.

It is, further, to this unequal reception of heat by land and water that we owe the phenomena of sea and land breezes. During day the atmosphere immediately above the land is more rapidly heated by radiation than that above the ocean, and hence the cooler sea-breeze sets in towards the land to restore the balance; while during the absence of the sun the more rapidly cooled land atmosphere sets out as a land-breeze towards the more slowly cooling surface of the ocean. In fine, it is mainly owing to the unequal distribution of sea and land, and to the different modes in which the two elements are affected by the sun's heat, that we owe the great essentials of climatic diversity—the air and ocean currents, the vapours and rains, of one region varying from the winds, currents, and moisture of another.

148. Whatever the temperature of the air, it is incessantly receiving *moisture* from the surface of the land and water. This vapour arises and is diffused through its mass in an invisible form, and becomes visible only when condensed into mists, fogs, rains, and snows. It arises alike from land and water, but most copiously and persistently, of course, from the surface of the water. The superficial moisture of the land, unless where clothed with vegetation, is rapidly converted into vapour; but the supply from this source is irregular and limited compared with that derived from the water. Surface for surface, salt water is less vaporisable than fresh; hence a slight check to evaporation from the vast and extended surface of the ocean. It rises from rivers, lakes, and seas in every latitude, but most abundantly, of course, within the tropics, where the sun's heat is the greatest. So great is the evaporation within that zone, that it has been estimated as annually sufficient to reduce the surface of the sea to a depth of 8 or 10 feet—a waste, of course, that is incessantly counterbalanced by influx from other latitudes. The warmer the air the greater its capacity for moisture; and this capacity for holding moisture in suspension increases up to the point of saturation, when any reduction in temperature produces condensation, and rain ensues.

149. As in the case of other meteorological phenomena, the circumstances connected with the production and retention of atmospheric vapour are extremely complicated: but in general terms it may be stated that heat is the grand promoter of evaporation; that a gentle current of air, by removing the pressure of the vapour as it is formed, is more favourable than a stagnant atmosphere; and that it is retarded by moist and cloudy conditions of the air, even though the sensation of heat be considerable. As previously mentioned, the humidity of the atmosphere is confined chiefly to its lower strata; and so rapidly does the quantity diminish as we ascend, that at the height of five or six miles it becomes all but inappreciable. The quantity of moisture evaporated from the surface of the globe may differ from day to day,

and from year to year ; but on the whole, and for any given number of years, it is returned again in the state of dew, rain, hail, or snow. Understanding the general conditions of the atmosphere as regards its heat and moisture, we may now proceed to its special effects in the production of winds, rains, and other kindred phenomena.

Winds—Constant, Periodical, and Variable.


150. As already explained, any current of the atmosphere produced by inequality of temperature or density is termed a *wind*—the denser and colder air rushing in to supply the place of the lighter and warmer. In other words, *wind is air in motion* ; and the most frequent cause of such motion is disparity of temperature between adjacent districts. As inequality of temperature is ever arising alike from general and from local causes, the occurrence of winds is incessant and universal, either along the earth's surface, or in the higher strata of the atmosphere. While the colder current is flowing *below* from the colder to the warmer region, a warm current is floating *above* from the warmer to the colder. As the homogeneity of the ocean is maintained by its currents, so the equilibrium and uniformity of the atmosphere is preserved by its winds. These winds, having different times, directions, and characters, are necessarily arranged into different classes, and distinguished by various designations. As regards times, those blowing constantly in one direction between, and within a few degrees beyond, the tropics (the Trades), are spoken of as *permanent winds* ; those blowing at certain periods (the Monsoons, &c.), as *periodical* ; and those obeying no fixed period, as *variable*. As concerns direction, they are distinguished by the points of the compass, as North, South, East, West, South-west, South-south-west, &c., according to the quarter from which they blow. And as regards inherent or climatic character, they are cold or hot, dry or moist, gentle or boisterous, healthful or unhealthful, according to locality and other accompanying conditions. One of the most obvious characteristics of winds, is their velocity, which may vary from a few miles to upwards of a hundred miles an hour ; and between these extreme rates we have every variety, from the gentlest zephyr to the most violent hurricane. According to meteorological authorities, a velocity of 7 miles per hour is regarded as a gentle air ; of 14, as a light breeze ; of 21, a good sailing breeze ; of 41, a gale ; of 61, a great storm ; of 82, a tempest ; and of 92, a hurricane, producing universal devastation—tearing up trees and sweeping away buildings.

151. In whatever character they occur, winds are important agents in the production and modification of climate. By their agency the moist and heated atmosphere of one region is transferred to another ; and on their agency also depend, in a great

measure, those currents of the ocean which are ever producing interchanges between the colder and warmer surface-waters of different latitudes. Besides these great climatological functions, they are intimately concerned in the production of rains and other aqueous phenomena ; while their incessant commotions tend to preserve the atmosphere in ever-healthy equilibrium. Geologically, winds have considerable effect in removing, piling up, and re-assorting all loose superficial matters, as the sand-dunes of the sea-shore and the sand-drift of the desert ; while through the agency of the ocean-waves, which are created by their power, important changes are produced along every shore of the world. In their gentler manifestations they assist in the fertilisation of plants and in the dispersion of their seeds ; while in their fiercer demonstrations their track is marked by ruin and devastation. By their impulse the commerce of distant nations is wafted from shore to shore ; and fickle and fitful as they may appear, man not unfrequently avails himself of their power to turn the wheels and shafts of his machinery.

152. The most remarkable of the *permanent, constant, or perennial* air-currents are the TRADE WINDS (so called from their influence on the trade or commerce of the world), which, within the torrid zone and a few degrees beyond it on either side, are ever sweeping round the globe in a westerly direction. This is occasioned by the fact that the region near the equator is most intensely heated—the other zones becoming colder and colder towards either pole. Under these circumstances, the air of the torrid zone becomes rarefied and ascends, while the colder and denser air sets in from either side to supply the deficiency. There are thus generated two great sets of currents—colder under-currents setting in towards the equator respectively from the north and south, and warm upper-currents flowing off from the equator towards either pole. If, then, the atmosphere were subject to no other influence than temperature, a *north wind* would always prevail along the earth's surface in the *northern hemisphere*, and a *south wind* in the *southern hemisphere*. But as the polar currents proceed towards the equator, they gradually come into zones having a greater velocity of rotation, and thus they are deflected in a westerly direction (as exhibited in the accompanying Sketch-Map), so as to become a *north-east wind* in the northern hemisphere, and a *south-east wind* in the southern.

153. In the Pacific, the north-east trade-wind may be said to prevail between the 25th and 2d degrees of N. latitude ; while the south-east trade ranges more widely between the 10th and 21st of S. latitude. In the Atlantic, on the other hand, the former is comprised between the 30th and 8th degree of N. latitude, and the latter, within the 3d of N. and the 28th of S. latitude. These limits, however, are not altogether stationary, but depend on the seasons—advancing towards the north during



the summer of the northern hemisphere, and receding to the south as the sun withdraws to the southern tropic. Within these somewhat fluctuating zones, the trade-winds are steady and perennial, travelling at the rate of from ten to twenty miles an hour, and wafting onwards in safety the merchandise of the globe. As they approach the continents, their courses become interrupted in consequence of the unequal heating of the land and water surfaces; hence within these coast-areas they assume the character of periodical, rather than of constant currents, being deflected at certain seasons from their normal north-east and south-east directions.

154. As the north-east and south-east trades approach the equator of temperature, their currents begin to fail, and this effect, augmented by the upward tendency of the highly heated air of that region, produces a zone or belt of calms, which fluctuates a few degrees north and south, according to the seasons. If the approaching trades be equally reduced in any locality, a dead calm will be the result; but as this is seldom the case, the zone is also characterised by short gusty winds, which vary in force and direction according as either main current prevails. This *zone of calms and variables*, as it is termed, is further characterised by the frequency and intensity of its thunderstorms—the crossing and collision of currents of unequal densities being favourable to the manifestation of electrical phenomena. In the Atlantic, the region of calms (see Sketch-Map) ranges in August between 3° and 12° N. Lat.; but in February it extends from $1^{\circ} 15'$ to 6° N. Lat. A portion of this zone near the Cape Verde Isles is known to sailors as “the rainy sea;” and is described as “a region doomed to continual calms, broken up only by terrific storms of thunder and lightning, accompanied by torrents of rain. A suffocating heat prevails, and the torpid atmosphere is disturbed at intervals by short and sudden gusts, of little extent and power, which blow from every quarter of the heavens in the space of an hour—each dying away ere it is succeeded by another. In these latitudes vessels are sometimes detained for weeks.” In the Pacific, the region of calms is comprised within the $2d$ degree of north and south latitude, near Cape Francis and the Galapagos Islands—a narrow belt separating the two trades, and characterised by the same phenomena as the calm zone of the Atlantic.

155. Besides the Trade-winds, which are the most persistent of all aerial currents, there are certain winds in the higher latitudes which also blow with considerable continuity. The heated air which ascends from the tropics, and flows off towards either pole, gradually descends as it proceeds; but as it passes from latitudes having a high to others having a less rotatory velocity, it is deflected from its original course, and assumes a south-easterly direction in the northern hemisphere, and a north-easterly in the southern. In this way the prevailing winds in

the higher latitudes of the northern hemisphere are from the *south-west*, and those in the southern hemisphere from the *north-west*; and though many causes—unequal distribution of sea and land, position of continents, irregularity of land-surface, &c.—tend to disturb this continuity, still, on the whole, there is a marked predominance of westerly winds. In our own islands, the fact must be sufficiently obvious to every observer; in the North Atlantic, the average packet voyage from New York to Liverpool is 23 days, while the return voyage requires 40 days; and in the vicinity of Cape Horn, westerly winds are three times more frequent than those from an easterly direction.

156. Among the constant currents of the atmosphere must also be enumerated the *north and south polar winds*, which, as mentioned in the theory of the Trade-winds, are continually flowing north and south from either pole. And further, as an under-current almost invariably implies the existence of an upper current in a contrary direction, we may notice also the *upper west wind of the tropics*, which, high above the Trades, seems constantly flowing in an opposite direction. Proofs of this upper tropical wind are found in the circumstance that dust ejected from the volcanoes of the West Indies has fallen on ship-deck several hundred miles to the eastward; and also in the often-observed fact, that a similar wind prevails near the summit of Teneriffe (12,000 ft.), while the regular trade-wind is blowing from an opposite direction below.

157. Of the *periodical winds* the most important are the *Monsoons* (from the Malay word *moussin*, signifying seasons), which in certain countries within and near the tropics blow from a certain quarter for one-half the year, and from an opposite point during the other half—the period of change being marked by calms, tempests, and variables. In other words, the monsoons are but the trade-winds interrupted in their regular action by the geographical peculiarities of the regions in which they occur. From April to October, the *south-west* monsoon prevails north of the equator, and the *south-east* in the southern hemisphere; but from October to April the *north-west* monsoon blows south of the equator, and the *north-east* in the northern hemisphere. Of course, the farther from the equator, the later in the season will the south-west and north-west monsoons occur; and thus it happens that in India, at Anjengo, on the Malabar coast, 8° 30' N. Lat., the south-west monsoon commences as early as the 8th of April; at Bombay, 10° N. Lat., about the 15th of May; in Arabia it commences a month earlier than on the coast of Africa; and in the northern part of Ceylon, fifteen or twenty days earlier than on the coast of Coromandel.

158. The cause of the monsoons is to be sought in the effect produced by the sun during his apparent annual progress from one tropic to the other, and has been thus explained:—"In the

Indian Ocean, for example, as the sun advances towards the north, the zone of greatest rarefaction recedes from the equator, and the north-east monsoon (which is nothing more than the trade-wind) then subsides, and is succeeded by calms and variable winds; but as the summer approaches, and the sun arrives at the northern tropic, the southern portions of the Asiatic continent become hotter than the ocean, and the humid air from the equatorial seas flows northward to the land; south-west winds will therefore arise, which prevail from the peninsula of India to the Arabian Gulf, until the time of the autumnal equinox. During the same period the south-east monsoon, in the southern hemisphere, tempers the heat of Lower Guinea, and brings rain to the shores of Brazil. The motions of the atmosphere, however, are reversed as the sun crosses the equator and approaches the southern tropic. Pouring his fervid rays upon Southern Africa, the vast tract of New Holland, and the splendid clime of Brazil, the air flows in from the north and north-west towards these highly heated regions, and winds from these quarters prevail for several months—the monsoon extending along the coast of Brazil from Cape St Augustine to the Isle of St Catherine. But now the influence of the sun is partially withdrawn from Southern Asia; it glows no longer beneath its vertical rays, and over the cooled earth the north-east monsoon resumes its wonted course."

159. Equally remarkable for their persistency, though local in their areas and limited in their times, are the *land and sea breezes*, which occur on almost every seaboard, but most notably, of course, within the tropics. As formerly explained, these breezes arise from the unequal heating of the land and water surfaces, and become most decided where this inequality is greatest. During day the land-surface, from its low conducting power, acquires a more elevated temperature than the adjacent waters, and the air above it, partaking of this heat, becomes rarefied and ascends, while the cooler and denser air from the ocean sets in as a *sea-breeze* to restore the equilibrium. This sea-breeze, especially in tropical latitudes, commences about nine in the morning, gradually increases in force till the middle of the day, and falls away as the sun declines in the afternoon. As evening approaches, the air over the surface of the land becomes more rapidly cooled by radiation than that over the water, and then a cool *land-breeze* sets out towards the ocean, blows freshly during the night, and dies away towards morning, when the sea-breeze again commences. The extent of these breezes is exceedingly variable. In some localities they blow only for a few miles out and in of the shore-line, while in others their bracing and refreshing influence is experienced for many leagues in either direction.

160. Owing to the mobility of the atmosphere, and the many

causes that may temporarily affect its temperature and density, the *variable winds* are exceedingly numerous and capricious—so capricious, that “fickle as the wind” has become an established simile in every extra-tropical country. We say *extra-tropical*, for within the tropics the Trades and Monsoons are the prevalent winds, and these, as we have seen, maintain a remarkable degree of permanency and periodicity. But in extra-tropical regions, where the force of the permanent winds becomes feeble, a thousand causes (as unequal distribution of land and water, nature of soil, irregularity of surface, &c.) occur to disturb these currents, and thus almost every district has its temporary winds—varying in direction, force, and duration. Fickle and uncertain as they may appear, they are, nevertheless, the results of law and law-directed forces; but these forces being so complicated, comparatively little has been determined in this department of meteorology. For this reason we shall merely advert to a few of the local winds that have become specially noticeable, either for their frequency or for their physical characteristics.

161. Generally speaking, in the higher latitudes of the northern hemisphere, a south wind is warm and moist, because it comes from warmer regions, and passes over a greater extent of ocean; while, on the contrary, and for opposite reasons, a north wind is cold and dry. It is for this reason that the westerly and south-westerly winds of Europe are humid and genial, and the north and north-easterly hard and ungenial. These major currents are of course much modified in their passage over certain localities, and hence the various characteristics assigned to the *Etesian Winds* (Gr. *etesios*, annual) that prevail very much in early summer all over Europe. One of the most noticeable of local winds is the *Simoom* (Arabic, hot and poisonous)—a hot suffocating blast that occurs in most countries bordering on sandy deserts, especially in certain parts of Asia and Africa, where its temperature has been noted as high as 120 and 130 degrees. Coming from the arid desert, and laden with the minutest particles, it often gives a reddish colour to the atmosphere, and thus forewarns the traveller to take shelter from the approach of its pestilential breath. In Turkey it is called the *Sameli*; in Egypt, *Khamsin* (fifty), because it usually continues fifty days; and in Guinea and Senegambia, *Harmattan*. The *Sirocco* (Arabic) is another hot, parching wind, that occasionally passes over Sicily and adjacent districts, and is supposed to originate in the Sahara, or great burning desert of Africa. The *Solano* (Lat. *sol*, the sun) is a similar south-west wind that occasionally visits the Spanish peninsula, and, blowing from the direction of the African deserts, is regarded as a modified sirocco.

162. In Buenos Ayres, *Pampero* is the name given to a violent west wind, which, traversing the arid plains of the Pampas, raises whirlwinds of dust, and carries them forward to the coast of the Atlantic. The *pamperos* seem to be portion of the return or

north-west trade-winds. In Peru, between the Cordilleras and the Andes, at the height of 12,000 feet, are vast tracts of desolate table-land, known by the name of the *Puna*. These regions are swept for four months in the year by piercing cold winds (the *Puna Winds*) from the snowy peaks of the Cordilleras, which are so extremely dry, and absorb moisture with such rapidity in their passage, as even to prevent putrefaction in dead animal bodies.

163. Among the most remarkable of the variable winds are those familiarly known as *Whirlwinds*, and technically as *Tornadoes*, *Typhoons*, and *Cyclones*. Properly speaking, a whirlwind is an aerial current that assumes a rotatory, whirling, or spiral motion. *Whirlwinds* are often of great violence, but fortunately of short duration, and are most frequently caused by the meeting of contrary winds, though sometimes by obstructions of the land, as precipitous mountains, narrow gorges, and the like. Their occurrence at sea produces *waterspouts*, and on the sands of the desert *sand-pillars*, and kindred phenomena. *Tornadoes* (Span. *tournar*, to turn) are merely violent whirlwinds, though the term is usually applied to such as are accompanied by lightning, thunder, and rain. *Typhoons* (Gr.) are literally also tempests or whirlwinds; but the name is specially given by navigators to the hurricanes that visit (generally from June to November) the seas of southern China and the adjacent archipelago of the Phillipine and Molucca islands. *Cyclone* (Gr. *kuklos*, a circle or whirl), on the other hand, is now the technical term applied by navigators to those rotatory hurricanes which occur most frequently between the equator and the tropics, and near the equatorial limit of the trade-winds. They sweep round and round with a progressive motion, their course describing a curve, and their violence being greater the narrower the limit of their whirl. In both hemispheres the rotation of a cyclone is *contrary* to that of the sun—those in the northern hemisphere moving counter, so to speak, to the motion of a watch hand; those in the southern following that motion.

Dew, Rain, Hail, Snow, &c.

164. Closely connected with the winds, and greatly influenced in their character by them, are the dews, rains, snows, and other aqueous phenomena which enter so largely into the determination of climate. The consideration of these subjects belongs especially to the province of Meteorology, but as much may be here recapitulated as will enable the learner to trace their connection with his immediate study of Physical Geography. As already stated, insensible vapour to a greater or less amount is always present in the atmosphere as one of its accidental ingredients. The warmer the air the greater its capacity for this moisture; hence the greater amount taken up during the heat of the day. After sunset the

earth and air lose their heat by radiation into space, but the former parts with it more rapidly; and hence on its cooled surface the moisture in the air is condensed as *dew*—just as the outside of a cold-water bottle is bedimmed or *bedewed* with moisture on being brought into the warmer air of a sitting-room. Dew is, therefore, the moisture insensibly deposited from the atmosphere on the surface of the earth; and what is termed the *dew-point* is that temperature of the atmosphere at which it begins to be precipitated. Dew never begins to be deposited upon the surface of any body until it is colder than the contiguous atmosphere; and the quantity deposited depends chiefly upon the humidity, serenity, and tranquillity of the atmosphere, as well as on the constitution, surface, and locality of the bodies receiving the moisture. Still, clear nights are, therefore, more favourable for the deposition of dew than windy and cloudy ones; and all substances like glass, silk, wool, grass, &c., which rapidly lose their own heat and slowly acquire that of others, are susceptible of being copiously bedewed; while substances like rocks and metals, which possess opposite qualities, contract but little dew. In temperate zones, where the frequent interchange of sun and shower preserves the earth from the extremes of heat and moisture, very little dew is deposited; but in tropical regions, where the day-heat is excessive, and no rain falls for months, the dews are most abundant and refreshing. But for this beneficent arrangement many intertropical countries would be altogether sterile and barren of vegetation.

165. When the temperature of the air is reduced below that of the invisible vapour it contains, the moisture becomes visible, and appears as *fogs*, *mists*, and other kindred phenomena. These fogs occur most abundantly along the courses of rivers, on mountain sides, over shoals and banks at sea, and generally along the seaboard of continents and islands; and this obviously in consequence of the unequal temperature of the contiguous lands and waters. Like dews and rains, fogs exercise a refreshing influence on vegetation, and in some countries, like Peru, their occurrence is periodical and regular. *Clouds* have been defined as “masses of visible aqueous vapour which float in the sky, or drift through it with the wind, assuming every variety of colour and form.” As clouds they exercise but an indirect effect on terrestrial climate, and it is only when they break, by further condensation, into rain, snow, and hail, that they come directly within the province of the geographer.

166. The precipitation of water from the atmosphere in the form of *rain* depends chiefly on the further condensation of clouds and fogs by the commingling of colder and warmer currents. The capacity of air for moisture decreases at a faster rate than the temperature, and thus the mingling of two currents only slightly differing in temperature may so reduce this capacity, that heavy rainfalls will take place even where there is no great decrease in atmospheric warmth. Where the temperature of the

air falls below the freezing point (32° Fahrenheit), the atmospheric vapour may be converted into *snow*; or if rain has been already formed in the upper air, the drops in passing through strata beneath the freezing point will be converted into *hail*. Rain, hail, and snow are thus to a certain extent convertible phenomena: hail occurring in all latitudes and climates; but snow only in the colder latitudes, and at elevations where the thermometer continues below 32°. Winds being the great natural agents by which the colder and warmer masses of the atmosphere are brought into combination, rains will occur most frequently where these currents are shifting and variable. Where the winds are constant and of equable temperature, rain seldom happens, unless at points where these currents impinge on lofty mountains, and carry up the air of the sea and plains into the colder regions of the atmosphere. As the capacity of the air for moisture increases with the temperature, it must happen (other things being favourable) that the amount of rainfall is greater in warm than in cold latitudes, and greater also in low-lying than in elevated districts.

167. For the foregoing reasons the amount of rainfall is greatest within the tropics, and decreases as we advance north or south into higher latitudes. It is also greater at the sea-level and moderate elevations than it is on lofty table-lands and mountains. In like manner more rain descends upon the coasts than upon the central regions of a country—the humid air from the ocean gradually parting with its moisture as it is borne farther and farther inland. In Britain, for example, where the prevalent winds are from the Atlantic, the annual rainfall on the west coast is about 37 inches, while on the east side of the island it rarely exceeds 25 inches. It must be observed, however, that though the annual rainfall within the tropics is greater than in higher latitudes, the number of rainy days are fewest within the tropics, or where the winds are constant, and most numerous in the higher latitudes, where the winds are shifting and variable. Thus, between 12° and 43° N. Lat., the number of rainy days is stated at 78; between 43° and 46°, 103; between 46° and 50°, 134; and between 50° and 60°, as many as 161. Of course, since the rainy days are fewer in low latitudes, and the annual amount of rainfall greater, the rains must be much more powerful—a fact sufficiently recorded by all travellers in tropical regions.

168. Understanding these principles, and bearing in mind what was stated regarding the constant, periodical, and variable winds, the learner will readily perceive—1. Why heavy and frequent rainfalls should take place within the so-called zone of calms, where evaporation is excessive, and the air often gusty and variable; 2. Why rain should seldom fall at sea within the region of the steady and equable trade-winds; 3. Why the rains within the range of the monsoons should be periodical, and the year divided into two seasons, a wet and a dry; 4. Why the rains in

the higher latitudes should occur at no fixed period, but be irregularly distributed throughout the year ; 5. Why some regions, like the central deserts of Africa and Asia and the coasts of Peru, should be altogether rainless ; and, 6. Why almost constant rains should occur in some countries, like Guiana ; while in others showers of excessive violence should occasionally fall, adding to their annual rainfall as much in the space of a few hours as ordinarily happen during the course of many months. As may be expected, the annual rainfall of these different regions will differ very widely ; and will be attended, of course, by proportionate results—heavy falls being beneficial in hot countries, and moderate supplies in temperate and milder zones. In the British Islands the annual fall (as ascertained by rain-gauge) ranges from 24 to 60 inches, or has an average about 36 inches ; while in tropical countries the mean is upwards of 200 inches. As much, however, as 229 inches has been noted in Dutch Guiana, 276 in Brazil, 302 at an elevation of 4200 feet in the Western Ghauts, south of Bombay ; and in the Khasia Mountains, at the head of the river-flats or Jheels of Bengal, upwards of 600 inches or 50 feet, have been registered by several observers. At the same place, Dr Hooker has recorded 30 inches in twenty-four hours ; 21 inches have been noted at Cayenne during the same period ; and 23 inches are not uncommon near Port Jackson in New South Wales.

169. Extending the facts alluded to in the preceding paragraphs, the rainfall of the globe may be arranged under three great heads—the periodical of the tropics ; the variable of the higher latitudes ; and the abnormal of certain districts where it occurs either in excess or is altogether absent. Within the tropics the rainy season commences at the shifting of the monsoons ; and as this change is dependent upon the position of the sun, it begins earlier in those regions that lie near the equator than in those more remote. In general terms, the rainy season in the northern half of the torrid zone may be said to commence with April and last till October, while the dry season extends from October till April. In the southern half this order is reversed—the dry season embracing from April till October, and the season of rain from October till April. In Africa, for instance, near the equator, the wet season begins in April and continues till October ; while in Senegambia it does not commence till June, and then lasts till November. In India, on the Malabar coast, the rains commence early in May, but do not reach Delhi till near the end of June. In the New World, also, the rain falls at Panama early in March, but it seldom appears in California before the middle of June. Of course, within the tropics as elsewhere, the regularity of the periodical rains is interrupted by the configuration of the land, mountain chains, and similar causes ; and hence the peculiarities that mark the times as well as intensities of the wet and dry seasons of such areas as the east coast of Africa, the

Red Sea, the Malabar coast, the Coromandel coast, and the coasts of Australia. On the Malabar coast, for example, the south-west monsoon (coming from the humid region of the equatorial ocean to supply the place of the highly rarefied air of the heated continent) is said to be "ushered in by terrific storms of thunder and lightning. The water pours down in torrents, and when the thunder has ceased, nothing is heard for several days but the rush of the descending rain, and the roar of the swelling streams. In a few days the storm ceases, and the earth, which before was withered by the glowing atmosphere of the dry season, is now, as if by magic, suddenly clothed with the richest verdure; the air above floats pure and balmy, and bright tropical clouds sail tranquilly through the sky. After this, the rains fall at intervals for the space of a month, when they again return with great violence. In July they attain their height, and from that time gradually subside until the end of September, when the season closes, as it began, in thunders and tempests." On the eastern, or Coromandel side, the order of things is reversed. The south-west monsoon, in passing over the western Ghauts and the central table-land of the Deccan, parts with all its moisture, and reaches the eastern side as a hot, dry wind; and it is not till the north-east monsoon begins to prevail in October that the rainy season is experienced on the coast of Coromandel.


170. Beyond the tropics the rains no longer occur at stated periods, but become *variable*—that is, are distributed throughout the year in a very irregular and uncertain manner. In some countries they occur most frequently in winter; in some, in spring and autumn; and in others, again, most abundantly in summer. Thus, the amount that falls in the west of England in winter is said to be eight times greater than that in summer; in Germany the winter and summer amounts are about equal; but in St Petersburg the winter fall is little more than a third of what descends in summer. Again, in Britain, there are more rainy days in winter than in summer; but in Siberia it rains four times as often in summer as in winter. The countries of Europe bordering the Mediterranean are generally regions of winter rains; while those of Western Europe are distinguished rather by the abundance of their autumnal rains. In Europe north of the Alps, the north-east winds (as coming from the higher and colder latitudes) are comparatively dry; while those from the south-west (from the humid expanse of the Atlantic) are warm and laden with moisture. On the eastern coast of North America the reverse holds good, and the north-east winds borne from the ocean give rise to the long-continued rain-storms of spring and autumn.

171. As might be expected, the districts of *excessive rainfall* lie within the tropics, and mainly in the immediate neighbourhood of the equator. In the New World, Brazil, Guiana, the West India Islands, Central America, and the shores of the Mexican Gulf, are all notable for their heavy and continuous rains;

while in the Old World, the coasts of Guinea and Senegambia, Eastern Africa, India, and the Indian Archipelago, are remarkable for similar phenomena. (See Rain Map in Atlas.) As some regions are celebrated for their rainy character, so others are equally remarkable for the entire absence of rainfall. Of these *rainless tracts* the more noted are the great desert lands of Africa, Arabia, Persia, and Mongolia—an almost continuous area, varying in breadth from the 15th to the 47th parallel, and in length from 16° west to 118° east longitude. Except on its borders this vast expanse is all but absolutely rainless, its heated atmosphere readily absorbing and converting into invisible vapour whatever humidity may be carried towards it from adjacent regions.

172. Of the other aqueous phenomena more immediately bearing on climate, the limits of an introductory sketch will only permit the briefest allusion to hoar-frost, snow, hail, and those accumulations of frozen water known as glaciers and icebergs. *Hoar-frost* is produced in the same manner as dew, and occurs chiefly in early spring and autumn, during serene nights, and when the surface of the earth falls below the freezing point. Whatever prevents the rapid radiation of heat (overhanging foliage, passing clouds and currents), arrests the formation of hoar-frost, which is often very destructive to tender plants and to buds and blossoms in early spring. *Snow* is the frozen moisture that descends from the atmosphere when the temperature of the air at the surface of the earth is near or below the freezing point of water. Of course, at the sea-level within the tropics, and for fifteen or twenty degrees beyond them in either hemisphere, snow is unknown (see Rain and Snow Map in Atlas); and it is only during winter that it falls in the higher latitudes, and at considerable elevations. In the polar regions, and at extreme heights in all latitudes, it becomes perennial; and this limit at which it remains unaffected by the heat of summer is known as the *snow-line*, or less accurately, perhaps, as the line of *perpetual congelation*. The lower the latitude—that is, nearer the equator—the higher the snow-line, which descends constantly, but somewhat irregularly, according to the nature of the situation, as we proceed towards either pole. The following diagram will convey some idea of its gradual descent from 16,000 feet at the equator down to the sea-level at the poles. From the equator to latitude 30° there is little variation in the height of the snow-line, unless there be some peculiarity as regards winds and moisture in the situation; but from 30° to 60° the descent is very rapid, and the limit rapidly approaches the sea-level, as was formerly noted in Chap. VI., on Mountains and Mountain-Systems. As hoar-frost may be said to be frozen dew, and snow frozen mist, so *hail* may be described as frozen rain. It occurs in all latitudes and at all seasons; and though the producing causes are not always discernible, it seems mainly dependent on the meeting of aerial

currents of greatly unequal temperatures, as well as on peculiar electrical conditions of the atmosphere. Falling in pellets from



the size of a pea to that of a pigeon's egg, hail-storms are frequently very destructive, but luckily are always restricted to limited areas.

173. As snow must accumulate on all mountain-ridges above the snow-line, as well as on the surface of all polar lands within the limits of perpetually frozen ground, it will become more or less compressed, and this compression will be greatly augmented by its partial meltings in summer, as also by the falling and freezing of rain and other atmospheric vapours. From this accumulation, compression, and re-freezing, arise avalanches, glaciers, icebergs, and other kindred phenomena. *Avalanches* (Fr.) are accumulations of snow, or of snow and ice, which are frequently precipitated with destructive violence from lofty mountain-ridges, like those of the Alps, into the valleys and plains below. They originate on the higher slopes, and begin to descend when the weight of their mass becomes too great for the declivity on which it rests, or when fresh weather destroys its adhesion to the surface. *Glaciers*, on the other hand, are accumulations of ice, or of ice and snow, which collect in the valleys and ravines of snowy mountains like the Alps and Himalayas, or on boreal uplands like those of Greenland, and which move downwards (partly by their own gravity and partly by the expansion of the freezing water that falls into their crevices and fissures) with a peculiar creeping motion, smoothing the rocks over which they pass, and leaving mounds of debris (*moraines*), lateral and terminal, as they melt away. In mountain-gorges, glaciers descend as ice-streams till, coming below the snow-line, they begin to melt and disappear; but on polar lands, where they cover almost the entire surface, they move downwards to the sea-shore, and there, losing their support, the advancing fronts break off and are floated away as *icebergs* (ice-mountains) and ice-floes (ice-islands). Such ice-masses have been met in the Arctic and Antarctic Oceans several miles in circumference, rising from 40 to

200 feet above the water, and loaded with stones and shingle. Some idea of their size may be formed from the fact that little more than a tenth of their bulk rises above the water, the specific gravity of ice being only 0.9. As they are floated by the polar currents to warmer latitudes they melt away, dropping their burden of boulders and debris on the bottom of the ocean. "Fields" and "packs" of such ice are familiar phenomena both in arctic and antarctic waters—the bergs of the north being seldom carried southward in the Atlantic beyond the 44th parallel of latitude, while those of the south are not unfrequently found northward as far as the Cape of Good Hope, or, on an average, at 10 degrees lower latitude.

174. As climatological agents, snow and ice have considerable influence on the regions in which they occur. Accumulating on mountain heights, they are the perennial sources of the springs and streams that descend to refresh and fertilise the thirsty lowlands; while in tropical countries the winds that pass over them are cooled, and flow downwards to temper the sultry atmosphere of the plains below. Masses of floating ice are productive of similar results, the seas and coasts in their vicinity being cooled by their presence, and fogs and rain-storms generated as they melt away. In the higher latitudes, on the other hand, where the winters are severe, snow forms a warm covering for the soil (the *snow-blanket*, as it is termed by farmers), and greatly defends vegetation from the rigour of the frost. Within the polar circle, also, the darkness of the long winter is considerably diminished by the *snow-sheen* or *snow-blink*—snow reflecting, instead of absorbing like the bare ground, the faint light that there proceeds from the sky. The grinding, grooving, and smoothing effects of glaciers on rock-surfaces, and the rock-transporting powers of icebergs, are facts now well known to Geology; and though their consideration belongs more immediately to that science, the results now and in time past cannot be altogether overlooked by the student of Physical Geography.

Causes affecting Climate.

175. In its technical sense, the term *climate* (Gr. *klima*, an inclination) denotes an imaginary belt of the globe parallel to the equator, and so called by the earlier geographers, because the differences of these climes depend on the inclination or obliquity of the sphere. Between the equator and the polar circle there were twenty-four such climatic belts—each depending on the increasing length of the day (half an hour's increase for each belt), and of course expanding in breadth from the equator towards the poles. From the polar circles to the poles there were six climes—the differences of the longest day in their case being counted by monthly periods instead of half-hours. The term is

now employed in a much wider sense, as embracing the entire *weather-conditions* of any district, and as such we shall now proceed to consider its relations. Although the climate of a locality is mainly dependent on its heat and moisture, yet so many circumstances tend to disturb and modify these conditions, that the subject becomes extremely complicated and difficult of determination. Latitude, height above the sea, distance from the ocean, nature of soil, distribution of land and water, direction of mountain-chains, winds and rains, currents of the ocean, cultivation, and the like, all exert their influence in modifying the weather-conditions or climate of any special locality. Our limits, however, will only permit a brief allusion to the more influential of these modifying causes.

176. As repeatedly stated, the main element in climate is the amount of heat received by the sun, and this diminishes (par. 144) according to the latitude or distance from the equator. Owing, however, to the obliquity of the earth's orbit, the regularity of this decrease is alternately interrupted in either hemisphere, and the respective length of day and night at opposite seasons becomes an important element in climatic diversity. Within the tropics day and night are all but equal—the longest day being little more than thirteen hours, and the shortest nearly eleven. In the latitude of Greenwich the longest day is nearly seventeen hours, and the shortest only seven; while, within the polar circles, there is a brief summer period when the sun never sets, and a corresponding winter season when he never rises. Bearing these facts in mind, it will be readily understood why the year, within the tropics, is divided into two seasons—a wet and a dry. The latter is regarded as the summer, and the former as the winter, but they are in direct opposition to the astronomical seasons, as the rains (par. 169) follow the sun. In some inter-tropical countries, as the West Indies, there are two rainy seasons and two dry seasons within the year. In the temperate zone, again, the year is divided into four seasons—spring, summer, autumn, and winter; but this regular succession of climatic change can hardly be considered as extending farther than from the 35th to the 60th parallels. In the frigid zones, on the other hand, only two seasons are known—a long and severe winter, during a portion of which the sun never rises above the horizon, and a brief but fervid summer when that luminary never sets. From these circumstances it will be at once perceived that the torrid zone is the region of greatest heat throughout the year, with comparatively little difference between its seasons; that the temperate zones stand next in order as regards the annual amount received, but experience great difference of temperature during the successive seasons; and that the frigid zones are regions of small annual heat, and at the same time subjected to seasonal extremes. Such is the general climatic order as regards latitude; though, owing to the smaller extent of land, the decrease of heat


as we depart from the equator is more rapid in the southern than in the northern hemisphere—so that, on a general average, the latter is about $3\frac{1}{2}^{\circ}$ warmer than the former.

177. Altitude is the next great modifier of climate, but owing to many correlative circumstances (prevalent winds, slope, proximity to the sea, &c.), its operation is not altogether uniform. As formerly stated, a decrease of 1° Fahr. takes place in the lower regions of the atmosphere for every 300 or 350 feet of ascent; but at great heights and in extreme latitudes the decrease is more rapid. As it is, under every latitude the loftier mountains (Himalayas, Alps, Andes, &c.) are perpetually covered with snow; the higher plateaux (Mexico, Bolivia, Armenia, Thibet, &c.) are several degrees colder than the contiguous lowlands; and in temperate zones such table-lands experience much greater extremes of summer's heat and winter's cold than the surrounding districts—having, like Spain, Armenia, and Persia, summers of tropical heat and winters of almost polar severity. Another great cause of modification is the unequal reception and radiation of heat by land and water (par. 146), by which islands and sea-coasts are rendered cooler in summer and warmer in winter than inland tracts—creating what has been termed *insular and continental climates*. Britain, New Zealand, and Tasmania enjoy, in this respect, insular climates; Germany, Central Russia, and Tartary, continental ones. Connected with this, and depending on the set of the trade-winds and ocean-currents, may be noticed the observation of Humboldt, that the continents and larger islands in the northern hemisphere are warmer on their western than on their eastern sides; while in the southern hemisphere the reverse holds good—the western being the colder and the eastern the warmer.

178. It would exceed the limits of an introductory outline to describe in detail the many causes concerned in the modification of climates; but besides those of latitude, altitude, and proximity to the sea, may be noticed the following:—1. The direction of mountain-chains, which, by intercepting cold winds, renders the countries on one side warmer than those of the other; and in like manner, by intercepting moist winds, favours the production of rain on the windward slopes and droughts on the leeward or sheltered declivities. The eastern slopes of the Andes, for example, intercept the humid trade-winds from the Atlantic; their counter-slopes on the Pacific side are arid and rainless. 2. The general inclination or slope of a district, as this may lie to the heat of the morning and noonday sun, or be turned to the feebler rays of his afternoon and evening declension. 3. The prevailing winds of a region, as these may be cold or warm, dry or humid. The westerly winds of our own island are humid and warm—the easterly are cold and dry; hence the greater rainfall and mildness of our western coasts as compared with the eastern. 4. As with winds, so, in like manner, with the influence of oceanic

currents. The Gulf Stream, by bringing warmth and moisture, mitigates the winter climate of Western Europe; the Arctic current, on the other hand, tempers the summer climate of the eastern shores of North America. 5. The direction of river-plains and valleys, as they open out to favourable winds and ocean-currents. The westerly reception of the Rhine basin produces a finer climate than the easterly trend of the Danube, even though the latter be several degrees farther south; and the southerly slope of the valley of the Rhone enjoys amenities unknown in the westwardly-trending basins of the Loire and Garonne. 6. Cultivation has also a marked effect on the climate of a country—the felling of forests, draining of lakes and morasses, and the like, being all favourable to greater dryness, warmth, and general amenity.

179. Under these circumstances, it will readily be perceived that the belts of climate can by no means correspond with the parallels of latitude, but that it requires a long series of observations to determine the *seasonal temperatures* of summer and winter, and also a careful average of these to ascertain the *mean yearly temperature* of any given locality. If the surface of the globe had been all water, or all land of equal altitude, the parallels of latitude would have determined the lines of climate; but this not being the case, the daily temperature, the summer and winter temperatures, as well as the mean annual temperature of any two places in the same latitude, may differ very considerably. In this way the maximum summer heat of an island may be several degrees below that of a continental country between the same parallels, while its winter temperature is many degrees higher. In this way, also, the mean annual temperature of some island or sea-coast may be equal to that of some inland district situated several degrees nearer the equator. A series of lines drawn through places having the same summer temperature show these variations at a glance, and are termed *isotheral lines* (Gr. *isos*, equal, and *theros*, summer), or lines of equal summer heat; those through places having the same winter temperature *isocheimōnal lines* (Gr. *isos*, and *cheimon*, *cheimonos*, winter); and those connecting places of the same mean annual temperature, *isothermal lines* (Gr. *isos*, and *thermè*, heat). The difference between the summer and winter temperatures may amount to 2°, 20°, 40°, or more degrees of Fahrenheit; but the isothermal lines show the mean amount of heat received throughout the year, and, of course, are the more correct indicators of the general climatic conditions of any given locality. In the accompanying Sketch-Map the isotherms are laid down for every ten degrees, and their bendings northward and southward (according to distribution of land and water, altitude, distance from sea, &c.) convey to the eye instructive proofs of the operating causes adverted to in the foregoing paragraphs.



RECAPITULATION.

In the preceding chapter attention has been directed to the Climatology of the globe—that is, to those weather-conditions on which its vegetable and animal life are so intimately dependent. The main medium of climate being the Atmosphere, it was necessary to advert to its nature and composition as an integral portion of our planet. As an aërial fluid, it consists of an admixture of 79 parts nitrogen and 21 oxygen, together with a small but variable proportion of carbonic acid gas, traces of ammonia, &c., and always holding in suspension less or more of aqueous vapour in a visible or invisible form. Constituted as plants and animals are, this composition is indispensable to their existence—the former assimilating carbonic acid and exhaling oxygen, the latter, in counterbalance, exhaling carbonic acid gas and consuming oxygen. Light and invisible as this aërial envelope may appear, it exerts a pressure on the earth's surface, at the sea-level, of about 15 lb. avoirdupois to the square inch—a pressure which is balanced by a column of mercury 30 inches in height; hence the *barometric column* of the meteorologist, and the *rise and fall of the barometer* according to changes in the weight of the atmosphere. As an elastic or compressible medium, its lower strata are denser or heavier than those at great elevations; and as its capacity for heat and moisture decreases with its rarity or attenuation, the higher regions of the air are colder and drier than those at a lower elevation. As the medium through which the light and heat of the sun are conveyed to the terraqueous surface, the atmosphere—partly owing to the varying inclination of the sun's rays, and partly to the unequal reception and radiation of heat by land and water—becomes variously heated in its different regions, and hence arise winds or aërial currents,—the warm air of one locality expanding and ascending, and the colder rushing in from all sides to supply the deficiency. The winds thus generated assume various directions and physical characteristics—the chief cause of modification being the different amounts of heat received by the different zones of the earth. Thus some are said to be *constant*, as the trade-winds of the tropics; others *periodical*, as the monsoons and the sea and land breezes; and others, again, *variable* and irregular, as the winds of the higher latitudes. In their physical characters they are governed chiefly by locality and the nature of the region from which they blow,

and are thus hot or cold, moist or dry, relaxing or invigorating—floating as a zephyr that scarcely disturbs the thistle-down, or sweeping as hurricanes that uproot forests and overturn buildings. The warmer the air, the greater its capacity for moisture ; but in every case there is a point beyond which it is incapable of sustaining more vapour in an invisible form. It is then said to be *saturated*, and in this state any cooling, by coming in contact with the colder earth, or by the contact and commingling of colder aerial currents, produces condensation into dews, fogs, rains, snow, hail, and other kindred phenomena. These aqueous phenomena,—whether descending as *dew* from the unequal temperatures of the earth and air—as *fogs* and *mists* from unequal temperatures of aerial strata—or as *rains*, by more sudden condensation,—are all essential to the vegetable and animal economies. Rains, like the winds on which they mainly depend, are almost constant within certain equatorial districts ; periodical within the regions of the monsoons ; and irregular in all the higher latitudes. Some tracts—as the Sahara, Egypt, Arabia, Persia, Mongolia, and Peru—are rainless, or all but rainless ; but in most countries the rainfall, though varying from month to month and year to year, is, in the long average, pretty regular and persistent. It will vary, of course, according to latitude, direction of prevailing winds, proximity to the ocean, direction of mountain-chains, and the like ; hence its great difference, even on opposite sides of the same island,—annual falls within temperate zones varying from 10 to 80 inches, and within tropical, from 100 to 500 or 600 inches. When the temperature of the air falls below the freezing-point, fogs and mists are converted into snow, and rain into hail—*snow* being mainly an extra-tropical and winter precipitation, *hail* occurring in all latitudes and at all seasons of the year. Within the tropics at the level of the sea snow is unknown ; in temperate zones it falls less or more during winter ; but in polar regions, and at great elevations above the sea, its presence is perennial in all latitudes. The altitude at which snow remains unaffected by the heat of summer (the *snow-line*) varies with the latitude—descending from 14,000 or 16,000 feet at the equator to near the sea-level at either pole. The latitudinal limit, north and south (the *snow-limit*), varies also according to the distribution of sea and land—receding and advancing as either element prevails. The perpetual presence of snow and ice on lofty mountain-ridges and polar uplands gives rise to *avalanches*, *glaciers*, and *icebergs*—phenomena whose geological influences are not less apparent than their climatological. The causes which

affect the *climate* or weather-conditions of any locality are thus extremely varied — latitude, altitude, distribution of land and water, proximity to the sea, prevalent winds, direction of mountain-chains, slope, nature of soil, cultivation, and the like, all more or less exerting their modifying influences. Heat and moisture, however, are the great regulators of climate; and thus, in general terms, climatological zones may be said to decrease in importance from the equator to the poles. Nevertheless, this decrease by no means coincides with the parallels of latitude; hence to determine the mean temperature of any locality, numerous thermometrical observations have to be made at hourly, daily, monthly, and yearly intervals. The results of such observations enable the meteorologist to connect places having the same summer, winter, and mean annual temperatures; and hence the scheme of exhibiting them at a glance by means of *isothermal*, *isoeheimonal* or *isoeheimal*, *isothermal*, and other lines.

XI.

PLANTS AND ANIMALS—THEIR DISTRIBUTION.

Life as affected by External Conditions.

180. HAVING noticed the general relations of the Land, Water, and Atmosphere, and the principal phenomena arising therefrom, we now turn to the Life by which they are respectively peopled. Hitherto we have dealt with the *inorganic* phases of nature; we have now to consider the *organic*. In the former case, the study involved consideration merely of *chemical and mechanical forces* acting from without; now we have to deal with the superaddition of *vital action* operating through peculiar organs from within. Under the term *Life* is embraced all that appertains to the vegetable and animal kingdoms—subjects which belong to the domain of Botany and Zoology, and only come under the notice of Geography in as far as they are dependent on external conditions for their position or distribution on the Globe. The origin, nature, and function of Life form the theme of Biology; its distribution and external relations become important considerations in the study of Physical Geography.

181. Whatever be the nature and origin of Life, it is clearly dependent for its continuance on the physical conditions by which it is surrounded. A little more heat or a little more cold, a little more moisture or a little more drought, and the plant flourishes or decays, the animal increases or dies. It is obvious, then, that on a globe having different zones of climate, having regions of excessive humidity and regions of excessive drought—and having, moreover, different areas of land and water—that life must be as diversified in its nature as the conditions under which it is destined to exist. The arrangement of plants into aquatic, terrestrial, and aerial—into flowering and flowerless—into trees, shrubs, herbs, grasses, mosses, lichens, sea-weeds, &c.—belongs to the province of Botany; and of animals into aquatic, terrestrial, and amphibious—into mammals, birds, reptiles, fishes, shell-fish, &c.—belongs in like manner to the province of Zoology. What the geographer has more especially to consider is their distribution over certain areas, the physical conditions appa-

rently concerned in their restriction to these areas, and the dependence of the one kingdom upon the other, as completing the economy of nature. For the sake of brevity, the term *flora* (Lat.) is employed to designate the plant-life of a region or epoch, and the term *fauna* its animal-life; hence we speak of the "Flora of South America" and the "Fauna of South America," as well as of the Flora and Fauna of the Tertiary, the Chalk, or any other geological period.

182. So far as the eye, or the eye aided by the microscope, can perceive, life is everywhere present—in the air, on the earth, and in the water, or even parasitic on and within other plants and animals. Unless, perhaps, among the perpetual snows and ices of the poles and lofty mountain-peaks, or in the extreme depths of the ocean, its manifestations are sufficiently apparent, and even in these situations some forms unknown to observation may find a permanent or temporary home. And yet, universal as life may appear, it is confined to the merest film of the terraqueous globe. A few thousand feet above the sea-level on land, and a few thousand feet beneath it in the waters, limit this *stratum of life* on either hand. Thickest at the equator, it thins out towards the poles; and densest near the sea-level, it becomes rarer and rarer the more it rises above or falls beneath this line of greatest intensity. Heat, light, moisture, and nature of soil are the great regulators of life on the land; heat, light, depth, nature of bottom, and saline composition, the main regulators in the ocean. Were it not for these causes there is no reason why the same forms of life should not prevail in every region from the equator to the poles, and from the shore-line to the darkest abysses of the ocean. The influence, however, of external conditions is insuperable. The palm of the tropics would dwarf and die in the temperate zone; the whale of the Greenland seas would perish in the waters of the equator; the rush that luxuriates in the marsh would wither if transferred to the arid upland; the shell-fish that swarm within the influence of the tides would die if submerged to the depth of a few hundred fathoms.

183. Every plant and every animal is, therefore, adapted by nature to the position it occupies, and within that position continues to fulfil its function so long as the surrounding conditions remain unchanged. Why one genus or kind should differ from another genus man may never know; and, at all events, the inquiry belongs to the general subject of biology and not to that of geography. Again, why certain forms should only appear in certain regions—the kangaroo, for instance, in Australia, the ostrich in Africa, and the llama in South America—while other regions seem equally fitted for their residence, is a question involving considerations of origin, and of geological alternations of land and water, that lie beyond the scope of our subject. What more immediately concerns geography, are the existing distribution of plants and animals, the conditions accom-

panying that distribution, and the question how far they are capable of being transferred to, and acclimatised in, other districts for the luxury and necessities of man. Some tribes have naturally a wider range than others; some, again, have a more elastic constitution, and are capable of enduring greater diversities of climate; and others, obeying the instincts of food, procreation, &c., migrate from the unfavourable season of one region to the favourable season of another. Of all animals man has the widest range—his superior intelligence enabling him to modify and overcome conditions that would be fatal to other creatures. Many, however, of the domesticated animals and cultivated plants have also considerable elasticity of constitution, and thus man has been enabled to carry them along with him over the greater part of the habitable globe. In this way we require to distinguish between the truly *indigenous* or native products of a country, and the *exotic* or imported, though many species have been so long transferred and re-transferred that it is now impossible to point to their original habitats. While, therefore, there is a natural apportioning of plants and animals to certain areas, and while external conditions are evidently the main regulators of this distribution, it must ever be borne in mind that man has already modified, and is continually modifying, this distribution by transferring, cultivating, and destroying, according as his wants and wishes may compel. His operations, however, can never extend beyond a mere modification; for over and above his control remain the great conditions of heat, light, moisture, &c., which ever govern the main geographical arrangements of plants and animals; and it is to these in particular that the limits of an introductory sketch will permit us to refer.

Plant Distribution.

184. As already mentioned, heat, light, and moisture are the principal conditions affecting the terrestrial distribution of vegetable life. These conditions have their greatest intensity near the equator; hence the greatest exuberance of vegetation within the tropics, and its gradual declension as we proceed towards either pole. As might be anticipated, however, this declension is governed more by the isotherms than by the parallels of latitude—the mean amount of annual heat being the predominant condition in vegetable distribution, though the amount of summer heat (isothermal) has also much to do with its ripening and perfection. The mean annual temperature of two places may be the same, and yet the summer temperature of the one may be 10 degrees higher than that of the other, while the winter temperature may be 15 or even 20 degrees lower. Much, therefore, in Botanical Geography, depends upon the amount of heat which a plant receives during the period of its greatest activity,

and this, in general, is regulated more by the monthly than by the annual isotherms. The zones of vegetation shading more gradually into each other than the astronomical zones (torrid, temperate, and frigid), botanists make a minuter subdivision of the earth's surface into equatorial, tropical, sub-tropical, warmer-temperate, colder-temperate, sub-arctic, arctic, and polar—each characterised by some peculiar feature, though partaking on either side of the forms that belong to the two adjacent zones.


185. 1. The *equatorial* zone, bounded by the isotherm of 79° , is characterised, wherever moisture is present, by its luxuriant vegetation. Succulent stems, large and showy flowers, gigantic parasites and climbers, arborescent grasses (canes, bamboos, &c.), orchids, palms, bananas, and the like, are typical of this belt, the greater portion of which, for reasons given in the preceding chapter, lies to the north of the equator. 2. The *tropical* zones, extending to the isotherm of $73\frac{1}{2}^{\circ}$ on either side of the equator, are marked by palms, bananas, pine-apples, tree-ferns, species of fig, pepper-shrubs, cotton, coffee, sugar-cane, &c.—there being fewer parasites and more underwood in the forests. 3. The *sub-tropical*, bounded by the isotherm of 63° , are characterised by a luxuriant growth of magnolias, laurels, myrtles, and figs, together with certain palms, zamias, cactuses, and arborescent euphorbias. In these zones, as in the damp regions of the preceding belts, vegetation is green throughout the year, and the climate, unless where rainless, is described as delightful. 4. The *warmer-temperate*, limited by $53\frac{1}{2}^{\circ}$, are still regions of evergreens, but are marked by the absence of palms—the dwarf palm of Europe, the palmetto of North America, and the Chilian palm, being, as it were, outlying forms from the sub-tropical zone. Deciduous forest-trees, oaks, chestnuts, &c., and figs, oranges, pomegranates, olives, and the vine, are typical of the warmer-temperate in the northern hemisphere; shrubby ferns, arborescent grasses, and araucariæ in the southern. 5. The *colder-temperate*, bounded by the mean annual temperature or isotherm of $42\frac{1}{2}^{\circ}$, is, in the northern hemisphere, the great zone of deciduous forest-trees, or those which shed their leaves in winter—hence the seasonal contrasts unknown in warmer regions. The characteristic vegetation of this zone is well seen in that of our own country, the north of France, and Germany—forests of coniferous trees (fir, pine, yew, &c.) and expanses of heath adding peculiar features to the area. The cultivation of wheat scarcely extends beyond this zone in the northern hemisphere; in the southern it is occupied chiefly by the ocean. 6. The *sub-arctic* zone, limited by the isotherm of 39° , is characterised by coniferous trees (pine, larch, spruce, juniper, &c.), poplar, beech, grasses, and heaths; and, on its northern limits, by birch, willow, and alder. 7. The *arctic* zone (which has no equivalent in the southern hemisphere, where the ocean alone prevails) is marked by the dwarf birch, alder, and willow; by occasional pines and firs; by grasses; and by numerous

lichens and mosses on its northern limits. In the American section, rhododendron, andromeda, and azalea are not unfrequent. 8. The *polar* zone is characterised chiefly by its flowerless plants—lichens and mosses—though, during its brief summer, species of *ranunculus*, *saxifrage*, *scurvy-grass*, *rush*, *willow*, &c., make their appearance. In this zone there are no trees nor bushes, nor any cultivation of plants for food.

186. Such, in general terms, is the characteristic distribution of vegetable life as we proceed from warmer to colder latitudes. But the learner is already familiar with the fact, that temperature decreases not only as we proceed from the equator towards the poles, but also as we ascend from the level of the sea into the higher regions of the atmosphere. This ascent—*hypsometical*, as it is technically termed (Gr. *hypsos*, height, and *metron*, measure)—is marked by analogous belts of vegetation; and at the equator, for instance, the traveller who ascends a lofty mountain passes through a flora much akin to that which marks the successive horizontal zones alluded to in the preceding paragraph. "We pass," says Herschel, "through the same series of climates, so far as temperature is concerned, which we should do by travelling from the same station to the polar regions of the globe; and in a country where very great differences of level exist, we find every variety of climate arranged in zones according to the altitude (hypsometical zones), and characterised by the vegetable productions appropriated to their habitual temperature." Thus Humboldt, in describing the South American Alpine flora, remarks:—"In the burning plains scarce raised above the level of the Southern Ocean, we find bananas, cycads, and palms in the greatest luxuriance; after them, shaded by the lofty sides of the valleys in the Andes, tree-ferns; and next in succession, bedewed by cool, misty clouds, cinchonas (Peruvian-bark trees) appear. When lofty trees cease, we come to aralias (ivies) and myrtle-leaved andromedas (heaths); these are succeeded by bejarias abounding in resin, and forming a purple belt around the mountains. In the stony region of the Paramos, the more lofty plants and showy flowering herbs disappear, and are succeeded by large meadows covered with grasses, on which the llama feeds. We now reach the bare traohytic rocks, on which the lowest tribes of plants flourish. Paramelias, lecidias, and leprarias (lichens), with their many-coloured thalli and fructification, form the flora of this inhospitable zone. Patches of recently-fallen snow now begin to cover the last efforts of vegetable life, and then the line of eternal snow begins."

187. In temperate latitudes, though the variety of vegetation be less, and the lower zones of tropical flora be necessarily wanting, similar phenomena present themselves. "We may begin the ascent of the Alps, for instance, in the midst of warm vineyards, and pass through a succession of oaks, sweet chestnuts, and beeches,

till we gain the elevation of the more hardy pines and stunted birches, and tread on pastures fringed by borders of perpetual snow. At the height of 1950 feet the vine disappears; and at 1000 feet higher, the sweet chestnuts cease to thrive; 1000 feet farther, and the oak is unable to maintain itself; the birch ceases to grow at an elevation of 4680 feet; and the spruce-fir at the height of 5900 feet, beyond which no tree appears. The rhododendron then covers immense tracts to the height of 7800 feet; and the herbaceous willow creeps 200 or 300 feet higher, accompanied by a few saxifrages, gentians, and grasses; while lichens and mosses struggle up to the imperishable barrier of eternal snow." The accompanying sketch exhibits, in general terms, the order of these ascending zones of vegetation, from the sea-level at the equator to the limit of perpetual snow.



188. Besides the great governing conditions of temperature, as dependent on latitude and altitude, there are others arising from light, slope, nature of soil, moisture, &c., which, though less general, are nevertheless equally imperative. The southern slopes of a mountain-range exposed to the full influence of the sun's rays exhibit not only a greater profusion of foliage and blossom, but a greater variety of species than is to be found on the northern and darker side. Even the southern side of a tree will make a larger amount of annual growth, and present a greater exuberance of flower and fruit than the northern. Again, the influence of soil is equally observable: hence we seek in vain on the clayey moorland for the species that form the rich and verdant carpeting of the calcareous district; or on the thirsty sand-dunes by the sea-shore for the flora that flourishes so freely on the alluvial meadow. Farther, some plants are truly littoral, and thrive best (like the

mangrove and cocoa-palm) within the influence of the sea-spray; while others become stunted and diseased if exposed, even for a season, to the breath of the sea-breeze. The effect of moisture on the distribution of vegetation is direct and perceptible, arid tracts being comparatively barren, while humid regions, and especially humid and warm ones, are noted for their luxuriance. Even in tropical countries, like South America, the dry season is one of torpidity, and corresponds to the winter of temperate zones—vegetation being dormant till the rainy season returns and once more renews its growth and foliage. It is owing to this influence of moisture that botanists distinguish between rainy and rainless regions, and between zones of summer rains, winter rains, and rains at all seasons—each zone being stamped by its own vegetable aspects.

189. As plants have a fixed and natural distribution on land, so also they have a similar distribution, horizontal and vertical, in the waters; though, in consequence of the greater uniformity of temperature, the marine areas are less marked and decided. *Horizontally*, the ocean has been divided into certain botanical provinces, of which our limits will merely permit the enumeration: 1. The Northern Ocean, from the pole to the 60th parallel; 2. The North Atlantic, between the 60th and 40th parallels—the great headquarters of *Fucus* proper; 3. The Mediterranean—a sub-region of the warmer temperate zone of the Atlantic, lying between the 40th and 23d parallels; 4. The tropical Atlantic, in which *Sargassum* abounds (Sargasso Sea, par. 103); 5. The Antarctic American regions, from Chili to Cape Horn, and the whole circumpolar ocean south of the 50° of S. latitude; 6. The Australian and New Zealand; 7. The Indian Ocean and Red Sea; 8. The Japan and China Seas, besides certain less decided provinces in the Pacific. *Vertically*, or *bathymetrically*, as it is termed when treating of the ocean (Gr. *bathys*, deep, *metron*, a measure), light, depth and nature of sea-bed seem to be the prime regulators of aquatic vegetation. In the limited areas of streams and fresh-water lakes the order of arrangement is less perceptible, but in the ocean each gradually-deepening zone is characterised by its own peculiar forms. Thus, in the British seas naturalists speak of a littoral, laminarian, coralline, and coral zone; and in the ocean generally, of a littoral, circum-littoral, median, infra-median, and abyssal or deep-sea zone. The nature of these bathymetrical subdivisions will be best understood by referring to our own shores, where the *littoral* is that which lies between high and low-water mark, and is characterised by such plants as the bladder-wrack, dulse, and cariceen; the *laminarian*, that which commences at low-water mark, and extends to a depth of from 40 to 60 feet, and marked, as its name implies, by the broad waving tangle (*laminaria*), and larger algæ; the *coralline*, that which extends from 90 to about 300 feet in depth, and where corallines luxuriate, and the common sea-weeds disappear; and the *coral*

zone, which lies between 300 and 600 feet, and is the region of the calcareous and stronger corals. In the abysses of the ocean diatoms (microscopic vegetable forms) alone occur; and generally as the corallines increase the true sea-weeds disappear, ordinary algae scarcely existing beyond a depth of 300 feet.

190. Admitting, however, in the fullest degree, the influence of heat, light, moisture, and the like, in the distribution of vegetable life, there still lies over and above them a primal arrangement, by which certain forms are naturally restricted to certain areas. This arrangement, which is seemingly not dependent on climate (for the plants thrive equally well when transferred to other areas), imparts a certain physiognomy to these regions; and thus botanists, entering more minutely into the geographical aspects of their science, subdivide the earth into *regions* and *provinces* according to their prevalent floras. Such subdivisions lie beyond the limits of our outline, but the learner will readily perceive their bearings when he considers—*first*, that some forms, like the tea-plant and camellia, are peculiar to Eastern Asia; some, like the eucalypti and casuarinæ, to Australia; others, like the magnolia, to the southern latitudes of North America, and so on: while, *second*, that every tribe of plants has a special aspect or physiognomy, and where such tribes prevail, that physiognomy will be imparted to the landscape. Wherever, therefore, certain orders are peculiar, and a certain number of genera and species prevail, this constitutes a botanical “region,” such as the region of saxifrages and mosses, the region of magnolias, the region of camellias and teas, region of palms, and so on, making in all some twenty-four regions, into which the earth’s surface has been botanically divided. In like manner with aspect or “physiognomy,” as the Palm form, the Banana form, the Mimosa form, Cactus form, Heath form, Grassy form, Willow form, &c.—there being in all twenty-two such forms, which are readily distinguished even by the eye of the unbotanical observer. In the same way it is not unusual to divide the Old and New World into agricultural zones, in something like the following order, from the tropics to the northern limit of profitable culture—1. Zone of rice and maize; 2. Maize and wheat; 3. Wheat, rye, buckwheat, pease, and beans; 4. Barley, oats, and potato.

191. Such, in general terms, is the distribution of plants over the surface of the globe. Though many families have a very narrow range, and naturally are never found beyond it, yet others having greater facilities for the dispersion of their seeds, and being, moreover, of a more elastic constitution, have a tendency to increase their area, and this often at the expense of other families that are destroyed by their presence. Others, again, possessing this greater adaptability, are transferred for their utility or ornament from region to region, and are now found *acclimatised* and growing luxuriantly in many countries to which they naturally would never have found their way. In

general, however, the great natural laws of distribution are supreme, and the majority of plants attain their perfection only in the habitats to which they originally belong. In densely-peopled and cultivated countries man is ever destroying, transferring, and acclimatising ; but, knowing the limit to which this power can be profitably exercised, he will cease to rear in one region what can be more abundantly and cheaply procured from another. It is thus that a knowledge of the geographical distribution of plants, and the laws on which that distribution depends, becomes a subject not only of scientific interest, but of true economic importance.

Animal Distribution.

192. Being influenced by climate, food, and other external conditions, animals, like plants, are necessarily less or more restricted to certain geographical regions. Endowed with greater powers of dispersion and locomotion, their limits are, perhaps, less precise than those of plants ; but in the main there is a similar horizontal and vertical arrangement of animal forms—from the equator to the poles, and from the sea-level to the loftiest heights of land, or to the greatest depths of ocean. The *fauna* of the tropics, taken in general terms, is more exuberant in kind, in size, strength, and beauty, than that of the temperate zones ; and this, again, more abundant than that of the arctic and antarctic regions. The more luxuriant and sheltered lowlands are peopled by races differing from those that inhabit the mountain-slopes, and those that affect the mountain-sides are distinct from those that find subsistence among the higher and colder elevations. In like manner, the creatures that throng the shallow shore are specifically different from those that are scattered through the deeper ocean.

193. In this way terrestrial animals may be arranged into a tropical, a temperate, and an arctic fauna—each shading to a certain extent into the other, but still, in the main, characterised by genera and species that do not naturally occur in the other sections. Thus, the Tropics are the great headquarters of the apes and monkeys ; of the lion, tiger, panther, and larger carnivora ; of the giraffe and zebra ; the elephant, rhinoceros, hippopotamus, and tapir ; the crocodile, turtle, boa, and larger reptiles ; the ostrich, flamingo, peacock, parrots, humming-birds, and generally of birds remarkable for their brilliant and variegated plumage. Insect life is also much more varied and exuberant in tropical than in colder latitudes—attaining its maximum in variety, in size, activity, and brilliancy of hue within the luxuriant regions of Brazil, Guinea, and the Indian Archipelago, and gradually declining towards either temperate zone. The Temperate zones, on the other hand, though marked on their warmer limits by the

presence of such tropical forms as the tiger, jackal, hyena, crocodile, &c., are, on the whole, the headquarters of such ruminants as the ox, bison, buffalo, goat, sheep, stag, and elk. Peculiar to them also are the Bactrian camel, the wild boar, wolf, fox, and beaver; the opossum in the northern hemisphere and the kangaroo in the southern; the eagle and falcon, turkey, goose, grouse, and pheasant, among birds; while reptiles become fewer and smaller the nearer we approach the arctic zone. The Arctic zone (for the antarctic is almost exclusively occupied by the ocean) is characterised by greater uniformity in its fauna, by few species but by numerous individuals, and generally by the quiet and sombre colouring both of its birds and quadrupeds. The reindeer, musk-ox, brown bear, polar bear, wolf, arctic fox, and sable, are peculiar to this region; the sea-fowl that frequent its summer seas are chiefly migrants from the waters of the colder-temperate zone; and reptile life is unknown. And here it may be observed that as the land in the northern hemisphere lies in great contiguous or all but contiguous masses, while in the southern it consists of far-separated spurs and patches, so there is a greater similarity between the fauna and flora of northern than of southern zones.

194. As with the terrestrial fauna, so also in a great measure with the marine, though at first sight there may seem no interruption to interchange and community of habitat. Variety of genera and species characterise the seas of the torrid zone; uniformity of species and immense numbers of individuals mark the fauna of the colder latitudes. The fishes and shell-fish of the tropics are noted for their varied and brilliant tints; those of the arctic regions are of uniform and sombre hues. The right whale never traverses beyond the cold waters of the higher latitudes in either hemisphere; the sperm whale, on the other hand, is unknown beyond the tropical areas of the Pacific. Unknown in the torrid zone, the seal and walrus occur in thousands in the colder-temperate and arctic regions. The headquarters of the sharks lie within the torrid zone; the tunny rejoices in the genial waters of the Mediterranean; while the cod, pilchard, herring, and salmon, attain perfection only within the colder waters of the higher latitudes. The constituents of the sea-water are nearly the same throughout, and yet the reef-building corals only elaborate their structures within the tropical and sub-tropical expanses of the ocean.

195. As with the horizontal areas of the water, so in like manner with its various depths. The *littoral zone* (par. 189) of our own seas, for example, is characterised, according as the bottom may be rocky, sandy, or muddy, by such shell-fish as the periwinkle, limpet, mussel, cockle, and razor-shell; the *lamina-rian* by star-fishes, the common sea-urchin, tubularia, modiola, and pullastra; the *coralline* by the disappearance of the ordinary shore shells, and the abundance of buccinum, fusus, trochus,

venus, pecten, and the like ; and the *coral zone* by forms of starfish, cidaris, and brachiopod mollusca that cannot exist in shallower waters. As to the extreme depths of ocean, they are generally presumed to be, like the extreme elevations of the land, altogether barren and lifeless. It is true that some recent deep-sea soundings in the North Atlantic would seem to point to a different conclusion ; but these, before final acceptance, require the confirmation of further and independent experiment.

196. It must ever be borne in mind, however, that obvious as may be the influence of external conditions on the distribution of animal life, there is over and above them an aboriginal dispersion that science in its present position is unable to account for. Why, for instance, should the kangaroo and ornithorhynchus be restricted to Australia, the hippopotamus and giraffe to Africa, the camel to the Old World, and the llama to the New ? But while these aboriginal differences exist and remain unaccounted for, the learner should remember that distant regions, having nearly the same conditions, are peopled by what are termed *representative species*—that is, species not zoologically identical, but merely representing one another in the economy of nature, and fulfilling similar functions. Thus, the camel of the Old World is represented by the llama of the New ; the lion and tiger of the Old World by the puma and jaguar of the New ; the ostrich of Africa by the rhea of South America and the emu of Australia ; and the crocodile of the Nile by the gavia of the Ganges and the alligator of the Amazon. It should also be borne in mind that local conditions of food, shelter, healthful position, freedom from enemies, and the like, are also operating causes in the distribution and dispersion of animal life ; and that where these are wanting, whole families will shift ground, or altogether disappear, even when the great conditions of temperature remain unchanged.

197. As in the vegetable world, so also (though to a less extent) in the animal, certain species have a greater elasticity of constitution, which enables them to subsist under a greater variety of conditions, and naturally, therefore, to enjoy a wider geographical range. Operating upon this principle, and a knowledge of climatology, man has been enabled to transfer from one region to another a considerable number of animals, either for the purposes of his convenience or luxury. All the domestic animals—horse, ass, ox, sheep, goat, pig, dog, cat, barn-fowl, &c. ; many birds, prized for their beauty or song ; and rats, mice, insects, and other creatures, considered as “pests and vermin,” have accompanied him over the habitable globe. His efforts in this respect—extirpating, transferring, and acclimatising—have been incessant ; and thus creatures naturally of widely distant habitats have been, and are still being, brought together into one common area. In this way the domestic animals of the Old World have been transferred to the New, where they were unknown at the time of its discovery by Columbus ; some of the New World fauna trans-

ferred to the Old ; and not only the domestic animals, but the birds, fish, and even shell-fish of Europe, transported to Australasia, where, within little more than a century, their genera were totally unrepresented. But as in the vegetable world, so in the animal, there is a limit to this system of transference, and man best studies his own interest and the comfort of the lower animals by fostering them mainly within their own native habitats, and sharing in their larger produce through the more profitable method of commerce and exchange.

Interdependence of Plants and Animals.

198. As plants and animals are alike dependent on external conditions, so both are, to a certain extent, dependent on one another. Both, for example, are dependent on the atmosphere, yet the oxygen which the plant exhales is inhaled by the animal, and the carbonic acid exhaled by the animal is absorbed and assimilated by the plant. The plant, rooted in the soil and casting abroad its leaves and branches in the atmosphere, though seemingly deriving the main elements of its growth from inorganic sources, is nevertheless stimulated into life and exuberance by the presence of organic decay, and many of the lower fungus-growths are found only where such decomposing matter is present. Herbivorous animals, as is well known, subsist directly upon plants, while the carnivorous prey upon the plant-feeders, and are thus also ultimately dependent on the vegetable world for their subsistence. Wherever vegetable life is varied and luxuriant, there animal life is marked by a corresponding variety ; hence the specific exuberance of the tropics compared with that of the colder latitudes. By the extirpation of certain plants, certain mammals, birds, and insects may be removed from a district ; while, on the introduction of some new exotic, animals hitherto unknown in that locality usually make their appearance. The law of circulation and interdependence is complete, and no portion of the circle could be removed without a corresponding change in the characters of the vegetable and animal kingdoms.

199. Again, though most plants have the inherent power of dispersing their own seeds, and are aided in this by winds and water-currents, yet many depend upon birds and mammals for their wider dispersion and increase, just as many depend upon insects for the fertilisation of their flowers. This wider dispersion creates a new source of subsistence for the animals that feed upon them, and thus the increased area of the one supplies a wider range to the other. Further, as some animals are fitted by their organisation for an arboreal existence, some for life on the grassy plain, and others for the shrubby thicket, the destruction of the tree, the planting of the plain, or the clearing of the thicket, would necessarily involve the destruction of these special

organisations. Still further, as some creatures are specially fitted to live on fruits, some on leaves, and others on roots, the disappearance of these specific supplies would necessarily be followed by the annihilation of the consumers. It is in this manner that plants and animals become co-dependent portions of one great vital plan, and that geographers, aware of these relations, can more intelligibly depict the aspects of nature by associating every fauna with its own appropriate and distinctive flora.

RECAPITULATION.

In the preceding chapter attention has been directed to the vital aspects of the globe—that is, to its vegetable and animal life, and their distribution over the land and through the waters. Leaving the nature and origin of Life to the Biologist, it has been shown that plants and animals are dependent for their continuance on the external conditions by which they are surrounded, and that any change in these conditions would materially affect, if not destroy, their existence. For this reason, plants and animals have a definite distribution over the globe; heat, light, and moisture being the great regulators of the one; climate and food the governing conditions of the other. In this way vital variety and exuberance culminate within the tropics and decline as we proceed towards either pole—declining also in an analogous manner as we ascend from the level of the sea into the higher elevations of mountains. Each zone or belt of the earth has thus its own special flora and fauna—that is, is characterised by genera and species not naturally occurring in other regions. As marked variations occur within narrower limits than the torrid, temperate, and frigid zones, botanists subdivide the earth's surface into equatorial, tropical, sub-tropical, warmer-temperate, colder-temperate, sub-arctic, arctic, and polar belts; and these, as might be expected, correspond with lines of temperature rather than with parallels of latitude. As in the land, so also in the waters, each zone of depth, from the shore seaward, has its characteristic forms—littoral, laminarian, coralline, and coral; the extreme abysses of ocean, like the extreme altitudes of land, being apparently barren and lifeless. Beyond these main horizontal and vertical arrangements of life, there is also an aboriginal dispersion of certain races over certain areas which science cannot account for; hence the subdivision of the earth's surface into botanical and zoological "regions" and "provinces,"

each subdivision being characterised by its own typical forms. These provinces form the special study of the botanist and zoologist; and the question how far the plants and animals of one region can be profitably transferred to another, becomes one of prime economical importance. Understanding the geographical conditions under which plants and animals naturally occur, and knowing, moreover, the intimate dependence between the vegetable and animal kingdoms, man, in his own migrations over the globe, will be better able to determine what to cultivate and what to extirpate, what to attempt acclimatising and what to continue in their own native habitats. Understanding, moreover, the relations that subsist between fauna and flora, he will, as a geographer, be better enabled to draw that intelligible picture of external aspects which it is the grand province of his science to depict.

XII.

ETHNOLOGY—RACES AND VARIETIES OF MAN.

Man as affected by External Conditions.

200. MAN, in whatever stage of civilisation he may appear, is always less or more influenced by the geographical conditions of the region he occupies. Unlike the lower animals, which either simply flourish under or succumb to these conditions, Man may struggle against and so far modify them; but still, to a great extent, his thoughts and actions, his industrial pursuits, his social polity, and religious beliefs, are all affected by the physical circumstances of his position. In savage life this influence is direct and perceptible; hence the difference between the semi-aquatic Esquimaux and the hunting Red Indian, though inhabiting the same continent; between the stationary vegetable-feeding islanders of the sunny Pacific, and the wandering omnivorous tribes of the scrubby plains of Australia. Even where civilisation has made some progress, it is these conditions that still mainly determine man's habits and pursuits—rendering the inhabitants of the grassy steppe nomades and herdsmen, the indwellers of the river-plain tillers of the ground and growers of grain, and the men of the sea-coast traders and adventurers. And higher still, where populations have been long settled and civilisation has assumed its most advanced aspects, climate, scenery, natural products, facilities for intercommunication and exchange, are ever exercising their influence—rendering one nation wealthy and independent, another bold and enterprising, and a third, it may be, isolated and stationary. Nor is it man's mere material condition that is thus affected; his religious sentiments, his poetic feelings, his love of liberty, his social government, are all less or more tinged by the nature of the physical characteristics of the country he inhabits.

201. That such is the case, the most cursory glance at the different nationalities of the world will readily convince; and though the inherent qualities of Race, for reasons we cannot now discover, may differ very widely, still over and above these qualities external conditions exert a direct and perceptible modifying influence.

The white men of Europe may differ physically and intellectually from the black tribes of Africa and the red races of North America ; but it may be fairly questioned whether the former would have ever exhibited their present activity and progress had it not been for the greater varieties of surface, climate, and general physical conditions that Europe, as a continent, enjoys. There can be no doubt that the moderate climate of Britain is more favourable to bodily and mental vigour than the relaxing temperature of the tropics ; and that the slight seasonal differences between our spring, summer, autumn, and winter, induce habits of continuous exertion and industry unknown in countries subjected to excessive summer's heat and winter's cold. But for our insular situation, our countrymen would never have been the traders and adventurers they have become ; and but for our natural supplies of coal, iron, and other metals, the mechanical and manufacturing character that now stamps the British nation would have been unattainable and impossible. So much does the general character of a people depend upon the physical conditions of the country they inhabit !

202. Even in their minor peculiarities, the different nations of the same great race are similarly affected by external conditions ; hence the obvious distinctions existing between the livelier and more versatile nations of southern Europe and the graver inhabitants of the north ; between the bold and independent mountaineers of Switzerland, Scotland, and Scandinavia, and the tamer occupiers of the central and eastern European plain. The language of everyday life is full of these distinctions, and this long before science had attempted their explanation. Thus we speak of the "dry, clear, exhilarating air" of one district, and the "damp, cloudy, and depressing atmosphere" of another : of the "dreary monotony" of one region, and the "charming variety" of another ; of the "awe-inspiring gloom" of the forest, and the "cheerful hues" of the open landscape. Indeed it is to the influence of situation that we are in a great measure to look for national peculiarities—these peculiarities diminishing the more that nations extend their range of intercommunication, and the less they are restricted to their own narrow boundaries.

Characteristics and Distribution of Races.

203. Whatever the influence of external conditions in modifying the characteristics of race, we find Man distributed at present over almost every region of the globe—wandering in savage freedom under the tropics, flourishing in busy communities within the temperate zones, and struggling in diminished numbers against the inclemencies of the polar regions. Within the tropics he is chiefly a vegetable feeder ; in the temperate zones he adopts a mixed vegetable and animal diet ; while within the polar circle

his diet is exclusively animal. But while in this respect he obeys, like plants and animals, the zonal arrangements of the world, unlike them the varieties of his race are distributed according to no law of latitudinal dispersion. As in districts of the same country we find differences of stature, physiognomy, dialect, and habits; so in the various countries of the same continent we find still wider differences in bodily appearance, mental constitution, language, and manners. Notwithstanding these well-known distinctions, there is among the inhabitants of certain regions a certain sameness in physical aspect, in colour of skin, in form of head and face, and also in mental disposition, that stamps them as distinct from the inhabitants of other regions; and hence arises the idea of *varieties* or *races* of the human species.

204. Without entering upon the vexed questions of man's origin and antiquity—whether he is a mere development from the lower animals, according to some great natural plan, or an entirely new creation—whether he has been six or sixty thousand years an inhabitant of the earth—and whether the varieties of our race have descended from a single pair, and been since modified by external conditions, or are the progeny of several independent pairs—it may be stated as the general opinion of naturalists that mankind form, in the zoological scale, a single species of a single genus. But though thus standing unique, and far exalted above other animals by his gifts of reason, moral perception, and religious sentiment, it is necessary and natural to divide mankind into several varieties according to their more prominent physical features; and Ethnology (Gr. *ethnos*, a race; *logos*, a discourse), extending the subject to minor features, language, and the like, still further subdivides these varieties into groups, tribes, and branches. The consideration of these minor distinctions (which are evidently produced by intermixture of races and the influence of external conditions) belongs more especially to Ethnology; our limits will merely permit a brief allusion to the *five* varieties or races into which the inhabitants of the globe have been arranged by the German philosopher, Blumenbach. These are the *Caucasian*, the *Mongolian*, the *American*, the *Ethiopian*, and the *Malay*; each being characterised by some peculiarity in colour of skin, eyes, nature of hair (curled, lank, woolly, frizzled) shape of skull, form of face, and general physiognomy. Of these characteristics colour of skin is one of the most obvious, and, though varying in shade even in the same race, is yet employed in everyday language as a main mark of distinction—the *Caucasian* being the *white*, the *Mongolian* the *yellow*, the *American* the *red*, the *Ethiopian* the *black*, and the *Malay* the *brown*. The distribution and more obvious characteristics of these respective races we epitomise from Blumenbach and other ethnologists:—

205. The *Caucasian* variety is dispersed over Turkey, Arabia, Persia, part of Tartary, Afghanistan, and Hindostan, in Asia; over Egypt, Abyssinia, and the Mediterranean seaboard, in Af-

rica; and over almost the whole of Europe—the Turks proper, the Magyars, Finns, and Laplanders, being of Mongolian origin. Within the last three centuries the race has spread from Europe over large areas of North and South America, South Africa, Australia, and New Zealand (see Ethnographic Map in Atlas); and wherever it has planted itself becomes the dominating power. This variety takes its name from the Caucasian mountain-range that stretches between the Caspian and Black Sea, because tradition points to that region as the place of its origin; and it is also known by the term *Indo-European*, from its spreading over India and Europe from the Ganges to the further shores of Iceland. The more important branches are the Hindoo, Persian, Arab, Circassian, Slavonic, Teutonic, and Celtic, with their various mixtures and alliances, which now constitute the nationalities of Southern Asia and Europe. The distinguishing characteristics of these Indo-Europeans are—a light-coloured skin, varying from fair to tawny or swarthy; red cheeks; copious, soft, flowing hair, generally curled or waving; ample beard; small, oval, and straight face, with features distinct; expanded forehead; large and elevated cranium; narrow nose; and small mouth. In stature the Caucasian is taller than any of the other varieties; of erect gait; with rounded, well-proportioned limbs; moderately small extremities; and light elastic step. The *White* race (for by this designation it is generally distinguished) has given birth to the most civilised nations of ancient and modern times; and has hitherto exhibited the intellectual and moral powers of human nature in their highest degree. Wherever the white man has established himself, the other races disappear before him. His proper field of development, however, seems to be the temperate zones in either hemisphere, for beyond them he degenerates physically and intellectually, and loses the higher characteristics of his race.

206. The *Mongolian* variety is spread, as the name implies, over the central and northern regions of Asia, China, Japan, Burmah, Siam, and Cochin-China; and includes, moreover, the scattered inhabitants of the Arctic seaboard, both in the Old and New World continents. It embraces the Mongols, Turks, Tibetans, Chinese, Indo-Chinese, Japanese, Kamtchatdales, Tungusians, Koriaks, and Samoiedes, in Asia; the Turks, Finns, and Lapps, in Europe; and the Esquimaux of the North American arctic regions. In the Mongolian, the skin is olive-yellow; the hair thin, coarse, and straight; little or no beard; head or cranium somewhat square-shaped; forehead rather low; face broad and flattened, with confluent features; high cheek-bones; eyes rather sunk, and obliquely directed towards the nose; wide and small nose; and thick lips. In stature the Mongolian is below the Indo-European; but in the true Tartar, Turk, and Chinese, the frame is broad, square-set, and robust, with high shoulders, and short and strong neck. In intellectual and moral capacity the

various branches of the race differ very widely ; but on the whole they are inferior, less energetic, and more stationary in their civilisation than the Caucasian variety. Their highest attainments appear in the Chinese and Japanese ; the Turks and Magyars having been so long amalgamated with Europeans as to assume a Western rather than an Oriental phase of civilisation.

207. Under the *American* variety, ethnologists comprise all the aboriginal races which peopled the New World prior to its discovery by Columbus in 1492. These are the Chippewyans, Sioux, Pawnees, Cherokees, and other tribes in North America ; and the Caribs, Solimoes, Guaranis, Araucanians, Patagonians, and Fuegians, in South America. In this race the skin is reddish or copper-coloured (hence the familiar designation of Red Indian) ; the hair is black, coarse, and lank ; beard scanty ; skull somewhat similar to the Mongolian, but narrower, and not so square ; forehead low and retreating ; cheek-bones prominent, but more rounded than those of the Mongol ; eyes sunk, and somewhat raised at their outer angle ; nose and other features rather prominent. In stature the North American Indian is rather tall ; spare and lithe in body ; and as a hunter, acute in his senses, and remarkable for his endurance of fatigue and insensibility to pain. In South America the race greatly degenerates ; the Guaranis, the Fuegians, and other tribes being amongst the most degraded of savages, their wretched appearance being in many instances aggravated by artificial distortion of the head and facial features. In intellectual and moral qualities the American Indians resemble in many respects the Mongolians. Like the Mongols, they have remained stationary, but at a much lower point of civilisation, if we except the ancient occupants of Mexico and Peru. In North America, the race is rapidly disappearing before the white settlers ; in South America, less fitted for the White, the inferior and more sparsely scattered tribes have been little interfered with.

208. The *Ethiopian* race, in one or other of its branches, inhabits the whole of Africa, with the exception of Egypt, Abyssinia, Tunis, Tripoli, and Morocco on the north, and Cape Colony on the south. It embraces Hottentots, Bushmen, Kaffirs, Negroes, Gallas, Tibboos, Mandingoes, and other tribes, differing widely among themselves in physical and mental aptitude, but all, from time immemorial, remaining in a barbarous or but very partially civilised state. In this variety, which derives its name from the Ethiopia of the ancients, the skin is black (hence the familiar designation, Negro—Lat. *niger*, black) ; the hair short, black, and woolly ; skull compressed on the sides, and elongated towards the front ; forehead low, narrow, and slanting ; cheek-bones very prominent ; nose broad and flat ; jaws projecting so as to make the upper front teeth oblique ; lips, especially the upper one, very thick. In stature there is great differences among the different branches of the race—some, like the Kaffir, being



1, Caucasian; 2, Mongolian; 3, Ethiopian; 4, Malay; 5, American.

of average size, and fairly formed; others, like the true negro, also of average size, but ungainly in form and limb, with large flat feet and hands, and shuffling awkward gait; and some, again, like the Hottentot and Bushmen, of stunted stature, and with slender ill-formed limbs. Intellectually, the Black race has ever remained in a rude and barbarous state; hence the subjection of one branch to another branch among themselves; and hence also their enslavement, from time immemorial, by the white variety. This is not the place to enter upon the question, how far the negro is capable of attaining the higher phases of civilisation; but the fact remains, that neither of himself, nor in any of his admixtures with other races, has he shown much aptitude for intellectual or social advancement.

209. The *Malay* race (so called from the Malayan peninsula) includes the widely-scattered and chiefly insular inhabitants of Malaysia, Australasia, and Polynesia. The population of these widely-spread districts differ, as might be expected, very widely from each other—those of Malaysia and Polynesia being of a brown or lighter colour, and somewhat resembling the Mongolians; while those of New Guinea and Australia (the Papuan negroes, as they are termed) are of a dark colour, and more closely approximated to the negro type. Intellectually, also, there is a similar difference, there being little in common between the Malay of the Indian Archipelago and the savage inhabitant of New Guinea; between the stunted and miserable native of Australia, and the daring, apt, and clever inhabitant of New Zealand. Taken, however, as described by ethnologists, the skin of the Malay varies from a light tawny to a deep-brown; hair black, crisp, and somewhat inclined to curl in the true Malay, and tufted and frizzled in the Papuan; head rather narrow; bones of the face large and prominent; nose full and broad towards the lips. In civilisation the Malay race has hitherto made little advancement. In the Indian Archipelago it has, like the Mongolian, long remained stationary; in Polynesia, some progress appears under tutelage of the white, though the areas are too widely separated ever to be of much importance; in New Zealand it seems reluctant to amalgamate with the new settlers; while in Australia it is rapidly dying out before the encroachment of the European immigrant.

210. Taking these varieties as aids to the *general* arrangement and consideration of mankind, it must be remembered that there are not only admixtures among conterminous races (Caucasian and Mongol, Mongol and Malay, Mongol and American), but admixtures also between immigrant and native races (Brazilians, Peruvians, Mexicans, &c.), which render sharp lines of demarcation impossible, and unnecessary as impossible. The consideration of these *minutiae* belongs more especially to Ethnography and Philology, by which not only shades of colour, types of skull, facial angles, and the like, are taken into account, but

dialects of language, mental peculiarity, and forms of government, are all considered in tracing the dispersion, affinities, and history of the human race. Such considerations lie beyond the scope of our Outline, and the learner may consider as sufficient for his purpose the *five* great varieties above enumerated, or even their abridgment into *three* (Caucasian, Mongolian, and Ethiopian—reckoning the American as a sub-variety of the Mongol, and the Malay a sub-variety partly of the Mongol and partly of the Negro), as all that is warranted or required at this stage of his progress.

Conditions of Civilisation and Progress.

211. Such is the usual subdivision of mankind into varieties or races, and such the existing distribution of these races over the surface of the globe. The subdivisions may to some extent be arbitrary; but as there are actual differences of colour, form of head, facial expression, and the like, and as these physical features are accompanied by strongly-marked differences in mental constitution, the distinctions in the main must be founded on nature. As to the prehistoric distribution of man, neither archæology nor geology, in the absence of reliable remains, can arrive at any certain conclusion. Within historical time, however, the various races, while peopling most densely the regions they now occupy, have ever been less or more encroaching on each other's domain—the inferior giving way to the superior and more civilised. The Ethiopian, in its numerous tribes and branches, has remained stationary in Africa. The Malay, chiefly an insular race, has spread itself, in one or other of its branches, over the islands of the Southern Ocean and Pacific. The Mongolian, while claiming Central and Eastern Asia as its headquarters, has spread partially into Europe, and largely along the entire seaboard of the Arctic Ocean. The White man, on the other hand, has partly repelled the Mongol, and, after spreading wave after wave over Western Asia, Northern Africa, and the whole of Europe, has within the last three centuries taken possession of the greater portion of North America (United States, Canada, Mexico, California, &c.); of part of South America (Brazil, Guiana, Peru, &c.); of South Africa; and of Southern Australia and New Zealand; while his influence is felt, less or more, in every region of the globe.

212. Wherever the means of subsistence can be obtained, there man will establish himself and increase in numbers—his increase being mainly regulated by the facility or difficulty of obtaining supplies. Where he can raise more than his own wants require (and this will depend very much on his knowledge of nature's laws and operations), or where he enjoys products not possessed by other localities, this surplus and these products

form subjects of barter and exchange, and thus he acquires wealth and the power to command luxuries. Raised above the mere physical struggle for existence, the higher faculties of his mind—imagination, invention, reflection, moral perception, and religious sentiment—begin to develop themselves, and man passes from the domain of savagery into the categories of civilisation. To trace the course of civilisation lies beyond the scope of our subject; but it is evident from what has been repeatedly stated, that its advancement (laying altogether aside the consideration of Race) depends, in a great measure, on the geographical or physical conditions by which it is surrounded. Wherever there exists a favourable climate, the means of subsistence, and opportunities of interchange and barter—in other words, wherever there are the objects and means of successful industry—there civilisation will manifest itself; and just as equitable laws, protection of property, freedom of action, and liberty of opinion are enjoyed, so civilisation will advance in a corresponding ratio. In absence of these adjuncts it has passed, and may yet pass, from nation to nation; but in the aggregate its maintenance from epoch to epoch has been secure—its progress seems illimitable.

RECAPITULATION.

In the preceding chapter attention has been briefly directed to the varieties and distribution of the human species—Ethnology or Ethnography being the science which treats of these distinctions. It has been shown that Man, though possessing a greater elasticity of constitution than most of the lower animals, and capable of enduring under almost every climate, is still, to a great extent, influenced by the external conditions by which he is surrounded, both in his physical and mental relations. Whether owing to these influences, or arising from aboriginal differences which science cannot explain, Man now appears in several *varieties* or *races*—each occupying well-marked territories on the globe, and distinguished by peculiarities of colour, form of head, facial expression, and other *physical* features, as well as by equally obvious *intellectual* and *social* qualities. These varieties are the *Caucasian* or *white*, the *Mongolian* or *yellow*, the *American* or *red*, the *Ethiopian* or *black*, and the *Malay* or *brown*—each embracing a great number of tribes, branches, and nationalities, differing in language, social polity, and other peculiarities. The Caucasians, or Indo-Europeans, inhabit the south-western section of Asia, the

northern belt of Africa, and nearly the whole of Europe, and have, in modern times, extended their dominion to large areas of North and South America, South Africa, South Australia, and New Zealand. The Mongolians, concentrated chiefly in Central and Eastern Asia, have partially penetrated into Eastern Europe, but are most extensively spread in scattered communities along the entire seaboard of the Arctic Ocean. The Malays, having their headquarters in Malaya and the Indian Archipelago, are spread, in one or other of their tribes, over Australia, as well as over all the island-groups that stud the bosom of the Pacific. The Ethiopians, though drafted hither and thither as the slaves of the white man, have been mainly stationary in Africa, the natural home and habitat of their race; while the Americans, or Red Indians, have been equally restricted to the New World continent. As a whole, the negro and Red Indian have made, and still make, the least progress in civilisation; hence their easier subjection by the higher races, and hence also their rapid disappearance before them. The Malay and Mongolian come next in order; and though some of their sections (Chinese, Japanese, &c.) have arrived at considerable eminence in the industrial arts, yet in both races the essentials of higher progress seem wanting, and hence their torpid and stationary aspect. By the white race alone do we find displayed the higher efforts of bodily and mental activity—hence in ancient times the civilisations of India, Assyria, Palestine, Egypt, Phœnicia, Greece, and Rome; and in modern times those of Western Europe—Austria, Germany, France, and Britain; of the United States and Canada; and, generally, of our colonies, in whatever region of the globe they may happen to be established. The conditions favourable to this civilisation are partly of a geographical or physical nature, and partly intellectual; and wherever the two are in fortunate conjunction, there the progress of the white race is certain, and illimitable as certain.

XIII.

GENERAL REVIEW AND DEDUCTIONS.

213. THE object of Physical Geography being not only to describe the external aspects of nature, but to determine the causes by which these aspects are produced, it has been necessary in the preceding chapters to deal with principles as well as details. Indeed, in an elementary outline the principles are of more value than the descriptions, the learner being readily enabled to extend the latter by his own readings of voyages and travels, but only intelligibly so when he has mastered the principles upon which the phenomena depend. The aim throughout has been to present our planet as subjected to general laws, believing that when these are understood there will be little difficulty in comprehending their modifications in local and limited areas. Let the learner clearly understand the origin, for example, of winds and rains—of tides and currents, and he will soon find his way to determine the causes that produce their modification in any special locality.

214. In accordance with these views, attention has been directed to the planetary relations of the earth—its figure, motions, dimensions, &c.—as on these depend its light and heat; its alternations of day and night, summer and winter; and, in general terms, all that gives rise to change and diversity in its external conditions. It was necessary, also, to consider the individual structure and composition of the globe, and the geological changes which, under the operation of aqueous, igneous, and meteoric forces, that structure has been and is still undergoing. Seizing upon its present terraqueous aspects (its lands and waters), the more immediate province of our science was to investigate their relative distribution—the disposition, highlands, and lowlands of the one, and the configuration, waves, tides, and currents of the other. Understanding the relations of land and water, and their necessary actions and reactions on each other, attention was next directed to the atmosphere by which both are surrounded, through which the light and heat of the sun are diffused to both, and in which are elaborated the winds, rains, frosts, snows, and other phenomena that constitute the essentials of weather and climate.

215. The consideration of this earth revolving through space as part of the solar system, composed superficially of land and water, surrounded by an atmosphere, and receiving heat, light, and other influences from the great central orb, constitute what may be termed the *physical or inorganic aspects* of geography. On the other hand, the study of the earth as peopled by plants and animals, and inhabited by man, capable of applying the whole to his own social and moral advantage, form its *vital or organic features*. The study of the vital world—the distribution of its plants and animals, the causes concerned in that arrangement, and the dependence of the one kingdom on the other, is one of the most attractive themes in geography. Still higher, however, and of more immediate interest, is that which deals with our own race—their physical and mental peculiarities, their dispersion over the globe, and the question how far geographical conditions may influence their social, mental, and moral progress.

216. But our science has its practical as well as theoretical importance—its economic as well as its scientific aspects. A knowledge of the distribution of sea and land—the winds, tides, and currents of the one, and of the highlands, lowlands, and climate of the other—is indispensable to successful navigation, commerce, and agriculture. The duty of determining the earth's mineral, vegetable, and animal products, and how far they can be rendered available for the purposes of everyday life, is not less imperative than the solution of its physical and vital problems. It is of vast importance to know, but it is not less important to learn how to apply. All knowledge of nature is good of itself, but its value is doubly enhanced when it is made to minister to our common humanity. Combining, therefore, its theoretical with its practical bearings, and remembering that every country is characterised by its own natural products, and has consequently something that no other country can supply, Physical Geography has paramount claims alike on the attention of the philosopher, the statesman, the sailor, the farmer, the merchant, and manufacturer.

217. Such, in general terms, is the aim and scope of Physical Geography; but it must be borne in mind that, subjected to the aqueous and igneous forces described in Chapter III., the aspects of the globe are continually undergoing change and modification. These changes may be all but imperceptible even for several generations; but in process of time they become sufficiently apparent, and stamp new features on the external conditions of the globe. The air and water, ever wearing and wasting the rocky surface in one district, and collecting the eroded material in another—the earthquake and volcano, elevating the solid crust in one region and depressing it in another, are ever changing the relative positions of sea and land; and thus the seas and lands of the future must differ from the seas and lands of the present, as

those of the present differ from what Geology assures us existed in time past. Every change of the external relations of sea and land implies a corresponding change in the nature and distribution of plants and animals; and here, again, the life of the future must differ from that of the present, as the present differs from that revealed to us by the researches of Geology.

218. In this way the student of Physical Geography must regard the earth and all its relations—physical and vital—as in a state of incessant change and progress. The changes may be slight—so slight as to pass unobserved—yet still they are not on that account less real or certain. Bearing these facts in mind, we can more readily comprehend why the rocks (the old sea-sediments) of one region should differ from those of another, why the mountain aspects of the one should differ from those of the other, and why one tract should be low and alluvial, while another is high and rocky—each feature being the product of a certain time and change in the earth's relations. It is also by bearing in mind this great feature of incessant change that we can account for the similarities and differences that exist between the flora and fauna of different regions—regions that may at one time have been continuously connected by land, though now widely separated by expanses of ocean. The plants and animals of Britain, for example, claim kindred with those of Europe, and this through land connections that existed long before their separation by the Strait of Dover. It is, in fine, by carrying this idea of mutation into all our reasonings, that Physical Geography becomes a more intelligible part of World-history—connecting the past with the present, and bearing the present into the inevitably approaching future.

219. As formerly stated, the chief agents concerned in the modification of the earth are the winds, rains, frosts, and snows of the atmosphere, which are ever loosening and disintegrating the exposed rocky surface; the streams, rivers, waves, tides, and ocean-currents, which are incessantly wearing and wasting and re-depositing the eroded materials along the bottom of the sea; and the earthquake and volcano, whose operations are alike incessant in upheaving and depressing the crust—now raising the sea-bed into dry land, and now submerging the dry land beneath the waters. The consideration of these agencies belongs more especially to Geology; but their results bear so directly on the terraqueous surface that they cannot possibly be overlooked by the geographical observer. Some idea of the magnitude of their combined influences may be formed by reflecting on the number of streams and rivers that are ever coursing over the earth's surface—or the extent of coast-line exposed to the action of waves and currents—and on the areas in Iceland, Italy, Central Asia, Indian Archipelago, Phillipine, Japan, and Aleutian Islands, Mexico, West Indies, Andes, and the Pacific islands (see Map of Volcanoes in Atlas), that are subjected to earthquake and volcanic

disturbance. Notwithstanding these continual operations, their results, for the time being, do not greatly affect the general relations of the earth; and thus Geography is enabled to depict, within appreciable limits, its existing aspects in an intelligible and available form.

220. It is this intelligible account of existing appearances, and the causes concerned in their production, that constitutes the sum and substance of our science. And just as this account is founded on correct observation, and harmonises with sound induction, so the more rapidly will Physical Geography attain to perfection. "It is not enough for it" (in the eloquent language of M. Guyot) "coldly to *anatomise* the globe, by merely taking cognisance of the arrangement of the various parts which compose it. It must endeavour to seize those incessant mutual actions of the different portions of physical nature upon each other—of inorganic upon organised beings, upon man in particular, and upon the successive developments of human societies; in a word, studying the reciprocal action of all these forces, the perpetual play of which constitutes what may be called the Life of the Globe, it should, if I may venture to say so, take up its Physiology." Understanding it in this light, the scheme of nature becomes invested with new interest, its various portions assume new harmony and unity, and man, in his physical, social, and moral development, stands in a clearer and closer relationship to the whole.

GLOSSARIAL INDEX.

*. The figures refer to the paragraphs of the text in which the particular term or subject occurs.

- ABYSSINIAN mountain-system, character of, 57.
 Acclimatisation of vegetable forms, 191.
 African mountain-systems, description of, 57.
 Agricultural zones of vegetation, 190.
 Aiguilles (Fr.), or needle-tops, defined, 47.
 Alleghany Mountains, Appalachian system, 58.
 Alpine mountain-system and its chains, 51.
 Amazon, leading characteristics and tributaries, 125.
 American variety of human race, 207.
 Andes, mountain-system of, described, 61.
 Animals and plants, mutual dependence of, 198.
 Animals, distribution of, over the globe, 192-197.
 Antipodes (Gr. *anti*, opposite; *pous*, *podos*, foot), 202.
 Apennine and Sub-Apennine mountains, 51.
 Appalachian mountain-system, nature of, 59.
 Aqueous agents of change, 216.
 Aralo-Caspian basin, its characteristics, 181.
 Archipelagoes, arrangement of islands in, 41.
 Arctic and Antarctic Oceans, areas and leading features, 82.
 Arctic and Antarctic circles, 19.
 Arctic zone of vegetation, character of, 185.
 Armenia, mountain plateaux of, 64.
 Asiatic mountain-systems, western section, 53; south-eastern system, 54; eastern system, 55; northern system, 56.
 Atlantic Ocean, area and leading features, 79; currents of, 98-101.
 Atlas system of mountains, character of, 57.
 Atmospheric agents of change, 216.
 Atmosphere, its nature and constitution, 139-143; pressure of, 141; extent of, 140; heat and moisture of, 144-150; composition, density, and character of, 17.
 Avalanches, nature of, 173.
 BALKAN Mountains, Alpine system, 51.
 Bathymetrical zones of vegetation, 189.
 Basin, river-basin, drainage-basin, &c., 116.
 Beach or strand, definition of, 32.
 Boiling-point of water, 141.
 Bolivian table-land, character of, 66.
 Botanical regions and provinces, 190.
 Brahmapootra, leading characteristics of, 128.
 Brazilian mountain-system, described, 62.
 Breezes, sea and land, origin of, 159.
 British mountain-system and its members, 50.
 CALMS, zone of, and variable winds, 154.
 Carboniferous or coal formation, character of, 28.
 Carnivorous and herbivorous, dependence between, 198.
 Carpathian mountain-chain, Alpine system, 61.
 Carse, or alluvial river-flats of Scotland, 68.
 Caspian Sea, depressed area of, 76.
 Castilian or Spanish table-land, 65.
 Caucasian mountain-system, its nature, 52.
 Caucasian variety of human race, 205.
 Centrifugal force, effects of, on earth's form, 12.
 China, principal rivers of, 127; the low river-plains of, 70.
 Circles, greater and lesser, definition of, 20.
 Civilisation, conditions of its progress, 210, 211.
 Climate, causes affecting its character, 175-179.
 Climates, insular and continental, 147, 177.
 Climatology of the globe, 139-179.
 Clouds, formation and character of, 165.
 Coast-line, importance of irregularity in, 41; nature and character of, 33.
 Colorado river, characteristics of, 127.
 Columbia river, chief features of, 127.
 Cone, mountain-cone, character of, 47.
 Continent (Lat. *con*, and *tenens*, holding together), 34.
 Continental river-basins, features of, 181.

- Continents or quarters of the globe, areas of, 34.
 Continents, their contour, relief, and position, 40-45.
 Contour or horizontal disposition, 41, 42.
 Contouring, method of, Recapitulation, p. 50.
 Coral zone of life in the ocean, 189, 195.
 Coralline zone of life, explanation of, 189, 195.
 Co-tidal lines, map of, 92.
 Counter-currents, nature and definition of, 93.
 Craig and tail, phenomenon of, described, 47.
 Cretaceous or chalk formation, character of, 28.
 Crust of the earth, structure and formation of, 22-30.
 Cultivated plants, extension of, 197.
 Currents of the ocean, nature and courses of, 98-107.
 Cyclones, or circular wind-storms, 163.
 DANUBE, the, leading characteristics of, 123.
 Day, cause and duration of, 10.
 Dead Sea, remarkable depression of, 96.
 Deccan, triangular table-land of, 64.
 Deltas and alluvial river-flats, 68.
 Density of globe, and of its rocky crust, 15.
 Devonian, or old red sandstone system, 28.
 Dew and dew-point, nature of, 164.
 Dome, mountain-dome, form and definition, 47.
 Domesticated animals, extension of, 197.
 Dovrefeld Mountains, Scandinavian system, 52.
 EARTH, the, its motions, daily and annual, 10; figure and dimensions, 11-14; density, temperature, &c., 15-17; technical subdivisions, 18-21.
 Earthquake-waves, destructive nature of, 89.
 Earthquakes, as agents of terrestrial change, 216.
 Elevation, mean, of the respective continents, 45.
 Elevation of land, effects of, on temperature, 43.
 Equatorial zone of vegetation, limits of, 185.
 Equator (Lat. *æquus*, equal), defined.
 Estuary (Lat.) or firth, definition of, 37.
 Etesian or annual winds of Europe, 161.
 Ethiopian variety of mankind, 208.
 Ethnology, races and varieties of man, 200-210.
 Europe, area of, 34; mountains of, 50-52; plains of, 69; rivers of, 122, 123.
 Evaporation of moisture from sea and land, 146.
 Exotic and indigenous, meaning of the terms, 188.
 FAUNA (Lat.), definition of, 181.
 Florida, or narrow rocky inlets of the sea, 37.
 Flora (Lat.), application of the term, 181.
 Fogs, nature and cause of, 165.
 Formations composing the earth's crust, 28.
 Fossils, their nature and geological value, 28.
 Frazer river, main features of, 127.
 Ganges, characteristics and tributaries of, 128.
 Geography, definition of, 1; Mathematical, Political, Descriptive, and Physical, 2; Physical, 3.
 Geology and Physical Geography, connection between, 30.
 Germanic plain, its physical features, 69.
 Ghants, eastern and western, mountain-chains, 54.
 Glaciers, nature and formation of, 173.
 Gobi or Shamo, arid desert of, 64.
 Grampian mountain-range, British system, 50.
 Granitic rocks, nature and composition of, 29.
 HARMATTAN or hot wind of Senegambia, 161.
 Heat, absorption and radiation of, 146, 147; origin and nature of, 172.
 Herbivorous and carnivorous, dependence between, 198.
 Highlands of the world, 46-66.
 Hill and hillocks, definition of, 47.
 Himalayan Mountains, description of, 54.
 Hindostan, great river-plain of, 70.
 Hindu Koh or Hindoo Kosh Mountains, 53.
 Hoang-ho or Yellow river of China, 127.
 Hoar-frost, formation and character of, 172.
 Hurricanes, nature and origin of, 163.
 Hypsometrical zones of vegetation, 183, 184.
 IBERIAN, Hesperian, or Spanish mountain system, 50.
 Icebergs, nature and formation of, 173.
 Igneous agents of change, 216.
 Indian Ocean, area and leading features, 81; currents of, 105.
 Indigenous and exotic, definition of the terms, 183.
 Indo-European race of mankind, 205.
 Indus, the, its tributaries and delta, 128.
 Inorganic and organic, distinction between, 180.
 Iran or Persia, table-land of, 64.
 Irawady, main river of Burmah, 128.
 Isocheimnal or Isocheimonal lines, defined, 179.
 Isothermal lines, explanation of, 179.
 Isothermal lines, nature of, explained 179.
 Isthmus, definition of, 85.

- KAROS** or arid terrace-lands of South Africa, 65.
Khamain or hot wind of Egypt, 161.
Kiölen or **Kölen** Mountains in Scandinavia, 52.
Kuen-lun Mountains, characteristics of, 55.
LAGOONS or **Lagunes** (Ital.), shallow sea lakes, 37.
Lakes and lacustrine areas, 132-138; lakes, their functions, 133; lakes, the principal, 184-188.
Laminarian zone of vegetation, 189; of animals, 195.
Land, its natural subdivisions and terms, 35; its superficial configuration, 40-45; its highlands, 46-66; its lowlands, 67-76.
Land or continental hemisphere, defined, 33.
Land and water, action and reaction of, 39; distribution of, 31-39: relative proportions of, 32.
Langefeld Mountains in Norway, 52.
La Plata, leading features of, 125.
Latitudes, low, high, and middle, meaning of, 21.
Latitude and longitude, definitions of, 21.
Life-stratum, definition of the term, 182.
Littoral (lat. *litus*, the shore) zone of life, 189, 195.
Llanos or grassy plains of the Orinoco, 68, 73.
Lochs, sea-arms or rocky inlets of the sea, 37.
Longitude and latitude, definitions of, 21.
Lowlands or plains of the world, 67-76.
MALAY variety of human race, 209.
Man, as affected by external conditions, 200-202.
Marine fauna, distribution of, 194; flora, 189.
Meridians (Lat. *meridies*, mid-day), defined, 20.
Mesopotamia (Gr. *mesos*, middle; *potamos*, river), plain of, 70.
Metamorphic or crystalline schists, nature of, 28.
Mexican table-land, nature and aspect of, 66.
Missouri, leading features of, 126.
Mississippi, characteristics and tributaries of, 126.
Mists, origin and nature of, 165.
Modifying causes operating on the earth, 214, 216.
Mongolian variety of human race, 206.
Monsoons, origin and nature of, 167.
Month, cause and duration of, 10.
Moon, motions and dimensions of, 10.
Mountains, their characteristics and origin, 46, 49.
Mountains and mountain-systems, 47, 48.
Murray river, of South Australia, 129.
NEOZO variety of human race, 208.
New red sandstone system, character of, 28.
Nile, the, characteristics of, 124; delta of, 68.
OASIS, or islet of vegetation, in the Sahara, 71.
Ocean, the, and its subdivisions, 36.
Ocean, saltness of, 84; density of water, 86; depth, 87; temperature, 89; colour and luminosity, 90; waves, 91; tides, 94; currents, 98-107; areas of, 81.
Ohio, the, chief characteristics of, 126.
Old red sandstone system, nature and composition, 28.
Oolitic system, nature and composition of, 28.
Organic and inorganic nature defined, 180.
Orinoco, the leading characteristics of, 125.
PACIFIC Ocean, area and leading features, 80; currents of, 106.
Palms, zone and regions of, 187.
Pampas or thistly flats of La Plata, 68, 73.
Pampero, or hurricane wind of the Pampas, 162.
Parimé mountain-system, described, 62.
Passes, mountain-passes, and defiles, 47.
Patagonia, desert terrace-lands of, 73.
Peninsula (Lat. *pene*, almost; *insula*, island), 35.
Peninsulas, general southward bearing of, 41.
Permian or lower new red sandstone system, 28.
Physical geography, definition of, 3, 4; theoretical and practical importance of, 6; hints to the learner, 7.
Physiognomy of vegetation in certain areas, 190.
Plains and deserts, characters of, 67-73.
Planetary relations of the earth, 8-21.
Planets and planetoids, 9.
Plants, special distribution of, 184.
Plants and animals, distribution of, 180-199; mutual dependence of, 198.
Plateaux or table-lands, 64-66.
Polar zone of vegetation, definition of, 185.
Polders or embanked meadow-flats of Holland, 69.
Poles, north and south, defined, 18.
Post-tertiary formation, character of, 28.
Prairies (Fr.) or plain-lands of North America, 68, 72.
Profile and contour, illustration of, p. 50.
Puna winds of the Andes, 162.
RAIN, formation and amount of, 166-173.
Rainfall, constant, periodical, and variable, 169-172.
Rainless tracts of the Old and New World, 171.
Regions of vegetable distribution, 190.
Representative species, meaning of, 196.
Review of facts and principles, 210-217.

- Rhine, the, characteristics, 123.
 Rhone, the, leading features of, 123.
 Rivers and river-systems, 115-131; characteristics of, 115-121; technicalities, 115.
 Roads and roadsteads for anchorage, 37.
 Rocky Mountains, members of, 60.
- SAHARA**, the, great arid desert of Africa, 71.
 Salt lakes and inland seas, 135.
 Sand pillars, origin and nature of, 163.
 Sargasso (Span.) or Grassy Sea, 103.
 Sarmatian mountain-system, 52.
 Sarmatian plain, characteristics of, 69.
 Satellites (Lat. *satellites*, an attendant), 9.
 Scandinavian mountain-system, its nature, 52.
 Sea and land breezes, origin of, 159.
 Seaboard, nature and character of, 32.
 Seas, their areas and configurations, 77-83.
 Seasons, nature and cause of, 11.
 Sea-water, its composition and character, 77, 84-90.
 Selvas (Lat. *selva*), wooded flats of the Amazon, 68, 73.
 Shore-line, nature and definition, 32.
 Siberian plain, physical features of, 69; rivers, 122.
 Silurian system, nature and composition of, 28.
 Simoom, or hot wind of the deserts, 161.
 Sirocco, or hot wind of Sicily and Italy, 161.
 Slope and counter-slope, definition of, 44.
 Snow-blink of polar regions, 174.
 Snow-blanket, protecting power of, 174.
 Snow, formation and nature of, 172.
 Snow-line, definition of, 145, 172.
 Solano, or hot south wind of Spain, 161.
 Solar system, members of, 9.
 Springs, origin and characteristics of, 109-112.
 Stannovoi Mountains, North-eastern Asia, 55.
 Steppes, or grassy plains of Northern Asia, 69.
 St Lawrence, the, characteristics of, 126.
 Straths, or ancient river-plains of Scotland, 68.
 Stratified rocks, their origin and character, 24.
 Streams, origin and characteristics of, 113-114.
 Sub-arctic zone of vegetation, 185.
 Sub-tropical zone of vegetation, character of, 185.
- TABLE-LANDS** or plateaux, 64-66.
 Tartary, desert table-lands of, 64.
 Taurus and Anti-Taurus mountains, 53.
 Temperate (warmer and colder) zones of vegetation, 185.
- Temperature of globe, superficial and internal, 16.
 Terrestrial fauna, distribution of, 193; flora, 185.
 Tertiary formation, structure and composition of, 28.
 Thian-shan or Celestial mountain-range, 56.
 Tibet, elevated plateau of, 64.
 Tides, origin, nature, and courses, 94-97.
 Titicaca lake, features of, 137.
 Trade-winds, nature and origin of, 152.
 Trappean rocks, nature and composition of, 29.
 Triassic or upper new red sandstone system, 28.
 Tropical zones of vegetation, character of, 185.
 Tropics, the (Gr. *trōpē*, a turning), defined, 19.
 Tundras, or frozen peat-bogs of Siberia, 68, 69.
 Turan, plain or steppe-land of, 70.
 Typhoons, origin and nature of, 163.
- UNSTRATIFIED** rocks, their origin and character, 24.
 Ural or Uralian mountain-system, 52.
- VALDAI** hills of Central Europe, 52.
 Valleys and minor depressions of the land, 74-76.
 Varieties of human race, 203-206.
 Vertical relief or elevation, 43-45.
 Vine, regions of, described, 187.
 Volcanic and earthquake areas, 216.
 Volcanic rocks, nature and composition of, 29.
 Volcanoes as agents of terrestrial change, 216.
- WATER**, the, its oceans and seas, 77-107; its rivers and lakes, 108.
 Water or oceanic hemisphere, defined, 33.
 Waterspout, nature and cause of, 163.
 Waves, nature and character of, 91, 92.
 Western, Pacific, or Rocky Mountain system, 60.
 Whirlwinds, nature and cause of, 163.
 Winds, constant, periodical, and variable, 150-164.
- YABLONOI** Mountains, North-Eastern Asia, 55.
 Yang-tse-kiang, leading features of, 127.
 Year, cause and duration of, 10.
- ZAMBESI** or Zambesé, great river of Eastern Africa, 129.
 Zones, torrid, temperate, and frigid, 19.
 Zones of vegetation, 185; of animals, 193.

THE END.

PRINTED BY WILLIAM BLACKWOOD AND SONS, EDINBURGH.

EDUCATIONAL WORKS

PUBLISHED BY

WILLIAM BLACKWOOD & SONS.

Fourteenth Edition.

EPITOME OF ALISON'S HISTORY OF EUROPE, for the Use of Schools AND YOUNG PERSONS. Post 8vo, pp. 604, price 7s. 6d. bound in leather.

In compiling this Epitome, it has been specially held in view to omit or suppress no fact of the slightest importance, and to limit the abridgment to the condensation of the minor and accessory details; and it is trusted that an adherence to this rule, while it has produced a work in which the interest of the narrative never flags, has also secured a history of the time in all essential particulars as complete as the more voluminous records of it.

A Chronological Table has been added of all the principal events, so arranged as to give a clear idea of the order in which they succeed each other; and a full Table of Contents, containing a synopsis of the subjects treated of in the body of the work.

Atlas to Epitome of the History of Europe, Eleven Coloured Maps, by A. KEITH JOHNSTON, F.R.S.E. In quarto, price 7s.

Second Edition, post 8vo, with Index, price 9s.

HISTORY OF FRANCE, from the Earliest Times to 1848. By the Rev.

JAMES WHITE, Author of "The Eighteen Christian Centuries."

"This book makes an attempt to furnish a readable account of the country with which we are in closest neighbourhood, and yet of whose history the generality of us know less than of that of almost any other kingdom. It aims at something higher than a mere epitome, for it founds itself on a great deal of various reading, and gives results more than abstracts. At the same time, it devotes sufficient space to any occurrences which seem to have a general bearing on the progress or character of the nation. But it does not profess to be very minute in its record of trifling or unimportant occurrences, nor philosophic in searching out the causes of obscure events."—*Author's Preface.*

"Contains every leading incident worth the telling, and abounds in word-painting, whereof a paragraph has often as much active life in it as one of those inch-square etchings of the great Callot."—*Athenæum.*

Fourth Edition, with Index.

THE EIGHTEEN CHRISTIAN CENTURIES. By the Rev. James White, Author of the "The History of France." In post 8vo, price 7s. 6d.

"He has seized the salient points—indeed, the governing incidents—in each century, and shown their received bearing as well on their own age as on the progress of the world. Vigorously and briefly, often by a single touch, has he marked the traits of leading men; when needful, he touches slightly their biographical career. The state of the country and of society, of arts and learning, and, more than all, of the modes of living, are graphically sketched, and upon the whole with more fulness than any other division."—*Spectator.*

"By far the best historical epitome we have ever perused, and it supplies a great want in this knowing age."—*Atlas.*

FORTIFICATION: FOR OFFICERS OF THE ARMY AND STUDENTS OF MILITARY HISTORY. By Lieut. HENRY YULE, Bengal Engineers. With Illustrations. 8vo, 10s. 6d.

CHOIX DES MEILLEURES SCÈNES DE MOLIÈRE, avec des Notes de Divers Commentateurs, et Autres Notes Explicatives. Par Dr E. DUBUC. Fcap. 8vo, price 4s. 6d.

TEXT-BOOKS ON GEOLOGY,

By DAVID PAGE, F.G.S.

I.

A New Edition, being the **FIRST**,

Enlarged and more fully illustrated, price 1s. 9d., cloth.

INTRODUCTORY TEXT-BOOK OF GEOLOGY. With Glossarial Index.

"Of late it has not often been our good fortune to examine a text-book on science of which we could express an opinion so entirely favourable as we are enabled to do of Mr Page's little work."—*Athenaeum*.

II.

A New Edition, being the **THIRD**,

Enlarged and more fully illustrated, price 6s.

ADVANCED TEXT-BOOK OF GEOLOGY, Descriptive and Industrial. With Glossary of Terms.

"An admirable book on Geology. It is from no invidious desire to underrate other works—it is the simple expression of justice—which causes us to assign to Mr Page's 'Advanced Text-book' the very first place among geological works addressed to students, at least among those which have come before us. We have read every word of it with care and with delight, never hesitating as to its meaning, never detecting the omission of anything useful in a popular and succinct exposition of a rich and varied subject. The *ordonnance* of its material is clear, masterly, and philosophical. The exposition is often eloquent, without ever striving after rhetorical effect. The information is lucidly yet briefly given. . . . The three passages (quoted) sufficiently indicate the quality of the book; its merits as a text-book can only be estimated by the student himself. If it does not drive many a student, hammer in hand, into quarries and railway cuttings, for immediate experience of geological phenomena, nothing will."—*Leader*.

"It is therefore with unfeigned pleasure that we record our appreciation of his 'Advanced Text-Book of Geology.' We have carefully read this truly satisfactory book, and do not hesitate to say that it is an excellent compendium of the great facts of Geology, and written in a truthful and philosophic spirit."—*Edinburgh Philosophical Journal*.

"We know of no introduction containing a larger amount of information in the same space, and which we could more cordially recommend to the geological student."—*Athenaeum*.

III.

Price Sixpence,

THE GEOLOGICAL EXAMINATOR. A Progressive Series of Questions adapted to the Introductory and Advanced Text-Books of Geology. Prepared to assist Teachers in framing their Examinations, and Students in testing their own Progress and Proficiency.

IV.

In Crown Octavo, price 6s.

HANDBOOK OF GEOLOGICAL TERMS AND GEOLOGY.

"Every science, like every manual art, must have its own technicalities and modes of expression: new objects require new names, and new facts new phrases to express their relations. There is no avoiding this necessity in any progressive branch of human knowledge; and the only thing that can be done to lessen the difficulty—next to the rigid exclusion of whatever seems superfluous—is to explain these terms in brief and simple language. This I have endeavoured to do, chiefly with a view to the requirements of the general reader, at the same time appending such details as might render the volume an acceptable Handbook of Reference to the student and professed Geologist. Thus the ordinary reader will generally find the information he requires in the first and second sentences of a definition; what follows is addressed more especially to the professional inquirer—to the student, miner, engineer, architect, agriculturist, and others, who may have occasion to deal with geological facts, and yet who might not be inclined to turn up half-a-dozen volumes, or go through a course of geological readings, for an explanation of the term in question.

"Such is the aim and object of this 'Handbook of Geological Terms.' I lay claim to little more than the arrangement of the matter, which has been gleaned and sifted from many sources—care having always been taken to present the science in its newest aspects, and to express its facts in the clearest and simplest language."—*Author's Preface*.

V.

THE PAST AND PRESENT LIFE OF THE GLOBE. Being a Sketch in

Outline of the World's Life-System. By DAVID PAGE, F.G.S. With Fifty Illustrations, Drawn and Engraved expressly for this Work. In crown octavo, price 6s.

"The object of this excellent and unpretending little work is clearly set forth in the title. It contains a pictorial and historical sketch of the life-scenes of our earth, from the earliest times until now, as inferred from the geological records. It is perspicuous in style, prettily illustrated, furnishing a very complete compendium of the subject. The spirit of the book is good. Mr Page is evidently a philosopher who thinks it no disgrace to be a Christian—a man of science who believes in an Author of science. He is no 'developmentalist;' and his arguments concerning successions of life are singularly acute and conclusive. Some of his positions are strikingly original; as, for instance, his views on recurrent cycles of temperature. He holds creation to be still in progress, presiding over the constant introduction of new species. And as each advance in creation has been, on the whole, upward, so our author anticipates a still higher order of animated creatures on our earth, perhaps also a higher destiny for man. We strongly recommend this little book to our readers, and its calm and temperate spirit to many of our modern writers."—*British Quarterly Review*.

"Mr Page is favourably known as the author of three or four useful elementary geological books, each of which we have noticed in due order of appearance. Together with the one now published, they form a serviceable series for primary instruction."—*Athenæum*.

INTRODUCTORY TEXT-BOOK OF PHYSICAL GEOGRAPHY. By David

PAGE, F.G.S. With Engravings. 1s. 9d.

THE CHEMISTRY OF COMMON LIFE. By Professor Johnston. A New

Edition, edited by G. H. LEWES. Two vols. crown 8vo, with Engravings. Price 11s. 6d.

"Mr Johnston's book is a book for the people; there is hardly a fact or a principle that it would not be for the benefit of the richest as well as the poorest to know."—*Athenæum*.

"All will concur in admiring the profound thought which has ennobled so many familiar things, and has even tinged the commonest processes of household life with the hues of novelty and surprise. The work deserves to be universally read."—*British Quarterly Review*.

"An invaluable reading class-book, and we earnestly recommend it for that purpose."—*English Journal of Education*.

THE PHYSIOLOGY OF COMMON LIFE. By George Henry Lewes. Two

vols. crown octavo, illustrated with numerous Engravings on Wood. Price 12s.

INSTRUCTIONS FOR THE ANALYSIS OF SOILS. By Professor Johnston.

Fourth Edition, crown octavo, 2s.

AINSLIE'S LAND-SURVEYING. A New and Enlarged Edition, embracing

RAILWAY, MILITARY, MARINE, and GEODETICAL SURVEYING. By W. GALBRAITH, M.A. In octavo, with plates in quarto, price 21s.

"The best book on Land-Surveying with which I am acquainted."—WM. RUTHERFORD, LL.D., F.R.A.S., *Royal Military Academy, Woolwich*.

Fifty-Fifth Edition, price 1s.

CATECHISM OF AGRICULTURAL CHEMISTRY AND GEOLOGY. By

Professor JOHNSTON.

"The extent to which this little Catechism has been circulated at home, its translation into nearly every European language, and its introduction into the Schools of Germany, Holland, Flanders, Italy, Sweden, Poland, and South and North America, while it has been gratifying to the Author, has caused him to take additional pains in improving and adding to the amount of useful information in the present edition."—*Preface*.

In crown 8vo, with Illustrations, price 1s.

A CATECHISM OF PRACTICAL AGRICULTURE. By Henry Stephens, F.R.S.E., Author of the "Book of the Farm," &c.

"Teachers will find in this little volume an admirable course of instruction in practical agriculture—that is, the outlines, which they may easily fill up; and by following the hints given in Mr Stephens's preface, the course will scarcely fail to be quite interesting, as well as of great practical benefit. Landed proprietors and farmers might with propriety encourage the introduction of this work into schools, and also have it put into the hands of servants."—*Aberdeen Journal*.

Seventh Edition, fcap. 8vo, price 6s. 6d.

ELEMENTS OF AGRICULTURAL CHEMISTRY AND GEOLOGY. By Professor JOHNSTON.

"Nothing hitherto published has at all equalled it, both as regards true science and sound common-sense."—*Quarterly Journal of Agriculture*.

In post 8vo, price 5s.

ELEMENTARY ARITHMETIC. By Edward Sang, F.R.S.E.

This Treatise is intended to supply the great desideratum of an intellectual instead of a routine course of instructions in Arithmetic.

The foundations of the science are laid with scrupulous care, and are thus able to support a firm and elevated superstructure; so that while the pupil is guided to the acquisition of expertness in performing the operations, his mind is engaged in the study of those principles on which they rest.

In crown 8vo, price 5s.

THE HIGHER ARITHMETIC. By the same Author. Being a Sequel to "Elementary Arithmetic."

"We know, indeed, of no more complete philosophy of pure arithmetic than they contain; they are well worthy of Sir John Leslie's favourite pupil. It is almost needless to add, that we consider the reasoning of these volumes both thorough and close, and the expression of that reasoning uniformly simple and clear."—*Edinburgh Weekly Review*.

"To all who are interested in the science of numbers and its practical uses, we commend Mr Sang's ingenious and elaborate treatise on the Elementary and the Higher Arithmetic."—*Literary Gazette*.

Price Sixpence, for the Waistcoat Pocket.

FIVE PLACE LOGARITHMS. Arranged by E. Sang, F.R.S.E.

SIR WILLIAM HAMILTON'S LECTURES ON METAPHYSICS. Edited

by the Rev. H. L. MANSIE, B.D., LL.D., Waynflete Professor of Moral and Metaphysical Philosophy, Oxford; and JOHN VEITCH, M.A., Professor of Logic, Rhetoric, and Metaphysics, St Andrews. Second Edition. In 2 vols. octavo, price 24s.

SIR WILLIAM HAMILTON'S LECTURES ON LOGIC. Edited by Professors

MANSIE and VEITCH. In 2 vols. octavo, price 24s.

INSTITUTES OF METAPHYSIC. The Theory of Knowing and Being.

By JAMES F. FERRIER, B.A. Oxon., Professor of Moral Philosophy and Political Economy, St Andrews. Second Edition. Crown octavo, price 10s. 6d.

DESCARTES ON THE METHOD OF RIGHTLY CONDUCTING THE

REASON, and Seeking Truth in the Sciences. Translated from the French. 12mo, 2s.

DESCARTES' MEDITATIONS, AND SELECTIONS FROM HIS PRIN-

CIPLES OF PHILOSOPHY. Translated from the Latin. 12mo, 3s.

NEW GENERAL ATLAS.

Dedicated by Special Permission to Her Majesty.

THE ROYAL ATLAS

OF

MODERN GEOGRAPHY.

IN A SERIES OF ENTIRELY ORIGINAL AND AUTHENTIC MAPS.

By A. KEITH JOHNSTON,

F.R.S.E. F.R.G.S.; Author of the "Physical Atlas," &c.

With a complete Index of easy reference to each Map, comprising nearly
150,000 Places contained in this Atlas.

In Imperial Folio, hf.-bd. in russia or morocco, £5, 15s. 6d.

**The only ATLAS for which a PRIZE MEDAL was awarded at the
International Exhibition, 1862.**

FROM THE "TIMES," DEC. 27, 1861.

NO one can look through Mr Keith Johnston's new Atlas without seeing that it is the best which has ever been published in this country. To those who have a mean idea of our British map-makers perhaps this is not saying much. Let us therefore add that the new Atlas takes away from us a reproach, and is worthy of a place beside the best of the Continental ones. It is a fit successor to that other production of Mr Johnston's, the *Physical Atlas*, which is quite unique among such works for beauty of execution and richness of information. In the present volume we have, at a charge of five and a half guineas, about 50 most carefully prepared and highly finished maps, together with a special index to each, containing all the names that appear in it. The price, therefore, of these remarkably accurate and beautiful maps is no more than 2s. for each—a fabulous sum, if we consider the amount of work which they contain. "The plan of this work," says Mr Johnston, "was deliberately considered years before it was commenced. Arrangements were carefully made for rendering available an immense store of the richest materials in maps, charts, and books, collected at great labour and expense, and the most approved method of constructing, drawing, engraving, and printing the maps was adopted. Nearly five years of constant labour have been devoted to its preparation and production, but it embodies the results of a systematic study of practical geography in its many departments, extending over a quarter of a century." The work thus contains the latest information presented to us in the most attractive form, every new appliance in cartography being turned to the best account.

It is a small matter, but it will indicate the numberless little touches by which Mr Keith Johnston has added to the usefulness of his Atlas, if we say that on

opening the volume we have the pleasure of seeing the maps numbered on the back as well as on the front. In the ordinary atlases, when we open the volume and see two blank pages before us, we know not where we are, and have to turn over the page in order to find out. Here, at whatever page we open, we see a number before us, and are therefore spared a little trouble. The next point we observe is a peculiar use of light blue ink on all the maps. Wherever water is to be represented—either the sea-coast, or a river, or a lake—the water-line is printed in blue ink; and not only is the configuration of this colour, but so also are the accompanying names. The result is certainly most satisfactory. Much confusion is got rid of, and the outlines of land and water stand out clear and beautiful to the eye. At a glance we can see the whole watercourse of a country; and are saved from the bewildering crowdedness of the old style of map. As we proceed in our examination we find that Mr Keith Johnston, for the easier discovery of places, has lettered the sides of each map, in correspondence with an index, so that by a reference to the letters on two adjacent sides we can see on which of the squares formed by the lines of latitude and longitude any name is to be found. This is easier than giving the latitude and longitude in the index.

There is not much to be done with the first of the maps in any general atlas—namely, that of the World in Hemispheres. Let us, therefore, turn to the much more improvable chart of the World on Mercator's Projection. On this Mr Johnston has put forth all his skill, and he has succeeded in producing something very valuable. It is especially valuable for Englishmen that the map is so arranged as to include the British Islands twice over. They appear at the extreme east and the extreme west of the map, so that, looking east or west, we can at once see the relations of England to the other parts of the globe. Over and above this, we cannot help noticing the part which the ocean plays in Mr Johnston's maps. According to the old system the sea is a blank, except where it is relieved by islands. Here all the great currents and streams are displayed, and the routes of vessels from port to port are elaborately indicated. Still more is this the case in a valuable map of the Basin of the North Atlantic Ocean. We have here the entire seaboard of North America, with that of the northern portion of South America, for one side of the Atlantic; and we have the seaboard of Western Europe, as well as of Northern Africa, for the other. Between these we have the entire island system of the North Atlantic, the Gulf Stream, and all the other currents of the ocean, together with the routes traversed by every class of vessel on the voyage across the Atlantic. And here we are reminded that this is one of a class of maps which it is difficult to get. It is unusual to give it in the ordinary atlases. Equally unexpected in a collection of this kind is that most valuable map, the Basin of the Mediterranean, showing at a glance all the shores of this sea, and their relation to those of the Black Sea. To show how completely every detail is worked out, we may state, that in the corners and sides of this map there are inserted seven elaborate plans of different places—a plan of the port and roads of Marseilles, one of the port and gulf of Genoa and of the Straits of Gibraltar, one of the Maltese Islands, one of the harbour and town of Valetta, one of the lagoons of Venice, and one of the port of Alexandria. Nearly all the maps, indeed, have these supplementary plans. Thus, that of France has plans of the city and suburbs of Paris, of the island of Corsica, and of France as divided into provinces before the Revolution. So the map of Italy has plans of Rome and of Naples, together with their environs; and a most elaborate map of the Basin of the Baltic has a plan of St Petersburg and its suburbs. We find the same carefulness in every map. All our colonial settlements are given with remarkable fulness. With regard to the Map of India, perhaps there may be some difference of opinion, not indeed as to the excellence of the chart, but as to the spelling of the names. It is at first puzzling to see the Mysore spelt "Maisur," and Lucknow "Lakhnanu." Mr Johnston says that this vexed question has caused him much anxiety, and that he has simply adopted the scientific method of spelling introduced by Sir William Jones, and used by the Asiatic and other learned societies in the *Trigonometrical Survey*, as well as by the author of Murray's *Handbook for India*. At all events, he has done his best to prevent any confusion by printing the received spelling of the name in conjunction with the scientific orthography. Perhaps he might have made an exception in favour of certain well-known names, but we are not disposed to speak of his omission to do so as an unpardonable crime. The advance of this Atlas upon its predecessors is so marked that it is hypercritical to dwell upon a point of spelling which any one who will use the map can in a few minutes master.

OPINIONS OF THE PRESS.

Athenæum, August 10, 1861.

Under the name of "The Royal Atlas of Modern Geography," Messrs Blackwood and Sons have published a book of maps, which for care of drawing and beauty of execution appears to leave nothing more to hope for or desire. Science and art have done their best upon this magnificent book. Mr A. Keith Johnston answers for the engraving and printing: to those who love clear forms and delicate bold type we need say no more. All that maps should be, these maps are: honest, accurate, intelligible guides to narrative or description. A very good feature is the Index of Names attached to each plate, with the easy mechanical arrangement of lines by which any particular name that may be sought is at once found on the map. Of the many noble atlases prepared by Mr Johnston and published by Messrs Blackwood and Sons, this Royal Atlas will be the most useful to the public, and will deserve to be the most popular.

Morning Herald.

There is scarce any science which has of late made such progress as that of chartography. But the culmination of all attempts to depict the face of the world appears in the "Royal Atlas," than which it is impossible to conceive anything more perfect. The only way to test the value of maps is to select those spots which one thoroughly knows, and from them to judge of the remainder. Never did an atlas stand this searching examination so well as the folio work just published by the Messrs Blackwood, and bearing on its title the honoured name of Keith Johnston. We have carefully gone over the coast-lines and the interior delineation of several portions of Europe, and of North and South America, with which we happen to be familiar, and, so far as we have examined, there is not merely not an error, but the absolute perfection of accuracy. . . . The most magnificent geographical work that has ever issued from the press. Considering the labour expended on it, and the style in which it is got up, its cheapness is not its greatest marvel.

Guardian.

This is, beyond question, the most splendid and luxurious, as well as the most useful and complete, of all existing atlases. To a habitual consulter of maps (which every reader of history and every man of science ought to be) there can hardly be a greater luxury than these beautiful sheets, which it is a pleasure for the eye to travel over, and upon which a vast quantity of the finest and most delicate work represents a more than corresponding mass of various information. In one large but not cumbrous folio volume Mr Keith Johnston has given us as perfect an atlas as can well be desired, embracing the results of all the most recent discoveries in every part of the world, as well as most of the territorial changes that have lately occurred in Europe. . . . A close examination of several different parts of Europe which are very familiar to us, satisfies us of the remarkable accuracy with which the work is done; it is very rarely that we miss even a new road, where, according to the scale of the map, a road should have been given.

OPINIONS OF THE PRESS.

Saturday Review.

The completion of Mr Keith Johnston's *Royal Atlas of Modern Geography* claims a special notice at our hands. While Mr Johnston's maps are certainly unsurpassed by any for legibility and uniformity of drawing, as well as for accuracy and judicious selection, this eminent geographer's Atlas has a distinguishing merit in the fact that each map is accompanied by a special index of remarkable fullness. The labour and trouble of reference are in this way reduced to a minimum. The number of places enumerated in the separate indices is enormous. We believe indeed that every name which appears in the maps is registered in the tables; and as each place is indicated by two letters, which refer to the squares formed by the parallels of latitude and longitude, the method of using the index is extremely easy and convenient. We know no series of maps which we can more warmly recommend. The accuracy, wherever we have attempted to put it to the test, is really astonishing.

Examiner.

There has not, we believe, been produced for general public use a body of maps equal in beauty and completeness to the Royal Atlas just issued by Mr A. K. Johnston. In beauty and clearness of engraving it has never been surpassed, and the fullness of information in such maps as the five representing the United Kingdom (England and Scotland each being enlarged so as to fill two maps, upon which the name of every hamlet has been entered), or the two representing India, is marvellous. We have a mechanical perfection of name-engraving in the crowded map of the North of England and the Southern sheet for Scotland; the last named being an especial labour of love, which includes even the footpaths, records sites of battles, and notes minutely many physical features of the country, is one of which a Scotchman may be proud. Something of beauty as well as much clearness is given to the engraving by the use of blue ink instead of black for the name of every lake, river, canal, harbour, bay, or other form of water. There are also ten or twelve forms and sizes of letters used to express the character and relative importance of the places named.

Globe.

This is one of those elegant and complete works which have given Mr Johnston his high reputation. The maps of this Atlas were originally published in parts, and at the time we expressed our high opinion of the plan and execution; we have now only to repeat that opinion. The beauty of the work equals its accuracy. Exquisitely engraved, coloured with clearness and decision, roads, rivers, and mountains distinctly marked, all the maps are grateful to the eye, and some are really beautiful pictures; now they are all bound together, they form one of the handsomest as well as one of the most useful volumes in the shelves of a library. The maps are "up to date" in point of modern discovery. They are original, and have been compiled from authentic sources, generously placed at Mr Johnston's disposal, not only by our own but by many foreign Governments.

OPINIONS OF THE PRESS.

From *Nouvelles Annales des Voyages*, &c., January 1861.

Ainsi disposé, l'Atlas Royal de Keith Johnston, dont S. M. la Reine Victoria a daigné accepter l'hommage, offre un répertoire des plus complets de l'état de la science géographique à notre époque. Il nous a paru digne du haut patronage sous lequel il est placé, et nous pensons qu'il est destiné à rendre d'utiles services aux savants comme aux gens du monde.

V. A. MALTE-BRUN.

Scotsman.

To go over the maps *seriatim*, noticing additions and improvements, would be tedious and somewhat unprofitable. An almost daily reference to, and comparison of it with others, since the publication of the first part some two years ago until now, enables us to say, without the slightest hesitation, that this is by far the most complete and authentic atlas that has yet been issued. In the whole of that time we failed to detect more than two or three errors, and these were so very trifling that we have forgotten what they were.

Bulletin de la Société de Géographie.

Mais entre les grande atlas, je dois vous signaler de préférence l'Atlas Royal de Géographie Moderne publié par le savant auteur de l'Atlas Physique, M. Alexandre Keith Johnston. Exécutées par une personne qui s'est tenue au courant des découvertes dont s'est enrichie la connaissance du globe, et des changements dans la circonscription des états.

From the Address of Sir E. I. Murchison, President of the
Royal Geographical Society.

Having called your attention in days gone by to the improvements made in cartography by Mr A. Keith Johnston, and to his zealous and successful endeavours to lay before his countrymen on maps all the chief data of Physical Science, I have now the satisfaction of adverting to his last important work—a new General Atlas. Fifteen years have elapsed since he published his National Atlas: the author felt that the time was come for the production of an entirely new work, which would embrace all the recent discoveries and all the territorial changes. In accomplishing his task, Mr Johnston has thoroughly succeeded in placing before the public a series of sheets in each region on a very convenient scale, and also by a judicious selection of names, arranged on a special index accompanying each map, which at once directs the observer to the position of any place. But that which most pleases my eye and instructs me, is the remarkable distinctness which is given to every water-course, lake, canal, or railroad by the use of light blue ink; by this process the orography and skeleton of every country stand out in clear relief, the outlines being never intermingled or confusing the eye as in old maps. In short, this beautiful Atlas will I have no doubt be generally approved, and its sale will I trust reward the author for his long and arduous labours.

CONTENTS.

Plate 1.	THE WORLD (in HEMISPHERES),	With Index to 1400 Places.
2.	CHART OF THE WORLD ON MERCATOR'S PROJECTION,	1340 "
3.	EUROPE,	2120 "
4.	BASIN OF THE NORTH ATLANTIC OCEAN,	1010 "
*5, 6.	ENGLAND (Two Sheets),	10,600 "
*7, 8.	SCOTLAND (Two Sheets),	9250 "
*9.	IRELAND,	5270 "
10.	FRANCE, IN DEPARTMENTS,	4406 "
*11.	SPAIN AND PORTUGAL,	4100 "
*12.	BASIN OF MEDITERRANEAN SEA,	2170 "
*13, 14.	NORTHERN ITALY AND SOUTHERN ITALY (Two Sheets),	6230 "
*15.	SWITZERLAND, and the ALPS of SAVOY and PIEDMONT,	4907 "
*16.	BELGIUM, } One {	2100 "
	THE NETHERLANDS, } Sheet {	2200 "
	DENMARK and ICELAND,	2000 "
17.	HANOVER, BRUNSWICK, MECK- LENBURG, and OLDENBURG, } One Sheet {	1180 "
*18.	SOUTH-WESTERN GERMANY,	4470 "
*19.	PRUSSIA,	2550 "
*20, 21.	AUSTRIAN EMPIRE (Two Sheets),	6300 "
	TURKEY IN EUROPE,	2280 "
23.	GREECE and the IONIAN ISLANDS,	2187 "
*24.	SWEDEN and NORWAY (SCANDINAVIA),	1630 "
25.	BASIN OF THE BALTIC SEA,	1830 "
26.	EUROPEAN RUSSIA,	3070 "
27.	SOUTH-WEST RUSSIA,	3740 "
28.	ASIA,	3900 "
29.	TURKEY IN ASIA (ASIA MINOR) and TRANSCAUCASIA,	2850 "
*30.	PALESTINE,	3100 "
31.	PERSIA and AFGHANISTAN,	2150 "
*32, 33.	INDIA (Two Sheets),	7500 "
*34.	CHINA and JAPAN,	2420 "
35.	OCEANIA,	2500 "
*36.	SOUTH AUSTRALIA, NEW SOUTH WALES, and VICTORIA,	1980 "
37.	AFRICA,	3850 "
38.	NORTH-WESTERN AFRICA, } One {	1340 "
	SOUTHERN AFRICA, } Sheet {	1250 "
39, 40.	EGYPT, NUBIA, ABYSSINIA, and ARABIA PETRÆA (Two Sheets),	2840 "
41.	NORTH AMERICA,	2740 "
*42, 43.	CANADA, NEW BRUNSWICK, NOVA SCOTIA, and NEWFOUNDLAND (Two Sheets),	3070 "
*44, 45.	UNITED STATES OF NORTH AMERICA (Two Sheets),	5675 "
46.	WEST INDIES and CENTRAL AMERICA,	1170 "
*47, 48.	SOUTH AMERICA,	5400 "

Each Plate may be had separately, with its Index, price 3s. ; and those marked
thus * mounted on linen in cloth cases for the pocket, if in One Sheet, 4s. 6d. ;
in Two Sheets, 8s.

GEOGRAPHICAL WORKS

BY

ALEX. KEITH JOHNSTON,

F.R.S.E. F.R.G.S.

Geographer to the Queen for Scotland.

THE PHYSICAL ATLAS

OF

NATURAL PHENOMENA.

A NEW AND ENLARGED EDITION.

Consisting of 35 large and 7 small Plates, printed in Colours; and 145 folio Pages of Letterpress, including an Index containing upwards of 16,000 References.

In Imperial Folio, hf.-bd. in russia or morocco, £8, 8s.

THE object of the Physical Atlas is to explain, in a popular and attractive manner, the results of the discoveries and researches of travellers and philosophers in all the important branches of Natural Science, embracing an investigation into the laws which regulate the formation of the surface of the globe, the successive changes which this surface has undergone, the varied influences of climate and temperature to which it is subjected, the effects of these several conditions on the existing species of plants and animals, as exemplified in their geographical distribution, and the combined action of the whole on the present races, national characteristics, means of subsistence, commercial pursuits, religious peculiarities, health and progressive improvement of MAN. And while the facts it enunciates are deduced from inquiries conducted in accordance with the soundest principles of inductive reasoning, they are, by a novel application of the principles of chartographic delineation, aided by colours, signs, and diagrams, so simply and forcibly conveyed through the eye to the mind, as to be comprehended with ease and recalled without effort. The book, therefore, has this great recommendation to the general reader, that it may be studied with advantage without any previous training; and as each of the branches into which it is divided, while elucidating the whole, is complete in itself, it may be overtaken in consecutive portions according to taste or requirement; while the complete Index, *now added to this edition*, will render it instantly available for reference on any of the multifarious subjects on which it treats.

THE PHYSICAL ATLAS

LIST OF PLATES.

GEOLOGY AND OROGRAPHY.

MAP

- 1.—THE GEOLOGICAL STRUCTURE OF THE GLOBE. Price 15s.
- 2.—THE PHYSICAL FEATURES OF EUROPE AND ASIA. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 3.—THE MOUNTAIN SYSTEMS OF EUROPE. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 4.—GEOLOGICAL MAP OF EUROPE. By Sir RODERICK IMPEY MURCHISON, D.C.L., F.R.S., &c. &c.; and JAMES NICOL, F.R.S.E., &c. Price 10s. 6d.
- 5, 6.—GEOLOGICAL AND PALÆONTOLOGICAL MAP OF THE BRITISH ISLANDS. In Two Sheets. By Professor EDWARD FORBES. Price 15s.
- 7.—THE PHYSICAL FEATURES OF NORTH AND SOUTH AMERICA. By A. K. JOHNSTON, F.R.S.E., &c.; and Professor H. D. ROGERS. Price 7s. 6d.
- 8.—GEOLOGICAL MAP OF THE UNITED STATES AND BRITISH NORTH AMERICA. By Professor H. D. ROGERS, Boston, U.S. Price 10s. 6d.
- 9.—ILLUSTRATIONS OF THE GLACIER SYSTEMS OF THE ALPS. By A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.
- 10.—THE PHENOMENA OF VOLCANIC ACTION. By A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.
- 11.—COMPARATIVE VIEWS OF REMARKABLE GEOLOGICAL PHENOMENA. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.

HYDROGRAPHY.

- 12.—THE PHYSICAL CHART OF THE ATLANTIC OCEAN. By A. K. JOHNSTON, F.R.S.E., &c.; and Professor H. D. ROGERS. Price 15s.
- 13.—PHYSICAL CHART OF THE INDIAN OCEAN. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 14.—PHYSICAL CHART OF THE PACIFIC OCEAN. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 15.—TIDAL CHART OF THE BRITISH SEAS. By J. SCOTT RUSSELL, and A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 16.—THE RIVER SYSTEMS OF EUROPE AND ASIA. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 17.—THE RIVER SYSTEMS OF NORTH AND SOUTH AMERICA. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.

METEOROLOGY AND MAGNETISM.

- 18.—DISTRIBUTION OF HEAT OVER THE GLOBE. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 19.—GEOGRAPHICAL DISTRIBUTION OF THE CURRENTS OF AIR. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 20.—HYETOGRAPHIC OR RAIN-MAP OF THE WORLD. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 21.—HYETOGRAPHIC OR RAIN-MAP OF EUROPE. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 22.—MAP OF LINES OF EQUAL POLARISATION IN THE ATMOSPHERE. By Sir DAVID BREWSTER, K.H., F.R.S., &c. &c. Price 7s. 6d.
- 23.—TERRESTRIAL MAGNETISM. By Colonel SABINE, R.A., V.P.R.S., &c.

THE PHYSICAL ATLAS.

BOTANICAL GEOGRAPHY.

MAP

- 24.—GEOGRAPHICAL DISTRIBUTION OF THE MOST IMPORTANT PLANTS YIELDING FOOD. By ARTHUR HENFREY, F.R.S., &c. Price 7s. 6d.
- 25.—GEOGRAPHICAL DISTRIBUTION OF INDIGENOUS VEGETATION. By ARTHUR HENFREY, F.R.S., &c., and A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.

ZOOLOGICAL GEOGRAPHY.

- 26.—GEOGRAPHICAL DISTRIBUTION OF MAMMALIA OF THE ORDERS QUADRUMANA, EDENTATA, MARSUPIALIA, AND PACHYDERMATA. By A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.
- 27.—GEOGRAPHICAL DISTRIBUTION OF MAMMALIA OF THE ORDER OF CARNIVORA. By A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.
- 28.—GEOGRAPHICAL DIVISION AND DISTRIBUTION OF MAMMALIA OF THE ORDERS RODENTIA AND RUMINANTIA. By A. K. JOHNSTON, F.R.S.E., &c., and G. R. WATERHOUSE, Esq. Price 10s. 6d.
- 29.—GEOGRAPHICAL DIVISION AND DISTRIBUTION OF BIRDS. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 30.—GEOGRAPHICAL DIVISION AND DISTRIBUTION OF REPTILIA. By A. K. JOHNSTON, F.R.S.E., &c. Price 7s. 6d.
- 31.—THE DISTRIBUTION OF MARINE LIFE. By EDWARD FORBES, F.R.S. Price 10s. 6d.

ETHNOLOGY AND STATISTICS.

- 32.—ETHNOGRAPHIC MAP OF EUROPE. By Dr GUSTAF KOMBST and A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.
- 33.—ETHNOGRAPHIC MAP OF GREAT BRITAIN AND IRELAND. By Dr GUSTAF KOMBST and A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.
- 34.—MORAL AND STATISTICAL CHART. By A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.
- 35.—THE GEOGRAPHICAL DISTRIBUTION OF HEALTH AND DISEASE IN CONNECTION CHIEFLY WITH NATURAL PHENOMENA. By A. K. JOHNSTON, F.R.S.E., &c. Price 10s. 6d.

EXTRACTS FROM REVIEWS, &c.

Sir John Herschell.

The "Physical Atlas" of Mr Keith Johnston—a perfect treasure of compressed information.

Examiner.

There is no map in this noble Atlas upon which we might not be tempted to write largely. Almost every one suggests a volume of reflection, and suggests it by presenting, in a few hours, accurate truths which it would be the labour of a volume to enforce in words, and by imprinting them, at the same time, upon the memory with such distinctness that their outlines are not likely afterwards to be effaced. The "Physical Atlas" is a somewhat costly work, reckoning it only by its paper; but upon its paper is stamped an amount of knowledge that could scarcely be acquired without the reading of as many books as would cost seven times the price.

THE PHYSICAL ATLAS.

Edinburgh Review.

The more extensive and elaborate "Physical Atlas of Natural Phenomena," by Mr A. K. Johnston, of which it would be difficult to speak in terms above the mark of its actual merits, embraces every part of the subject. It delineates to the eye as well as the mind, and far better than by any verbal description, those complex relations of Physical Phenomena on the globe which are the true foundations of Physical Geography.

Quarterly Review.

We know no work containing such copious and exact information as to all the physical circumstances of the earth on which we live, nor any of which the methods are so well fitted for the instruction of those who come ignorantly to the subject.

Bulletin de la Société de Géographie.

Là se termine cet immense travail. Nous avons indiqué rapidement les diverses parties de l'Atlas; nous devons ajouter que chaque carte est accompagnée d'un texte explicatif, qui fait connaître l'état présent de la science. De telles publications font la gloire d'un pays, et nous voyons avec un profond sentiment de regret la France devancée par l'Allemagne et l'Angleterre dans cette voie si belle et si féconde.

Proceedings of Royal Geographical Society.

Our Associate, Mr Alexander Keith Johnston, has completed the new edition of his superb "Physical Atlas." The publication of the first edition of this great work, some ten years since, had the effect of introducing into this country almost a new era in the popular study of Geography, through its attractive and instructive illustration of the prominent features of the science. This second edition is to some extent entirely a new work, owing to the additions and improvements which have been introduced. I have only to refer to the names of Murchison, Forbes, Brewster, Ami Boué, and Berghaus, to stamp the high character of the work; but I must not omit to mention, among new contributions, the Geological Map of Europe by Sir Roderick Murchison and Professor Nicol, that of America by Professor Rogers, Colonel Sabine's Map of Terrestrial Magnetism, the Distribution of Marine Animals by the lamented Professor Edward Forbes, and the addition of a large General Index adds materially to the utility of this extensive Compendium of Natural Geography.

THE PHYSICAL ATLAS.

FOR THE USE OF COLLEGES, ACADEMIES, AND FAMILIES.

Reduced from the Imperial Folio.

BY A. KEITH JOHNSTON, F.R.S.E. &c.

This Edition contains 25 Maps, including a Palæontological Map of the British Islands, with Descriptive Letterpress, and a very copious Index. In Imperial Quarto, half-bound morocco, £2, 12s. 6d.

Four Sheets Imperial, beautifully printed in Colours,
A GEOLOGICAL MAP OF EUROPE,

EXHIBITING THE DIFFERENT SYSTEMS OF ROCKS, ACCORDING TO THE
LATEST RESEARCHES, AND FROM INEDITED MATERIALS.

By **SIR R. I. MURCHISON, D.C.L. F.R.S. &c.**

Director-General of the Geological Survey of Great Britain and Ireland;

And **JAMES NICOL, F.R.S.E. F.G.S.**

Professor of Natural History in the University of Aberdeen.

Constructed by **ALEX. KEITH JOHNSTON, F.R.S.E. &c.**

Geographer to the Queen, Author of the "Physical Atlas," &c.

Scale, $\frac{1}{4,800,000}$ of nature, 76 miles to 1 inch.

Size, 4 feet 2 by 3 feet 5 inches. Price, in Sheets, £3, 3s.;
in a Cloth Case, 4to, £3, 10s.

**FIRST SKETCH OF A NEW GEOLOGICAL
MAP OF SCOTLAND.**

By **SIR RODERICK I. MURCHISON, D.C.L. &c.;**

And **ARCHIBALD GEIKIE, F.G.S.**

Constructed by **A. KEITH JOHNSTON.**

Price 5s.

GEOLOGICAL MAP OF SCOTLAND.

FROM THE MOST RECENT AUTHORITIES AND PERSONAL OBSERVATIONS.

By **JAMES NICOL, F.R.S.E. &c.**

Professor of Natural History in the University of Aberdeen.

With Explanatory Notes. The Topography by **ALEXANDER KEITH
JOHNSTON, F.R.S.E., &c.** Scale, 10 miles to an inch.

In Cloth Case, price 21s.

A SMALL GEOLOGICAL MAP OF EUROPE.

FROM KEITH JOHNSTON'S SCHOOL "PHYSICAL ATLAS."

Printed in Colours, price Sixpence.

A GEOLOGICAL MAP OF THE BRITISH ISLES.

FROM THE SAME.

Printed in Colours, price Sixpence.

New and Enlarged Edition, Imperial 8vo, half-bound, price 12s. 6d.

SCHOOL ATLAS OF GENERAL AND DESCRIPTIVE GEOGRAPHY,

EXHIBITING THE ACTUAL AND COMPARATIVE EXTENT OF ALL THE COUNTRIES
IN THE WORLD, WITH THEIR PRESENT POLITICAL DIVISIONS.

Constructed with a special view to the purposes of sound instruction.

CONTENTS. •

Map

1. The World in Hemispheres, with
Tables of the Heights of Moun-
tains and Lengths of Rivers.
2. Europe.
3. England and Wales.
4. Scotland.
5. Ireland.
6. Belgium and the Netherlands.
7. Denmark.
8. France (in Provinces and Depart-
ments).
9. Switzerland.
10. Western Germany and Prussia.
11. Austrian Empire.
12. Italy.
13. Spain and Portugal.

Map

14. Turkey and Greece.
15. Sweden and Norway.
16. Russia and Poland.
17. Asia.
18. India, Siam, and the Malay Penin-
sula.
19. Oceania (Australasia, Polynesia,
&c.)
20. Turkey in Asia, Persia, &c.
21. Canaan or Palestine.
22. Africa and Arabia.
23. North America.
24. United States and Canada.
25. Central America and West India
Islands.
26. South America.

New and Enlarged Edition, Imperial 8vo, half-bound, price 12s. 6d.

SCHOOL ATLAS OF PHYSICAL GEOGRAPHY.

In this Atlas of Physical Geography the subject is treated in a more simple and elementary manner than in the previous works of the Author—the object being to convey broad and general ideas on the form and structure of our Planet, and the principal phenomena affecting its outer crust. Printed in colours.

CONTENTS.

Plate

1. Illustrations of Chartography and Climatography.
2. Geological Map of Europe.
3. of the British Isles.
4. The Mountains, Table-Lands, Plains, and Valleys of Europe.
5. of Asia.
6. of Africa.
7. of N. America.
8. of S. America.
9. Distribution of Earthquakes and Volcanoes over the Globe.
10. Chart of the Ocean-Currents of the World.
11. The Principal Lakes of the Globe, showing their comparative Sizes,
Depth, &c.
12. Chart of the River Systems of the World.
13. Climatological Chart, showing the Temperature of the Globe.
14. Map of the Constant and Periodical Winds of the Globe.
15. The Distribution of Rain and Snow over the Globe.
16. Map of the Distribution of Vegetable Life over the Globe.
17. of Animal Life
18. of the Different Races of Man.
19. Map showing the Distribution and Extent of the Different Forms of
Religion over the Globe.

New and Enlarged Edition, Imperial 8vo, half-bound, 12s. 6d.
SCHOOL ATLAS OF CLASSICAL GEOGRAPHY.

COMPRISING, IN TWENTY PLATES, MAPS AND PLANS OF ALL THE IMPORTANT COUNTRIES AND LOCALITIES REFERRED TO BY CLASSICAL AUTHORS.

Constructed from the best Materials, and embodying the Results of the most recent Investigations. With an Index.

Map**CONTENTS.**

1. Plan of Rome, and Illustrations of Classical Sites.
2. Orbis Terrarum (et Orb. Homeri, Herodoti, Democriti, Strabonis, Ptolemæi).
3. Hispania.
4. Gallia.
5. Insulæ Britannicæ (et Brit. Strabonis, Brit. Ptolemæi, &c.)
6. Germania, Vindelicia, Rætia, et Noricum.
7. Pannonia, Dacia, Illyricum, Moesia, Macedonia, et Thracia.
8. Italia Superior et Corsica.
9. Italia Inferior, Sicilia, et Sardinia (et Campaniæ, Syracusæ, Roma).
10. Imperium Romanum (et Imp. Rom. Orient. et Occid.)
11. Græcia (et Athenæ, Marathon, Thermopylæ).
12. Græcia a Bello Peloponnesiaco, usque ad Philipp. II. (et Mantinea Leuctra, Platea).
13. Asia Minor (et Campus Trojæ, Bosphoros, Troas Ionia).
14. Syria et Palestina (et Hierosolyma, &c.)
15. Armenia, Mesopotamia, Babylonia, Assyria (et Iter Xenophontis).
16. Regnum Alexandri Magni (et Granicus, Issus, Arbela).
17. Persia et India (et India Ptolemæi).
18. Ægyptus, Arabia, et Æthiopia (et Ægyptus Inferior).
19. Africa (et Carthago, Alexandria, Numidia, et Africa Propria).
20. Europe, showing the general direction of the Barbarian Inroads during the Decline and Fall of the Roman Empire.

Index.

In Imperial 8vo, half-bound, price 12s. 6d.

SCHOOL ATLAS OF ASTRONOMY.

Edited by J. R. HIND, Esq., F.R.A.S., &c.

With Notes and Descriptive Letterpress to each Plate, embodying all recent Discoveries in Astronomy.

Eighteen Maps, printed in colours, by a new process.

Map**CONTENTS.**

1. The Celestial Sphere—Refraction—Parallax—Aberration of Light—Phases of the Moon—of the Inferior Planets, and of Saturn's Ring.
2. Axial Rotation of the Earth—Day and Night—Her Annual Revolution in the Ecliptic—The Seasons—The Tides.
3. The Solar Spots—Rotation of the Sun—His apparent magnitude from the various Planets—The Zodiacal Light.
4. Telescopic appearance of the Moon.
5. Eclipses of the Sun, and Phenomena attending them.
6. Eclipses of the Moon.
7. The Solar or Planetary System.
8. Position of the Earth in the Solar System.
9. Transits of Mercury—Telescopic appearances of the Planets—Their relative dimensions.
10. The Comet.
11. Double Stars—Binary Systems—Coloured Stars—Clusters—Distribution of the Stars—The Via Lactea.
12. Nebula.
- 13 to 18. Maps of the Stars.

*Separate Maps of the General, Classical, and Physical Atlases, price 6d.:
 of the Astronomical, price 8d. each.*

New and Cheaper Edition, demy 8vo, price 5s., half-bound,
**ELEMENTARY SCHOOL ATLAS OF GENERAL
 AND DESCRIPTIVE GEOGRAPHY.**

FOR THE USE OF JUNIOR CLASSES.

Including a Map of Canaan and Palestine, and a General Index.

CONTENTS.

Map	Map
1. The World in Hemispheres, with Tables of the Heights of Moun- tains and Lengths of Rivers.	10. Spain and Portugal.
2. Europe.	11. Ottoman Empire in Europe and Asia.
3. England and Wales.	12. Sweden and Norway.
4. Scotland and Ireland.	13. Russia and Poland.
5. France (in Provinces and Depart- ments) and Switzerland.	14. Asia and East India Islands.
6. Belgium, Netherlands, and Den- mark.	15. India—Australia.
7. Prussia and Western Germany.	16. Africa and Arabia.
8. Austria.	17. North and South America.
9. Italy.	18. United States and Canada.
	19. Central America and West India Islands.
	20. Canaan and Palestine.

TWELVE OUTLINE PROJECTIONS, corresponding in
 Scale with the Plates of KEITH JOHNSTON'S General Atlas—viz.,
 Europe, Asia, Africa, North America, South America, England,
 Scotland, Ireland, France, Spain, Italy, and Palestine. Price 2s. 6d.
 Single Projections, 8d. each.

OPINIONS REGARDING THE SCHOOL ATLASES.

"They are as superior to all School Atlases within our knowledge, as were the
 larger works of the same Author in advance of those that preceded them."—*Edu-
 cational Times*.

"Decidedly the best School Atlases we have ever seen."—*English Journal of
 Education*.

"The *Physical Atlas* seems to us particularly well executed. . . .
 The last generation had no such help to learning as is afforded in these excellent
 elementary maps. The *Classical Atlas* is a great improvement on what has usu-
 ally gone by that name; not only is it fuller, but in some cases it gives the same
 country more than once in different periods of time. Thus it approaches the
 special value of a historical atlas. . . . The *General Atlas* is wonderfully full
 and accurate for its scale. . . . Finally, the *Astronomical Atlas*, in which Mr
 Hind is responsible for the scientific accuracy of the maps, supplies an admitted
 educational want. No better companion to an elementary astronomical treatise
 could be found than this cheap and convenient collection of maps."—*Saturday
 Review*.

"The plan of these Atlases is admirable, and the excellence of the plan is rivalled
 by the beauty of the execution. . . . The best security for the accuracy
 and substantial value of a School Atlas is to have it from the hands of a man like
 our Author, who has perfected his skill by the execution of much larger works,
 and gained a character which he will be careful not to jeopardise by attaching
 his name to anything that is crude, slovenly, or superficial."—*Scotsman*.

"Mr Johnston, whose labours in this department of science have won him
 great renown, has, by the publication before us (The School Physical Atlas), done
 much to facilitate and encourage the study of physical geography. . . . It
 consists of eighteen beautiful illustrations, exhibiting the leading natural character-
 istics of every portion of the globe. . . . Nothing could be better calcu-
 lated to illustrate the various elementary treatises on physical geography which
 have lately appeared."—*Athenaeum*.

Hand Atlases:

Being the Maps of KEITH JOHNSTON'S School Atlases, on Large Paper, and half-bound, full size, Imperial Quarto.

PHYSICAL GEOGRAPHY,

ILLUSTRATING, IN A SERIES OF ORIGINAL DESIGNS, THE ELEMENTARY FACTS OF GEOLOGY, HYDROLOGY, METEOROLOGY, AND NATURAL HISTORY.

In Imperial Quarto, half-bound morocco, price 21s. In Portfolio, with each Map separate, mounted on Canvass, price 16s. 6d.

CLASSICAL GEOGRAPHY,

COMPRISING, IN TWENTY PLATES, MAPS AND PLANS OF ALL THE IMPORTANT COUNTRIES AND LOCALITIES REFERRED TO BY CLASSICAL AUTHORS.

In Imperial Quarto, half-bound morocco, price 21s. In Portfolio, with each Map separate, mounted on Canvass, price 16s. 6d.

New and Enlarged Edition.

GENERAL AND DESCRIPTIVE GEOGRAPHY,

EXHIBITING THE ACTUAL AND COMPARATIVE EXTENT OF ALL THE COUNTRIES IN THE WORLD, WITH THEIR PRESENT POLITICAL DIVISIONS.

In Imperial Quarto, half-bound morocco, price 21s. In Portfolio, with each Map separate, mounted on Canvass, price 16s. 6d.

ASTRONOMY,

COMPRISING, IN EIGHTEEN PLATES, A COMPLETE SERIES OF ILLUSTRATIONS OF THE HEAVENLY BODIES, DRAWN WITH THE GREATEST CARE FROM ORIGINAL AND AUTHENTIC DOCUMENTS.

By ALEX. KEITH JOHNSTON, F.R.S.E. &c.

Edited by J. R. HIND, F.R.A.S. &c.

In Imperial Quarto, half-morocco, price £1, 1s.

"To say that Mr Hind's Atlas is the best thing of the kind is not enough—it has no competitor."—*Athenæum*.

"The Atlas is undoubtedly the most beautiful work of its class that has ever been published, and in several respects the most instructive."—*The Astronomer Royal*.

"I am delighted with the *Atlas of Astronomy* which has just arrived; . . . it is exceedingly valuable, both as a complete system of the science and as a work of art of the most perfect and beautiful kind; indeed all your works, so eminently useful and admirable, do honour to our country."—*Mrs Somerville*.

COMPANION TO KEITH JOHNSTON'S ATLASES.

MANUAL OF MODERN GEOGRAPHY,

MATHEMATICAL, PHYSICAL, AND POLITICAL.

Embracing a Complete Development of the River-Systems of the Globe.

By the Rev. ALEXANDER MACKAY, F.R.G.S.

Crown 8vo, with a very complete Index, pp. 760, price 7s. 6d. bound.

OPINIONS.

Annual Address of the President of the Royal Geographical Society (Sir Roderick I. Murchison), 27th May 1861.

We must admire the ability and persevering research with which he has succeeded in imparting to his Manual so much freshness and originality. In no respect is this character more apparent than in the plan of arrangement by which the author commences his description of the physical geography of each tract by a sketch of its true basis or geological structure. The work is largely sold in Scotland, but has not been sufficiently spoken of in England. It is, indeed, a most useful school-book in opening out geographical knowledge.

English Journal of Education.

Of all the Manuals on Geography that have come under our notice, we place the one whose title is given above in the first rank. For fulness of information, for knowledge of method in arrangement, for the manner in which the details are handled, we know of no work that can, in these respects, compete with Mr Mackay's Manual.

Saturday Review.

Mr Mackay's Manual is meant to accompany the School General Atlas. It contains a prodigious array of geographical facts, and will be found useful for purposes of reference.

Spectator.

In the two essential points of completeness and compactness, this is by far the best geographical Manual with which we are acquainted. . . . Mr Mackay's Manual fully deserves, and we heartily hope that it will meet with, a success commensurate with the vast amount of time and labour which he must have expended on its compilation.

Daily News.

This volume combines fulness with compactness in an extraordinary degree. It is eminently practical in its arrangement, while ample care is taken throughout to place the facts in their scientific relation. . . . The volume will be found very useful either in the school or the counting-house.

Critic.

This is perhaps the most elaborate and carefully-written text-book of geography which has ever come under our notice. D'Anville himself would, we think, were he living, give it his meed of warm approbation.

Educational Guardian.

A valuable companion and worthy addition to the series of Atlases by Keith Johnston, issued from the same establishment. . . . Pupil-teachers, students in training-colleges, &c., will find it both interesting and valuable. They have here all they can, as students, ever require; and, as a text-book, it may be recommended as being as far in advance of Cornwell's and Sullivan's treatises as these were superior to the catechisms and grammars which they have superseded.

Athenæum.

The plan upon which it is composed is good, great stress being laid upon physical geography, and the river-systems of the globe being completely developed in a manner never before attempted.

W. BLACKWOOD & SONS, EDINBURGH AND LONDON.



•

1

1

1

1

1

1

1

1

1

1

1

1

1

